

# L<sup>A</sup>T<sub>E</sub>X TEMPLATE

XIAOXIANG ZHOU

## CONTENTS

1. Degree of a Map Induced by Divisors with Weighted Coefficients	1
References	2

### 1. DEGREE OF A MAP INDUCED BY DIVISORS WITH WEIGHTED COEFFICIENTS

Let  $C$  be a smooth projective curve over  $\mathbb{C}$  of genus 4, and view  $C$  as a subvariety of its Jacobian  $A := \text{Jac}(C)$  via the Abel–Jacobi map. Consider the map

$$f : C \times C \times C \longrightarrow A \quad (p_1, p_2, p_3) \longmapsto 2p_1 + 3p_2 + 4p_3.$$

What is the degree of  $f$  onto its image? In other words, for a general point  $a \in \text{Im}(f)$ , can we prove that  $\#f^{-1}(a) = 1$ ?

This question is related to Brill–Noether theory, but I have not found an answer to this specific case, which concerns divisors with multiplicities. Insights from the tropical version might provide helpful ideas.

## REFERENCES

- [1] Jens Niklas Eberhardt.  $K$ -motives and Koszul duality. *Bulletin of the London Mathematical Society*, 54(6):2232–2253, 2022.
- [2] Ravi Vakil. The rising sea: Foundations of algebraic geometry. *preprint*, 2017.

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, BERLIN, 12489, GERMANY,  
*Email address:* `email:xiaoxiang.zhou@hu-berlin.de`