

Master thesis

Contents

1	Variety and stratification	5
1.1	Initial case: \mathcal{F} and $\mathcal{F} \times \mathcal{F}$	5
1.2	quiver and Weyl group	7
1.3	algebraic group and Lie algebra	7
1.4	typical variety	7
1.5	stratification and T -fixed points	7
2	K-theory and cohomology theory	9
3	Localization theorem	11
4	Excess intersection formula	13
5	From formula to diagram	15
6	Generalization	17
7	Atiyah-Segal completion theorem	19

Chapter 1

Variety and stratification

1.1 Initial case: \mathcal{F} and $\mathcal{F} \times \mathcal{F}$

We introduce the complete flag variety to give a bird's eye view on the whole section. Actually, the entire difficulty is bundled in this example.

Fix $n \geq 1$, we denote $\mathrm{GL}_n := \mathrm{GL}_n(\mathbb{C})$, B , T , N , W be the standard Borel subgroup, standard torus, unipotent subgroup, Weyl group respectively, i.e.,

$$\mathrm{GL}_n = \begin{pmatrix} * & \dots & * \\ \vdots & \ddots & \vdots \\ * & \dots & * \end{pmatrix} \quad B = \begin{pmatrix} * & \dots & * \\ & \ddots & \vdots \\ 0 & & * \end{pmatrix} \quad T = \begin{pmatrix} * & & 0 \\ & \ddots & \\ 0 & & * \end{pmatrix} \quad B = \begin{pmatrix} 1 & \dots & * \\ & \ddots & \vdots \\ 0 & & 1 \end{pmatrix}$$

$$W := N_{\mathrm{GL}_n}(T)/T \cong S_n$$

Definition 1.1.1 (flag). *For a finite dimensional \mathbb{C} -vector space V , a flag of V is an increasing sequence of subspaces of V :*

$$F : 0 \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq V_k = \mathbb{C}^n.$$

F is called a complete flag if $\dim V_i = i$ for all i , otherwise F is called a partial flag.

Definition 1.1.2 (complete flag variety). *The complete flag variety \mathcal{F} is defined as*

$$\begin{aligned} \mathcal{F} &= \mathrm{GL}_n / B \\ &\cong \{\text{complete flags of } \mathbb{C}^n\} \\ &= \{0 \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq V_n = \mathbb{C}^n \mid \dim V_i = i\} \\ &\cong \{\text{Borel subgroups of } \mathrm{GL}_n\} \\ &= \{gBg^{-1} \mid g \in \mathrm{GL}_n\} \end{aligned}$$

Remark 1.1.3.

1. We implicitly give the base point of \mathcal{F} , which is not considered as the data of \mathcal{F} . Fix a standard basis of \mathbb{C}^n by $\{v_1, \dots, v_n\}$, we define the standard flag

$$F_{\text{Id}} : 0 \subseteq \langle v_1 \rangle \subseteq \langle v_1, v_2 \rangle \subseteq \dots \subseteq \langle v_1, \dots, v_n \rangle = \mathbb{C}^n.$$

2. We have the natural GL_n -action on \mathcal{F} , which is considered as the data of \mathcal{F} . For $g \in \text{GL}_n$, we define the flag attached to g :

$$F_g \triangleq gF_{\text{Id}} : 0 \subseteq \langle gv_1 \rangle \subseteq \langle gv_1, gv_2 \rangle \subseteq \dots \subseteq \langle gv_1, \dots, gv_n \rangle = \mathbb{C}^n.$$

Especially, for $w \in W = N_{\text{GL}_n}(T)/T \cong S_n$, the flag attached to w

$$\begin{aligned} F_w : 0 \subseteq \langle \tilde{w}v_1 \rangle \subseteq \langle \tilde{w}v_1, \tilde{w}v_2 \rangle \subseteq \dots \subseteq \langle \tilde{w}v_1, \dots, \tilde{w}v_n \rangle &= \mathbb{C}^n \\ 0 \subseteq \langle v_{w(1)} \rangle \subseteq \langle v_{w(1)}, v_{w(2)} \rangle \subseteq \dots \subseteq \langle v_{w(1)}, \dots, v_{w(n)} \rangle &= \mathbb{C}^n \end{aligned}$$

does not depend on the choice of the lift $\tilde{w} \in N_{\text{GL}_n}(T)$ of w .

Readers can verify that $\{F_w | w \in W\}$ are all T -fixed points of \mathcal{F} , while $\{wBw^{-1} | w \in W\}$ are all Borel subgroups of G containing the standard torus T .

3. For $n = 2$, $\mathcal{F} \cong \mathbb{P}^1$. We encourage readers to use \mathbb{P}^1 as a toy example for the whole theory.

interpretation	GL_n/B	flags	Borel subgroups
base point	Id	F_{Id}	B
GL_n -action	left multiplication	$\{V_i\} \mapsto \{gV_i\}$	conjugation
general point	g	F_g	gBg^{-1}

\mathcal{F} is a well-studied variety, and has many combinatorial properties. For example, from the well-known Bruhat decomposition,¹

$$\text{GL}_n \cong \bigsqcup_{w \in W} BwB$$

We get a stratification of \mathcal{F} by B -orbits:

$$\mathcal{F} = \text{GL}_n/B \cong \bigsqcup_{w \in W} BwB/B$$

The B -orbit BwB/B is called the Schubert cell, denoted by \mathcal{V}_w . Since

$$\mathcal{V}_w = BwB/B \cong B/(B \cap wBw^{-1}) \cong \mathbb{A}^{l(w)},$$

the Schubert cell is an affine space of dimension $l(w)$.

As a result, we know a lot of information of \mathcal{F} :

¹For the most time the formula does not depend on the lift of w , so we abuse the notation of $w \in N_{\text{GL}_n}(T)/T$ and $\tilde{w} \in N_{\text{GL}_n}(T)$.

$H^i(\mathcal{F}; \mathbb{C})$	0	2	4	6	8	10	12
1	1						
2	1	1					
3	1	2	2	1			
4	1	3	5	6	5	3	1
5	1	4	9	15	20	22	20

G	Orbit	G -fixed points
GL_n	$\mathcal{F} \cong GL_n / B$	\emptyset
B	$\mathcal{V}_w \cong B / (B \cap wBw^{-1})$	$\{F_{\text{Id}}\}$
T	–	$\{F_w \mid w \in W\}$

- 1.2 quiver and Weyl group
- 1.3 algebraic group and Lie algebra
- 1.4 typical variety
- 1.5 stratification and T -fixed points

Chapter 2

K-theory and cohomology theory

Chapter 3

Localization theorem

Chapter 4

Excess intersection formula

Chapter 5

From formula to diagram

Chapter 6

Generalization

Chapter 7

Atiyah-Segal completion theorem