

## Part I

# *K*-theory of the Steinberg varieties



## Part II

# Affine pavings of partial flag varieties



## Chapter 7

# Preliminaries for Affine Pavings

The goal of the two chapters is to find an affine paving of the partial flag variety  $\text{Flag}_{\mathbf{f}}(X)$ , which generalizes the variety  $\text{Flag}_{\mathbf{d}}(X)$  defined in 1.5.9. We have constructed some stratifications of most varieties by orbits in Subsection 1.6.1 and 1.6.2, and some are even affine pavings. For  $\text{Flag}_{\mathbf{f}}(X)$ , we have no natural group actions, so we can not find affine pavings by methods in Subsection 1.6.1 and 1.6.2. To achieve our goal, we extend methods from [12] and [16] as well as techniques from Auslander–Reiten theory.

In this chapter, we realize the partial flag variety  $\text{Flag}_{\mathbf{f}}(X)$  as a quiver Grassmannian of the extended quiver, prove some Ext-vanishing properties, and also give a short introduction of Auslander–Reiten theory.

**Setting 7.0.1.** *Throughout Chapter 7 and 8,  $R$  is a  $\mathbb{C}$ -algebra with unit, and  $\text{mod}(R)$  denotes the category of  $R$ -modules of finite dimension. For a representation  $X \in \text{rep}(Q)$ , we denote by  $X_i := e_i X$  the  $\mathbb{C}$ -linear space at the vertex  $i \in v(Q)$ . We denote by  $P(i)$ ,  $I(i)$  and  $S(i)$  the indecomposable projective, injective, simple modules corresponding to the vertex  $i$ , respectively.*

### 7.1 Extended quiver

In this section, we introduce the notion of extended quiver which allows to view partial flag varieties as quiver Grassmannians. Intuitively, a flag of quiver representations can be encoded as a subspace of a representation of the extended quiver.

**Definition 7.1.1** (Extended quiver). *For a quiver  $Q$  and an integer  $d \geq 1$ , the extended quiver  $Q_d$  is defined as follows:*

- *The vertex set of  $Q_d$  is defined as the Cartesian product of the vertex set of  $Q$  and  $\{1, \dots, d\}$ , i.e.,*

$$v(Q_d) = v(Q) \times \{1, \dots, d\}.$$

- *There are two types of arrows: for each  $(i, r) \in v(Q) \times \{1, \dots, d-1\}$ , there is one arrow from  $(i, r)$  to  $(i, r+1)$ ; for each arrow  $i \rightarrow j$  in  $Q$  and  $r \in \{1, \dots, d\}$ , there is one arrow from  $(i, r)$  to  $(j, r)$ .*

The extended quiver  $Q_d$  is exactly the same quiver as  $\hat{\Gamma}_d$  in [16, Definition 2.2]. The next definition is a small variation:


**Definition 7.1.2** (Strict extended quiver). *For a quiver  $Q$  and an integer  $d \geq 2$ , the strict extended quiver  $Q_{d,\text{str}}$  is defined as follows:*

- The vertex set of  $Q_d$  is defined as the Cartesian product of the vertex set of  $Q$  and  $\{1, \dots, d\}$ , i.e.,

$$v(Q_{d,\text{str}}) = v(Q) \times \{1, \dots, d\}.$$

- We have two types of arrows: for each  $(i, r) \in v(Q) \times \{1, \dots, d-1\}$ , there is one arrow from  $(i, r)$  to  $(i, r+1)$ ; for each arrow  $i \rightarrow j$  in quiver  $Q$  and  $r \in \{2, \dots, d\}$ , there is one arrow from  $(i, r)$  to  $(j, r-1)$ .

**Example 7.1.3.** The (strict) extended quiver for a Dynkin quiver  $Q$  of type  $A_4$  looks as follows.



Next, we define the quiver algebras for later use.

**Definition 7.1.4** (Algebra of an extended quiver). For an extended quiver  $Q_d$ , let  $\mathbb{C}Q_d$  be the corresponding path algebra, and  $I$  be the ideal of  $\mathbb{C}Q_d$  identifying all the paths with the same sources and targets. The algebra of the extended quiver  $Q_d$  is defined as

$$R_d := \mathbb{C}Q_d/I.$$

Similarly, we define the algebra  $R_{d,\text{str}} := \mathbb{C}Q_{d,\text{str}}/I$  for the strict extended quiver.

By abuse of notation, we often abbreviate  $R_d$  and  $R_{d,\text{str}}$  by  $R$ .

## 7.2 Canonical functor $\Phi$

We follow [16, 2.3] in this section with a few variations.

**Definition 7.2.1** (Partial flag variety). For a quiver representation  $X \in \text{rep}(Q)$  and an integer  $d \geq 1$ , we define

$$\text{Flag}_d(X) := \{0 \subseteq M_1 \subseteq \dots \subseteq M_d \subseteq X\}$$

as the collection of all partial flags of length  $d$ , and call it the partial flag variety associated to  $X$ .

**Definition 7.2.2** (Strict partial flag variety). For a quiver representation  $X \in \text{rep}(Q)$  and an integer  $d \geq 2$ , we define

$$\text{Flag}_{d,\text{str}}(X) := \{0 \subseteq M_1 \subseteq \dots \subseteq M_d \subseteq X \mid x.M_{k+1} \subseteq M_k\}$$

as the collection of all strict partial flags of length  $d$ , and call it the strict partial flag variety associated to  $X$ .

**Definition 7.2.3** (Grassmannian). Let  $R$  be the bounded quiver algebra defined in Definition 7.2.1 or 7.2.2. For a module  $T \in \text{mod}(R)$ , the Grassmannian  $\text{Gr}^R(T)$  is defined as the set of all submodules of  $T$ , i.e.,

$$\text{Gr}^R(T) := \{T' \subseteq T \text{ as the submodule}\}.$$

**Definition 7.2.4** (Canonical functor  $\Phi$ ). *The canonical functor  $\Phi : \text{rep}(Q) \longrightarrow \text{mod}(R)$  is defined as follows:*

- $(\Phi(X))_{(i,r)} := X_i$ ;
- $(\Phi(X))_{(i,r) \rightarrow (i,r+1)} := \text{Id}_{X_i}$ ;
- *Either*  $(\Phi(X))_{(i,r) \rightarrow (j,r)} := X_{i \rightarrow j}$  *for*  $R = R_d$ ,  
*or*  $(\Phi(X))_{(i,r) \rightarrow (j,r-1)} := X_{i \rightarrow j}$  *for*  $R = R_{d,\text{str}}$ .

The functor  $\Phi$  helps to realize a partial flag as a quiver subrepresentation.

**Proposition 7.2.5.** *For a representation  $X \in \text{rep}(Q)$ , the canonical functor  $\Phi$  induces isomorphisms*

$$\text{Flag}_d(X) \cong \text{Gr}^{R_d}(\Phi(X)) \quad \text{Flag}_{d,\text{str}}(X) \cong \text{Gr}^{R_{d,\text{str}}}(\Phi(X)).$$

*Proof.* The isomorphism maps a flag  $M : M_1 \subseteq \cdots \subseteq M_d$  to a representation  $\Phi'(M)$  with  $\Phi'(M)_{(i,r)} = M_{i,r}$  and obvious morphisms for arrows. The non-strict case is mentioned in [16, page 4] and the strict case works similarly.  $\square$

**Example 7.2.6.** *Consider the quiver  $Q : x \longrightarrow y \longleftarrow z \longrightarrow w$ , and let  $X : X_x \longrightarrow X_y \longleftarrow X_z \longrightarrow X_w$  be a representation. The varieties  $\text{Flag}_3(X), \text{Flag}_{3,\text{str}}(X)$  then arise as quiver Grassmannian as shown in Figure 7.1.*

In many cases, the proof of the strict case and the non-strict case is the same, so we often treat them in the same way. For example, we may abbreviate the formula in Proposition 7.2.5 as

$$\text{Flag}(X) \cong \text{Gr}(\Phi(X)).$$

### 7.3 Dimension vector

In this section we define some varieties indexed by dimension vectors. Recall Definition 1.2.7 and 1.2.11. Notice that every module  $T \in \text{mod}(\mathbb{C}Q/I)$  can be viewed as a representation of  $Q$ , so we automatically have a notion of dimension vector for  $T$ .

We can write the (strict) partial flag variety and Grassmannian as disjoint union of several pieces. Since  $v(Q_{d,(\text{str})}) = v(Q) \times \{1, \dots, d\}$ , any dimension vector  $\underline{\mathbf{f}}$  of  $R$  can be viewed as  $d$  dimension vectors  $(\underline{\mathbf{f}}_1, \dots, \underline{\mathbf{f}}_d)$ . Define

$$\begin{aligned} \text{Flag}_{\underline{\mathbf{f}}}(X) &:= \{0 \subseteq M_1 \subseteq \cdots \subseteq M_d \subseteq X \mid \underline{\dim} M_k = \underline{\mathbf{f}}_k\} && \subseteq \text{Flag}_d(X), \\ \text{Flag}_{\underline{\mathbf{f}},\text{str}}(X) &:= \{0 \subseteq M_1 \subseteq \cdots \subseteq M_d \subseteq X \mid x.M_{k+1} \subseteq M_k, \underline{\dim} M_k = \underline{\mathbf{f}}_k\} && \subseteq \text{Flag}_{d,\text{str}}(X), \\ \text{Gr}_{\underline{\mathbf{f}}}^R(T) &:= \{T' \subseteq T \text{ with } \underline{\dim} T' = \underline{\mathbf{f}}\} && \subseteq \text{Gr}^R(T). \end{aligned}$$

Then from the Proposition 7.2.5 we get

$$\text{Flag}_{\underline{\mathbf{f}}}(X) \cong \text{Gr}_{\underline{\mathbf{f}}}^{R_d}(\Phi(X)) \quad \text{Flag}_{\underline{\mathbf{f}},\text{str}}(X) \cong \text{Gr}_{\underline{\mathbf{f}}}^{R_{d,\text{str}}}(\Phi(X)).$$

Finally, we need to define the Euler form of two dimension vectors. For this we need to define the set of virtual arrows of the quivers  $Q_d$  and  $Q_{d,\text{str}}$ . Following Example 7.3.3, the virtual arrows of the quivers  $Q_3$  and  $Q_{3,\text{str}}$  are depicted in red.




Figure 7.1

**Definition 7.3.1** (Virtual arrows of the quiver  $Q_d$ ). For  $d \geq 1$ , the virtual arrows of the quiver  $Q_d$  is defined as a triple  $(va(Q_d), s, t)$ , where

$$va(Q_d) := a(Q) \times \{1, \dots, d-1\}$$

is a finite set, and  $s, t : va(Q_d) \rightarrow v(Q_d)$  are maps defined by

$$s((i \rightarrow j, r)) = (i, r) \quad t((i \rightarrow j, r)) = (j, r+1).$$

**Definition 7.3.2** (Virtual arrows of the quiver  $Q_{d, \text{str}}$ ). For  $d \geq 2$ , the virtual arrows of the quiver  $Q_{d, \text{str}}$  is defined as a triple  $(va(Q_{d, \text{str}}), s, t)$ , where


$$va(Q_{d, \text{str}}) := a(Q) \times \{2, \dots, d-1\}$$

is a finite set, and  $s, t : va(Q_{d, \text{str}}) \rightarrow v(Q_{d, \text{str}})$  are maps defined by

$$s((i \rightarrow j, r)) = (i, r) \quad t((i \rightarrow j, r)) = (j, r).$$

**Example 7.3.3.**





**Definition 7.3.4** (Euler form of  $R$ ). *Let  $R$  be a bounded quiver algebra defined in Definition 7.1.4. We denote*

$$\begin{aligned} v(R) &:= \{\text{vertices in } Q_d \text{ or } Q_{d,\text{str}}\}, \\ a(R) &:= \{\text{arrows in } Q_d \text{ or } Q_{d,\text{str}}\}, \\ va(R) &:= \{\text{virtual arrows in } Q_d \text{ or } Q_{d,\text{str}}\}. \end{aligned}$$

For two dimension vectors  $\underline{f}, \underline{g}$  of  $R$ , the Euler form  $\langle \underline{f}, \underline{g} \rangle_R$  is defined by

$$\langle \underline{f}, \underline{g} \rangle_R := \sum_{i \in v(R)} f_i g_i - \sum_{b \in a(R)} f_{s(b)} g_{t(b)} + \sum_{c \in va(R)} f_{s(c)} g_{t(c)}.$$

## 7.4 Ext-vanishing properties

We will show that some higher rank extension group are zero, which will be a key ingredient in the proofs of Section 8.1.

For a bounded quiver algebra  $R$  defined in Definition 7.1.4, we have a standard resolution for every  $R$ -module  $T$ :

$$\begin{aligned} 0 \rightarrow \bigoplus_{c \in va(Q)} Re_{t(c)} \otimes_{\mathbb{C}} e_{s(c)} T \rightarrow \bigoplus_{b \in a(Q)} Re_{t(b)} \otimes_{\mathbb{C}} e_{s(b)} T \rightarrow \bigoplus_{i \in v(Q)} Re_i \otimes_{\mathbb{C}} e_i T \rightarrow T \rightarrow 0 \\ r \otimes x \longmapsto \begin{matrix} rc_1 \otimes x + r \otimes b_1 x \\ -rc_2 \otimes x - r \otimes b_2 x \end{matrix} \qquad r \otimes x \longmapsto rx \\ r \otimes x \longmapsto rb \otimes x - r \otimes bx \end{aligned}$$

There are exactly two paths of length two from  $s(c)$  to  $t(c)$  for any virtual arrow  $c$ , which we denoted by  $b_1 c_1$  and  $b_2 c_2$  in the above. By definition, these paths are identified in  $R$ .

**Lemma 7.4.1.** *Let  $M, N \in \text{rep}(Q)$ .*

- (1)  $\text{gl. dim } R \leq 2$ ;
- (2) The functor  $\Phi : \text{rep}(Q) \rightarrow \text{mod}(R)$  is exact and fully faithful;
- (3)  $\Phi$  maps projective module to projective module, and maps injective module to injective module;
- (4)  $\text{Ext}_{\mathbb{C}Q}^i(M, N) \cong \text{Ext}_R^i(\Phi(M), \Phi(N))$ ;
- (5)  $\text{proj. dim } \Phi(M) \leq 1, \text{inj. dim } \Phi(M) \leq 1$ ;

*Proof.*

For (1), this follows from the standard resolution.

For (2), it follows by direct inspection, see [16, Lemma 2.3].

For (3), we reduce to the case of indecomposable projective modules, and observe that

$$\Phi(P(i)) = P((i, 1)), \quad \Phi(I(i)) = I((i, d)).$$

For (4), it comes from the fact that  $\Phi$  is fully faithful and maps projective module to projective module.

For (5), notice that the minimal projective resolution of  $M$  is of length 1, and  $\Phi(-)$  sends the projective resolution of  $M$  to the projective resolution of  $\Phi(M)$  by (3), thus we get  $\text{proj. dim } \Phi(M) \leq 1$ . The injective dimension of  $\Phi(M)$  is computed in a similar way.  $\square$

The following key lemma will be crucial later.

**Lemma 7.4.2.** *Let  $X, S \in \text{rep}(Q)$  and  $V \subseteq \Phi(X), W \subseteq \Phi(S), T \in \text{mod}(R)$ . Then  $\text{Ext}_R^2(W, T) = 0$  and  $\text{Ext}_R^2(T, \Phi(X)/V) = 0$ .*

*Proof.* The short exact sequence

$$0 \longrightarrow W \longrightarrow \Phi(S) \longrightarrow \Phi(S)/W \longrightarrow 0$$

induces the long exact sequence

$$\cdots \longrightarrow \text{Ext}_R^2(\Phi(S), T) \longrightarrow \text{Ext}_R^2(W, T) \longrightarrow \text{Ext}_R^3(\Phi(S)/W, T) \longrightarrow \cdots$$

By Lemma 7.4.1 (1) and (5),  $\text{Ext}_R^3(\Phi(S)/W, T)$  and  $\text{Ext}_R^2(\Phi(S), T)$  are both 0, so  $\text{Ext}_R^2(W, T) = 0$ .

Similarly, from the short exact sequence

$$0 \longrightarrow V \longrightarrow \Phi(X) \longrightarrow \Phi(X)/V \longrightarrow 0$$

we get the induced long exact sequence

$$\cdots \longrightarrow \text{Ext}_R^2(T, \Phi(X)) \longrightarrow \text{Ext}_R^2(T, \Phi(X)/V) \longrightarrow \text{Ext}_R^3(T, V) \longrightarrow \cdots,$$

so  $\text{Ext}_R^2(T, \Phi(X)/V) = 0$ .  $\square$

We will frequently use extension groups as well as long exact sequences, so we introduce some abbreviations. For  $Q$ -representations  $M, N$  and  $R$ -modules  $T, T'$ , we denote

$$\begin{aligned} [M, N]^i &:= \dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^i(M, N) & [M, N] &:= \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(M, N) \\ [T, T']^i &:= \dim_{\mathbb{C}} \text{Ext}_R^i(T, T') & [T, T'] &:= \dim_{\mathbb{C}} \text{Hom}_R(T, T') \end{aligned}$$

and write the Euler form as

$$\langle T, T' \rangle_R := \sum_{i=0}^{\infty} (-1)^i [T, T']^i = [T, T'] - [T, T']^1 + [T, T']^2.$$

**Lemma 7.4.3** (Homological interpretation of the Euler form). *For two  $R$ -modules  $T, T'$ , we have*

$$\langle T, T' \rangle_R = \langle \underline{\dim} T, \underline{\dim} T' \rangle_R.$$

*Proof.* Compute  $\langle T, T' \rangle_R$  by applying the functor  $\text{Hom}_R(-, T')$  to the standard resolution of the  $R$ -module  $T$ .  $\square$

## 7.5 A crash course on Auslander–Reiten theory

In this section, we will introduce concepts in Auslander–Reiten theory one by one: indecomposable representation, irreducible morphism, Auslander–Reiten translation, Auslander–Reiten sequence, Auslander–Reiten quiver, and minimal sectional mono. The main references for the material covered in this section are [4, 16].

**Definition 7.5.1** (Indecomposable module). *Fix an algebra  $R$ . A non-zero module  $M \in \text{mod}(R)$  is called indecomposable if  $M$  can not be written as a direct sum of two non-zero submodules. The set of all indecomposable modules is denoted by  $\text{ind}(R)$ .*

There are several descriptions of the indecomposable representations in special cases. For instance:

- By Gabriel’s theorem [11, Theorem 2.1], the functor **dim** yields a bijection from the indecomposable representations of a Dynkin quiver to the positive roots of the associated Lie algebra.

There is a unique indecomposable representation of maximal dimension vector which corresponds to the unique maximal positive root. This is shown in Table 7.1.

- By [4, Theorem 2, p34], in the affine case, the functor **dim** yields a surjective map from the indecomposable representations to the positive roots of the associated affine diagram. The map is  $\infty$ -to-1 when the root is imaginary, and is 1-to-1 when the root is real.<sup>1</sup>

We also have a unique minimal imaginary root  $\delta$  which controls the whole indecomposable representation theory, as shown in Table 7.1.

- All indecomposable representations of Dynkin quivers and all indecomposable representations of affine quivers corresponding to the positive real roots  $\alpha$  with  $\alpha < \delta$  or  $\langle \alpha, \delta \rangle \neq 0$  are rigid, i.e.,  $[M, M]^1 = 0$ . They are also bricks, i.e.,  $[M, M]^1 = 0$  and  $[M, M] = 1$ .<sup>2</sup>

Indecomposable representations form the vertices of Auslander–Reiten quiver, while irreducible morphisms form the arrows.

**Definition 7.5.2** (Irreducible morphism). *Given two indecomposable representations  $T, T' \in \text{mod}(R)$ , let*

$$\begin{aligned} \text{rad}(T, T') &:= \{f \in \text{Hom}_R(T, T') \mid f \text{ is not invertible}\} \\ &= \begin{cases} \text{Hom}_R(T, T') & T \not\cong T', \\ \text{Jac}(\text{End}_R(T)) & T \cong T'. \end{cases} \end{aligned}$$

*be the radical, and let*

$$\text{rad}^2(T, T') := \bigcup_{S \in \text{ind}(R)} \text{Im} [\text{rad}(T, S) \times \text{rad}(S, T') \longrightarrow \text{rad}(T, T')]$$

*be the subspace of  $\text{rad}(T, T')$ . A morphism  $f \in \text{Hom}_R(T, T')$  is called irreducible if  $f \in \text{rad}(T, T') \setminus \text{rad}^2(T, T')$ .*

The definition of irreducible morphism applies to any representation, and one can easily show that any irreducible morphism is either injective or surjective.

<sup>1</sup>The root  $\alpha \in \mathbf{dim}(Q)$  is called real if  $\langle \alpha, \alpha \rangle = 1$ , and called imaginary if  $\langle \alpha, \alpha \rangle = 0$ .

<sup>2</sup>Any rigid indecomposable module of a hereditary algebra is a brick.

Type	maximal positive real root(Dynkin)	minimal positive imaginary root $\delta$ (affine)
$A$	$1 - 1 - \dots - 1 - 1$	$\begin{array}{c} 1 \\ \swarrow \quad \searrow \\ 1 - 1 - \dots - 1 - 1 \end{array}$
$D$	$\begin{array}{c} 1 \\   \\ 1 - 1 - \dots - 2 - 1 \end{array}$	$\begin{array}{c} 1 \quad 1 \\   \quad   \\ 1 - 2 - \dots - 2 - 1 \end{array}$
$E_6$	$\begin{array}{c} 2 \\   \\ 1 - 2 - 3 - 2 - 1 \end{array}$	$\begin{array}{c} 1 - 2 \\   \\ 1 - 2 - 3 - 2 - 1 \end{array}$
$E_7$	$\begin{array}{c} 2 \\   \\ 1 - 2 - 3 - 4 - 3 - 2 \end{array}$	$\begin{array}{c} 2 \\   \\ 1 - 2 - 3 - 4 - 3 - 2 - 1 \end{array}$
$E_8$	$\begin{array}{c} 3 \\   \\ 2 - 3 - 4 - 5 - 6 - 4 - 2 \end{array}$	$\begin{array}{c} 3 \\   \\ 1 - 2 - 3 - 4 - 5 - 6 - 4 - 2 \end{array}$

Table 7.1: Roots which control all other roots.

**Definition 7.5.3.** Let  $R = \mathbb{C}Q/I$  be a bounded quiver algebra. We define the Nakayama functor  $\nu_R$ , Auslander–Reiten translation  $\tau_R$ , and inverse Auslander–Reiten translation  $\tau_R^{-1}$ , as follows:

$$\begin{aligned} \nu_R : \quad \text{mod}(R) &\xrightarrow{\text{Hom}_R(-, R^R)} \text{mod}(R^{op}) \xrightarrow{\text{Hom}_{\mathbb{C}}(-, \mathbb{C})} \text{mod}(R), \\ \tau_R : \quad \underline{\text{mod}}(R) &\xrightarrow{\text{Ext}_R^1(-, R^R)} \underline{\text{mod}}(R^{op}) \xrightarrow{\text{Hom}_{\mathbb{C}}(-, \mathbb{C})} \overline{\text{mod}}(R), \\ \tau_R^{-1} : \quad \overline{\text{mod}}(R) &\xrightarrow{\text{Hom}_{\mathbb{C}}(-, \mathbb{C})} \underline{\text{mod}}(R^{op}) \xrightarrow{\text{Ext}_{R^{op}}^1(-, R_R)} \underline{\text{mod}}(R). \end{aligned}$$

Here  $\underline{\text{mod}}(R)$  and  $\overline{\text{mod}}(R)$  denote the stable module categories. The objects are the same as in  $\text{mod}(R)$ , and their morphisms are modified by “collapsing” the morphisms passing through projective/injective modules to zero, i.e.,

$$\begin{aligned} \text{Mor}_{\underline{\text{mod}}(R)}(T, T') &:= \text{Mor}_{\text{mod}(R)}(T, T') / (f : T \rightarrow P \rightarrow T', P \text{ is projective}), \\ \text{Mor}_{\overline{\text{mod}}(R)}(T, T') &:= \text{Mor}_{\text{mod}(R)}(T, T') / (f : T \rightarrow I \rightarrow T', I \text{ is injective}). \end{aligned}$$

These modifications guarantee that the Auslander–Reiten translation  $\tau_R$  is indeed a functor. For convenience, we abbreviate  $\text{Mor}_{\underline{\text{mod}}(R)}$ ,  $\text{Mor}_{\overline{\text{mod}}(R)}$ ,  $\text{Mor}_{\text{mod}(R)}$  as  $\underline{\text{Hom}}_R$ ,  $\overline{\text{Hom}}_R$ ,  $\text{Hom}_R$ , and ignore the subscription  $R$  in the symbol  $\tau_R$ .

The Auslander–Reiten translation has many magical properties. For example,  $\tau_R$  induces the one-to-one correspondence between non-projective indecomposable representations and non-injective indecomposable representations. We would also frequently use the Auslander–Reiten formulas:  $((-)^{\vee} = \text{Hom}_{\mathbb{C}}(-, \mathbb{C})$  is the dual)

$$\begin{aligned} (\overline{\text{Hom}}_R(T, \tau T'))^{\vee} &\xrightarrow{\sim} \text{Ext}_R^1(T', T) \\ (\underline{\text{Hom}}_R(\tau^{-1} T, T'))^{\vee} &\xrightarrow{\sim} \text{Ext}_R^1(T', T) \end{aligned}$$

which is functorial for any  $T, T' \in \text{mod}(R)$ . Especially, when  $T$  is not injective,  $\overline{\text{Hom}}_R(T, \tau T') = \text{Hom}_R(T, \tau T')$ , we get  $[T', T]^1 = [T, \tau T']$ ; when  $T'$  is not projective,  $\underline{\text{Hom}}_R(\tau^{-1} T, T') = \text{Hom}_R(\tau^{-1} T, T')$ , we get  $[T', T]^1 = [\tau^{-1} T, T']$ .

For the Auslander–Reiten sequence there can be many equivalent definitions, and we only present one due to limitations of space.

**Definition 7.5.4** (Auslander–Reiten sequence). *For  $X \in \text{ind}(R)$  non-projective, an epimorphism  $g : E \rightarrow X$  is called **right almost split** if  $g$  is not split epi and every homomorphism  $h : T \rightarrow X$  which is not split epi factors through  $E$ . The short exact sequence*

$$0 \rightarrow \tau X \rightarrow E \xrightarrow{g} X \rightarrow 0$$

*is called an Auslander–Reiten sequence if  $g$  is right almost split.*

All the concepts introduced in this section can be clearly observed from the Auslander–Reiten quiver. In the Auslander–Reiten quiver the vertices are indecomposable representations, the arrows are irreducible morphisms among indecomposable representations, Auslander–Reiten translation is labeled as the dotted arrow, and the Auslander–Reiten sequence can be read by collecting all paths from  $\tau X$  to  $X$ . For instance, in Figure 7.2 we can get an Auslander–Reiten sequence

$$0 \rightarrow 12\bar{3}21 \rightarrow 12\bar{2}11 \oplus 11\bar{1}10 \oplus 01\bar{2}21 \rightarrow 12\bar{2}21 \rightarrow 0$$

of the corresponding quiver.

Finally we move forward to the definition of minimal sectional mono. The rest can be skipped until Lemma 8.3.1.

**Definition 7.5.5** (Sectional morphism). *Let  $Q$  be a quiver of Dynkin/affine type, and  $M, N \in \text{rep}(Q)$  be two indecomposable representations of  $Q$ , which are preprojective<sup>3</sup> when  $Q$  is affine. A morphism  $f \in \text{Hom}_{\mathbb{C}Q}(M, N)$  is called sectional if  $f$  can be written as the composition*

$$f : M = X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} \dots \xrightarrow{f_{t-1}} X_{t-1} \xrightarrow{f_t} X_t = N$$

*where  $f_i \in \text{Hom}_{\mathbb{C}Q}(X_{i-1}, X_i)$  are irreducible morphisms between indecomposable representations, and  $\tau X_{i+2} \not\cong X_i$  for any suitable  $i$ .*

**Remark 7.5.6.** Let  $f$  be a sectional morphism. If the underlying quiver  $Q$  is a Dynkin/affine quiver without oriented cycles, then  $X_0, \dots, X_t$  are uniquely determined, and  $f_1, \dots, f_t$  are unique up to constant.

**Lemma 7.5.7.** *Any sectional morphism  $f \in \text{Hom}_{\mathbb{C}Q}(M, N)$  is either surjective or injective.*

*Proof.* When  $Q$  is a quiver without oriented cycles, then  $[N, M]^1 \leq [M, \tau N] = 0$ , thus by [16, Lemma 7] we get the result; when  $Q$  is of type  $\tilde{A}$ , the result comes from [16, Lemma 51].  $\square$

**Definition 7.5.8** (Sectional mono, minimal sectional mono). *Let  $Q$  be a quiver without oriented cycles. A sectional morphism  $f \in \text{Hom}_{\mathbb{C}Q}(M, N)$  is called as a sectional mono if  $f$  is injective; a sectional mono is called minimal if  $f_t \circ \dots \circ f_{i+1} : X_i \rightarrow N$  are surjective for any  $i \in \{1, 2, \dots, t\}$ .*

---

<sup>3</sup>A representation  $M \in \text{rep}(Q)$  is called preprojective if  $\tau^k M$  is projective for some  $k \geq 0$ . Similarly, A representation  $M \in \text{rep}(Q)$  is called preinjective if  $\tau^{-k} M$  is injective for some  $k \geq 0$ .




Figure 7.2: The Auslander-Reiten quiver of the quiver

$$E_6 : \begin{array}{c} \bullet \\ \downarrow \\ \bullet \leftarrow \bullet \rightarrow \bullet \leftarrow \bullet \leftarrow \bullet \end{array}$$

Minimal sectional monos can also be clearly seen from the Auslander-Reiten quiver, and we can check if a sectional morphism is mono by comparing the dimension vectors. In the case of Example  $E_6$  in Figure 7.2, a non-zero morphism from  ${}^{00110}_1$  to  ${}^{11110}_1$  is a minimal sectional mono while a non-zero morphism from  ${}^{01100}_0$  to  ${}^{01211}_1$  is not, since a sectional morphism from  ${}^{01210}_1$  to  ${}^{01211}_1$  is also injective.

## Chapter 8

# Constructions of Affine Pavings

With tools in hand, we prove the main theorem 8.1.2 and 8.1.3 in Part II, and apply them to obtain the affine paving structure of partial flag varieties, see Theorem B.

### 8.1 Main theorem in Part II

In this section we state and prove the main theorems, which are essential in Section 8.2 and 8.4.

Let  $\eta : 0 \longrightarrow X \xrightarrow{\iota} Y \xrightarrow{\pi} S \longrightarrow 0$  be a short exact sequence in  $\text{rep}(Q)$ . Consider the canonical **non-continuous** map

$$\Psi : \text{Gr}(\Phi(Y)) \longrightarrow \text{Gr}(\Phi(X)) \times \text{Gr}(\Phi(S)) \quad U \longmapsto ([\Phi(\iota)]^{-1}(U), [\Phi(\pi)](U)).$$

Denote the set

$$\text{Gr}(\Phi(Y))_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} := \Psi^{-1}\left(\text{Gr}_{\underline{\mathbf{f}}}(\Phi(X)) \times \text{Gr}_{\underline{\mathbf{g}}}(\Phi(S))\right)$$

and let  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  be the map  $\Psi$  restricted to  $\text{Gr}(\Phi(Y))_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$ , i.e.,

$$\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} : \text{Gr}(\Phi(Y))_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} \longrightarrow \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X)) \times \text{Gr}_{\underline{\mathbf{g}}}(\Phi(S)).$$

*Remark 8.1.1.* Even though  $\Psi$  is not continuous,  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  is continuous. Moreover, for any dimension vectors  $\underline{\mathbf{f}}, \underline{\mathbf{g}}$ , the set

$$\text{Gr}(\Phi(Y))_{\geq \underline{\mathbf{f}}, \leq \underline{\mathbf{g}}} := \left\{ U \in \text{Gr}(\Phi(Y)) \mid \begin{array}{l} \underline{\dim}[\Phi(\iota)]^{-1}(U) \geq \underline{\mathbf{f}} \\ \underline{\dim}[\Phi(\pi)](U) \leq \underline{\mathbf{g}} \end{array} \right\}$$

is closed in  $\text{Gr}(\Phi(Y))$ . This gives us a filtration

$$0 = Z_0 \subset Z_1 \subset \cdots \subset Z_d = \text{Gr}_{\underline{\mathbf{h}}}(\Phi(Y))$$

with  $Z_i$  closed and  $Z_{i+1} \setminus Z_i$  isomorphic to  $\text{Gr}(\Phi(Y))_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  for some  $\underline{\mathbf{f}}, \underline{\mathbf{g}}$ . Therefore, from the affine pavings of  $\text{Gr}(\Phi(Y))_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  (for every  $\underline{\mathbf{f}}, \underline{\mathbf{g}}$ ) one can construct one affine paving of  $\text{Gr}_{\underline{\mathbf{h}}}(\Phi(Y))$ .

**Theorem 8.1.2.** *If  $\eta$  splits, then  $\Psi$  is surjective. Moreover, if  $[S, X]^1 = 0$ , then  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  is a Zarisky-locally trivial affine bundle of rank  $\langle \underline{\mathbf{g}}, \underline{\dim} \Phi(X) - \underline{\mathbf{f}} \rangle_R$ .*

**Theorem 8.1.3** (Generalizes [12, Theorem 32]). *When  $\eta$  does not split and  $[S, X]^1 = 1$ ,*

$$\text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} = \left( \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X)) \times \text{Gr}_{\underline{\mathbf{g}}}(\Phi(S)) \right) \setminus \left( \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X_S)) \times \text{Gr}_{\underline{\mathbf{g}} - \underline{\mathbf{dim}} \Phi(S^X)}(\Phi(S/S^X)) \right)$$

where

$$\begin{aligned} X_S &:= \max \{ M \subseteq X \mid [S, X/M]^1 = 1 \} \subseteq X, \\ S^X &:= \max \{ M \subseteq S \mid [M, X]^1 = 1 \} \subseteq S. \end{aligned}$$

Moreover,  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  is a Zarisky-locally trivial affine bundle of rank  $\langle \underline{\mathbf{g}}, \underline{\mathbf{dim}} \Phi(X) - \underline{\mathbf{f}} \rangle_R$  over  $\text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$ .

We will spend the rest of the section proving these theorems. We investigate the image as well as the fiber of  $\Psi$  respectively.

**Lemma 8.1.4** (Follows [12, Lemma 21]). *The element  $(V, W) \in \text{Gr}(\Phi(X)) \times \text{Gr}(\Phi(S))$  lies in the image of  $\Psi$  if and only if the canonical map  $\text{Ext}^1(\Phi(S), \Phi(X)) \longrightarrow \text{Ext}^1(W, \Phi(X)/V)$  maps  $\eta$  to 0.*

*Proof.* The canonical map is defined as follows:

$$\begin{array}{ccccccc} \eta \in \text{Ext}^1(\Phi(S), \Phi(X)) & 0 & \longrightarrow & \Phi(X) & \longrightarrow & \Phi(Y) & \xrightarrow{\Phi(\pi)} \Phi(S) \longrightarrow 0 \\ & & & \parallel & & \uparrow & \uparrow \\ & \text{Ext}^1(W, \Phi(X)) & 0 & \longrightarrow & \Phi(X) & \longrightarrow & \pi^{-1}(W) \longrightarrow W \longrightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \parallel \\ \bar{\eta} \in \text{Ext}^1(W, \Phi(X)/V) & 0 & \longrightarrow & \Phi(X)/V & \longrightarrow & \pi^{-1}(W)/V & \longrightarrow W \longrightarrow 0 \end{array}$$

so  $\bar{\eta} = 0$  if and only if the last short exact sequence splits, that means, there exists a submodule  $U \subseteq \Phi(Y)$ , such that  $\Phi(\pi)(U) = W$  and  $U \cap \Phi(X) = V$ .  $\square$

**Corollary 8.1.5.** *Resume the notations of Lemma 8.1.4 When  $\eta$  splits, then  $\Psi$  is surjective.*

**Lemma 8.1.6.** *The canonical map  $\text{Ext}^1(\Phi(S), \Phi(X)) \longrightarrow \text{Ext}^1(W, \Phi(X)/V)$  is surjective.*

*Proof.* By using the long exact sequence of extension groups and the fact that  $\text{Ext}^2(\Phi(S)/W, \Phi(X)) = 0$  and  $\text{Ext}^2(W, V) = 0$  by Lemma 7.4.2, the maps

$$\text{Ext}^1(\Phi(S), \Phi(X)) \longrightarrow \text{Ext}^1(W, \Phi(X)) \quad \text{Ext}^1(W, \Phi(X)) \longrightarrow \text{Ext}^1(W, \Phi(X)/V)$$

are both surjective. Thus the composition is also surjective.  $\square$

**Corollary 8.1.7.** *Let  $W \subseteq \Phi(S), V \subseteq \Phi(X)$  be  $R$ -submodules, then*

$$[W, \Phi(X)/V]^1 \leq [\Phi(S), \Phi(X)]^1 = [S, X]^1.$$

*In particular, when  $[S, X]^1 = 1$ , we get  $[W, \Phi(X)/V]^1 = 0$  or 1; when  $\eta$  generates  $\text{Ext}^1(S, X)$ , we get*

$$(V, W) \in \text{Im } \Psi \iff [W, \Phi(X)/V]^1 = 0.$$



In the case where  $\eta$  generates  $\text{Ext}^1(S, X)$ , we want to describe  $\text{Im } \Psi$  more precisely. For this reason we need to introduce two new  $R$ -modules:

$$\begin{aligned}\widetilde{X}_S &:= \max \{V \subseteq \Phi(X) \mid [\Phi(S), \Phi(X)/V]^1 = 1\} \subseteq \Phi(X), \\ \widetilde{S}^X &:= \max \{W \subseteq \Phi(S) \mid [W, \Phi(X)]^1 = 1\} \subseteq \Phi(S).\end{aligned}$$

$\widetilde{X}_S$  and  $\widetilde{S}^X$  are well-defined because of the following lemma:

**Lemma 8.1.8** (Follows [12, Lemma 27]).

- (i) Let  $V, V' \subset \Phi(X)$  such that  $[\Phi(S), \Phi(X)/V]^1 = [\Phi(S), \Phi(X)/V']^1 = 1$ . Then  $[\Phi(S), \Phi(X)/(V + V')]^1 = 1$ .
- (ii) Let  $W, W' \subset \Phi(S)$  such that  $[W, \Phi(X)]^1 = [W', \Phi(X)]^1 = 1$ . Then  $[W \cap W', \Phi(X)]^1 = 1$ .

*Proof.* We only prove (i). (ii) is similar.

From the short exact sequence

$$0 \longrightarrow \Phi(X)/(V \cap V') \longrightarrow \Phi(X)/V \oplus \Phi(X)/V' \longrightarrow \Phi(X)/(V + V') \longrightarrow 0,$$

we get the long exact sequence

$$\cdots \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V \cap V'}\right) \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V}\right) \oplus \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V'}\right) \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V + V'}\right) \rightarrow \cdots.$$

By Corollary 8.1.7,  $[\Phi(S), \Phi(X)/(V \cap V')]^1 \leq 1$ ,  $[\Phi(S), \Phi(X)/(V + V')]^1 \leq 1$ , and this forces  $[\Phi(S), \Phi(X)/(V + V')]^1 = 1$ .  $\square$

**Lemma 8.1.9** (Follows [12, Lemma 31(1)(2)], with the same proof). *Let  $\tau$  be the Auslander-Reiten translation.*

*Let  $f : X \rightarrow \tau S$  be a non-zero morphism,<sup>1</sup> then  $X_S = \ker(f)$ ;*

*also,  $\Phi(f) : \Phi(X) \rightarrow \Phi(\tau S)$  is a non-zero morphism,  $\widetilde{X}_S = \ker(\Phi(f))$ .*

*Proof.* For any  $M \subseteq X$ , we have

$$\begin{aligned}\text{Ext}^1(S, X/M)^\vee &\cong \overline{\text{Hom}}(X/M, \tau S) \\ &\cong \{g \in \text{Hom}(X, \tau S) \mid g|_M = 0\} \\ &\cong \begin{cases} \mathbb{C}, & M \subseteq \ker f \\ 0, & M \not\subseteq \ker f. \end{cases}\end{aligned}$$

so  $[S, X/M]^1 = 1$  exactly when  $M \subseteq \ker f$ . Thus  $X_S = \ker f$ .

For  $\Phi(f)$  it is similar. For any  $V \subseteq \Phi(X)$ , we have

$$\begin{aligned}\text{Ext}^1(\Phi(S), \Phi(X)/V)^\vee &\cong \overline{\text{Hom}}(\Phi(X)/V, \tau\Phi(S)) \\ &\cong \overline{\text{Hom}}(\Phi(X)/V, \Phi(\tau S)) \\ &\cong \{g \in \text{Hom}(\Phi(X), \Phi(\tau S)) \mid g|_V = 0\} \\ &\cong \begin{cases} \mathbb{C}, & V \subseteq \ker \Phi(f) \\ 0, & V \not\subseteq \ker \Phi(f). \end{cases}\end{aligned}$$

so  $[\Phi(S), \Phi(X)/V]^1 = 1$  exactly when  $V \subseteq \ker \Phi(f)$ . Thus  $\widetilde{X}_S = \ker(\Phi(f))$ .  $\square$

<sup>1</sup>Since  $X$  is not injective,  $[X, \tau S] = [S, X]^1 = 1$ ,  $f$  is uniquely determined up to a constant.

**Corollary 8.1.10.**  $\widetilde{X}_S = \Phi(X_S)$ . (since  $\widetilde{X}_S = \ker(\Phi(f)) = \Phi(\ker(f)) = \Phi(X_S)$ )

By a dual argument, one can show that  $\widetilde{S}^X = \Phi(S^X)$ .

**Lemma 8.1.11** (Follows [12, Lemma 31(6)]). For  $V \subseteq \Phi(X)$  and  $W \subseteq \Phi(S)$ , we have

$$[W, \Phi(X)/V]^1 = 0 \iff V \not\subseteq \Phi(X_S) \text{ or } W \not\supseteq \Phi(S^X).$$

*Proof.*  $\Leftarrow$ : Without loss of generality suppose  $V \not\subseteq \Phi(X_S)$ , then

$$V \not\subseteq \Phi(X_S) \iff [\Phi(S), \Phi(X)/V]^1 = 0 \Rightarrow [W, \Phi(X)/V]^1 = 0.$$

$\Rightarrow$ : If not, then  $V \subseteq \Phi(X_S)$  and  $W \supseteq \Phi(S^X)$ , and<sup>2</sup>

$$[W, \Phi(X)/V]^1 \geq [\Phi(S^X), \Phi(X)/\Phi(X_S)]^1 = [S^X, X/X_S]^1 = 1. \quad \square$$

**Corollary 8.1.12.** When  $\eta$  generates  $\text{Ext}^1(S, X)$ , we have

$$\text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} = \left( \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X)) \times \text{Gr}_{\underline{\mathbf{g}}}(\Phi(S)) \right) \setminus \left( \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X_S)) \times \text{Gr}_{\underline{\mathbf{g}} - \underline{\dim} \Phi(S^X)}(\Phi(S/S^X)) \right).$$

**Lemma 8.1.13.** For  $(V, W) \in \text{Im } \Psi$ , the preimage of  $(V, W)$  is a torsor of  $\text{Hom}_R(W, \Phi(X)/V)$ . Hence, there is a non-canonical isomorphism

$$\Psi^{-1}((V, W)) \cong \text{Hom}_R(W, \Phi(X)/V).$$

*Proof.* Recall the commutative diagram

$$\begin{array}{ccccccc} \eta \in \text{Ext}^1(\Phi(S), \Phi(X)) & 0 & \longrightarrow & \Phi(X) & \longrightarrow & \Phi(Y) & \xrightarrow{\Phi(\pi)} \Phi(S) \longrightarrow 0 \\ & & & \parallel & & \uparrow & \uparrow \\ & & & \text{Ext}^1(W, \Phi(X)) & 0 & \longrightarrow & \pi^{-1}(W) \longrightarrow W \longrightarrow 0 \\ & & & \downarrow & & \downarrow & \parallel \\ \bar{\eta} \in \text{Ext}^1(W, \Phi(X)/V) & 0 & \longrightarrow & \Phi(X)/V & \xrightarrow{\iota} & \pi^{-1}(W)/V & \xrightarrow{\pi'} W \longrightarrow 0 \end{array}$$

$\xleftarrow{\theta}$

When  $(V, W) \in \text{Im } \Psi$ ,  $\bar{\eta}$  is split, and each split morphism  $\theta$  give us an element in  $\Psi^{-1}((V, W))$ . If we fix one split morphism  $\theta_0$ , then the other split morphisms are all of the form  $\theta_0 + \iota \circ f$  where  $f \in \text{Hom}_R(W, \Phi(X)/V)$  (and this form is unique). So

$$\Psi^{-1}((V, W)) \cong \{\theta : \text{split morphism}\} \cong \text{Hom}_R(W, \Phi(X)/V). \quad \square$$

**Remark 8.1.14.** Any point  $(V, W) \in \text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  can be also viewed as a morphism

$$f : \text{Spec } \mathbb{C} \longrightarrow \text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} \subseteq \text{Gr}_{\underline{\mathbf{f}}}(\Phi(X)) \times \text{Gr}_{\underline{\mathbf{g}}}(\Phi(S))$$

where Grassmannian are viewed as moduli spaces over  $\mathbb{C}$ . Essentially by replacing  $\text{Spec } \mathbb{C}$  by any locally closed reduced subscheme  $\text{Spec } A$  of  $\text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  in Lemma 8.1.13, we can run the machinery of algebraic geometry, and mimic the proof of [12, Theorem 24] to show that  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  is a Zarisky-locally trivial affine bundle over  $\text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  when  $\eta$  generates  $\text{Ext}^1(S, X)$ .

*Proof of Theorem 8.1.2 and 8.1.3.* We have already computed  $\text{Im } \Psi$  in Corollary 8.1.5 and 8.1.12. In both cases  $\eta$  generates  $\text{Ext}^1(S, X)$ , so by Corollary 8.1.7 we get

$$\begin{aligned} (V, W) \in \text{Im } \Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}} &\iff [W, \Phi(X)/V]^1 = 0 \\ &\implies [W, \Phi(X)/V] = \langle W, \Phi(X)/V \rangle_R = \langle \underline{\mathbf{f}}, \underline{\dim} \Phi(X) - \underline{\mathbf{g}} \rangle_R. \end{aligned}$$

From Remark 8.1.14,  $\Psi_{\underline{\mathbf{f}}, \underline{\mathbf{g}}}$  is a Zarisky-locally trivial affine bundle.  $\square$

<sup>2</sup> $[S^X, X/X_S]^1 = 1$  follows from [12, Lemma 31(5)].

## 8.2 Application: Dynkin case

This section (plus Section 8.3) mainly focus on the proof of the following result:

**Theorem 8.2.1.** *For any Dynkin quiver  $Q$  and any representation  $M \in \text{rep}(Q)$ , the (strict) partial flag variety  $\text{Flag}(M) \cong \text{Gr}(\Phi(M))$  has an affine paving.*

Before discussing the proof of the affine paving property, we introduce some numerical concepts, which can be seen as a measure of the “complexity” of the representation.




Figure 8.1: The quantity  $\text{ord}_e$  for indecomposable representations in type  $E$  arranged in the Auslander–Reiten quiver <sup>3</sup>

For an **indecomposable** quiver representation  $M \in \text{rep}(Q)$ , we define the order of  $M$  by

$$\text{ord}(M) := \max_{i \in v(Q)} \dim_{\mathbb{C}} M_i.$$

When the quiver  $Q$  is of type  $E$ , we denote by  $e \in v(Q)$  the unique vertex which is connected to three other vertices, and the number

$$\text{ord}_e(M) := \dim_{\mathbb{C}} M_e = [P(e), M]$$

is equal to  $\text{ord}(M)$  unless  $\text{ord}_e(M) = 0$ .

The next lemma shows the affine paving property for representations of small order.

**Lemma 8.2.2** (Follows [16, Lemma 2.22]). *Suppose that the underlying graph of  $Q$  is a tree. For an indecomposable representation  $M \in \text{rep}(Q)$  with  $\text{ord}(M) \leq 2$ , the variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  is either empty or a direct product of some copies of  $\mathbb{P}^1$ . Especially, the partial flag variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  has an affine paving.*

*Proof.* For every  $i \in v(Q)$ ,  $\dim_{\mathbb{C}} M_i \leq 2$ . Since  $Q$  is a tree and  $M$  is indecomposable, for every  $b \in a(Q)$  satisfying  $\dim_{\mathbb{C}} M_{s(b)} = \dim_{\mathbb{C}} M_{t(b)} = 2$ , the map  $M_{s(b)} \rightarrow M_{t(b)}$  is an isomorphism. Therefore, when  $\text{Grf}(\Phi(M)) \neq \emptyset$ ,<sup>4</sup> we get the natural embedding

$$\mathrm{Gr}_{\underline{\mathbf{f}}}(\Phi(M)) \longrightarrow \prod_{\substack{i \in v(Q) \text{ s.t.} \\ \dim_{\mathbb{C}} M_i = 2 \\ \underline{\mathbf{f}}_{(i,r)} = 1 \text{ for some } r}} \mathbb{P}^1,$$

<sup>3</sup>Some representations  $M$  are hidden when  $\text{ord}_e(M) = 0$ . In [1] the Figure 8.1 is called the starting functions.

<sup>4</sup>This condition imposes very strong restrictions on  $\mathbf{f}$ .

and the information of non-vertical arrows in the extended quiver (see Example 7.1.3) just reduce the number of  $\mathbb{P}^1$ . Precisely, one need to carefully discuss three cases of  $M_i \rightarrow M_j$ :

$$K \hookrightarrow K^2 \quad K^2 \twoheadrightarrow K \quad \text{and} \quad K^2 \xrightarrow{\cong} K^2. \quad \square$$

Now we've nearly prepared every step of the proof of Theorem 8.2.1. By following the process in Figure 8.2, we now prove Theorem 8.2.1 assuming Claim 8.2.3. We will prove Claim 8.2.3 in Section 8.3.




Figure 8.2: The process of induction

**Claim 8.2.3.** *Suppose  $Q$  is of Dynkin type. For any indecomposable representation  $M \in \text{rep}(Q)$  with  $\text{ord}(M) > 2$ , the (strict) partial flag variety  $\text{Gr}(\Phi(M))$  has an affine paving.*

*Proof of Theorem 8.2.1.* First of all, for any indecomposable representation  $M \in \text{rep}(Q)$  we obtain an affine paving. This follows from Claim 8.2.3 when  $\text{ord}(M) > 2$ , and follows from Lemma 8.2.2 when  $\text{ord}(M) \leq 2$ .

The general case follows by induction on the dimension vector. The indecomposable representations  $\{N_i\}_{i \in v(Q)}$  of quiver  $Q$  can be ordered such that  $[N_i, N_j] = 0$  for all  $i > j$ . Therefore, every non-indecomposable representation  $M$  can be decomposed as the direct sum of two nonzero representations  $M_1, M_2$  satisfying  $[M_2, M_1]^1 = 0$ . By applying Theorem 8.1.2 to the short exact sequence

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0,$$

we get an affine paving from the affine pavings of  $M_1$  and  $M_2$ , see Remark 8.1.1.  $\square$

*Remark 8.2.4.* By the same technique one can show that, for Dynkin quiver  $Q$  and any representation  $M$  with  $\max_{i \in v(Q)} \dim_{\mathbb{C}} M_i \leq 2$ , the variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  has an affine paving.

This result does not depend on Claim 8.2.3.


### 8.3 Proof of Claim 8.2.3

The task of this section is to prove Claim 8.2.3. When the quiver  $Q$  is of type  $A$  or  $D$ , Claim 8.2.3 is trivially true since no indecomposable representation can have order bigger than two. So we only concentrate on type  $E$ .

The idea of the proof is as follows. For any indecomposable representation  $Y$  with  $\text{ord}(Y) > 2$ , we put  $Y$  into a short exact sequence

$$\eta : 0 \longrightarrow X \longrightarrow Y \longrightarrow S \longrightarrow 0$$

fulfilling the assumptions of Theorem 8.1.3, and then  $\text{Gr}(\Phi(Y))$  has an affine paving if  $\text{Im } \Psi$  has. If additionally the map  $X \hookrightarrow Y$  is a minimal sectional mono, then  $\text{Im } \Psi_{\underline{f}, \underline{g}}$  can be written as the product space, which makes  $\text{Im } \Psi$  easier to understand.



The next two lemmas tell us the existence of the desired short exact sequence.

**Lemma 8.3.1.** *For every indecomposable representation  $Y$  of type  $E$  with  $\text{ord}(Y) > 2$ , there is a minimal sectional mono  $f : X \longrightarrow Y$ .*

*Proof.* Just observe the Auslander–Reiten quiver. The chosen minimal sectional monos are represented in Figure 8.3. Notice that for the most time  $\text{ord}_e(-)$  is enough to guarantee the map to be a mono.  $\square$




Figure 8.3: Minimal sectional monos

**Remark 8.3.2.** The condition  $\text{ord}(Y) > 2$  in the lemma can not be removed.

**Lemma 8.3.3.** *Let  $X \hookrightarrow Y$  be a minimal sectional mono, and  $S := Y/X$  be the quotient. Then we have the short exact sequence*

$$\eta : 0 \longrightarrow X \longrightarrow Y \longrightarrow S \longrightarrow 0$$

$\begin{array}{c} [M, N] \\ [M, N]^1 \\ M \end{array}$	$N$			
	X	Y	S	
X	1 0	1 0	0 0	
Y	0 0	1 0	1 0	
S	0 1	0 0	1 0	

Table 8.1

and the dimensions of extension groups among  $X, Y, S$  are as shown in the Table 8.1.

In particular,  $S$  is indecomposable and rigid;  $[S, X]^1 = 1$ , so  $X_S$  and  $S^X$  are well-defined.

*Proof.* Since every indecomposable representation of Dynkin quiver is a brick, we get  $[X, X] = [Y, Y] = 1$  and  $[X, X]^1 = [Y, Y]^1 = 0$ . By the definition of minimal sectional mono, we get  $[X, Y] = 1, [Y, X] = 0$  and  $[X, Y]^1 = [Y, X]^1 = 0$ . By applying the functors  $[Y, -], [-, S], [X, -], [-, X], [-Y]$  to the short exact sequence  $\eta$  we get the results.  $\square$

In the following two lemmas we will describe the representations  $S^X$  and  $X_S$  more clearly.

**Lemma 8.3.4.** *Take the same notations as in Lemma 8.3.3. Then  $S^X = S$ .*

*Proof.* Let  $\iota : N \rightarrow S$  be a proper non-zero subrepresentation of  $S$ , we need to prove that  $\iota^*\eta : 0 \rightarrow X \rightarrow Y' \rightarrow N \rightarrow 0$  splits.

$$\begin{array}{ccccccc}
 \iota^*\eta : & 0 & \longrightarrow & X & \hookrightarrow & Y' & \longrightarrow & N & \longrightarrow & 0 \\
 & & & \parallel & & \downarrow h & & \downarrow \iota & & \\
 \eta : & 0 & \longrightarrow & X & \longrightarrow & Y & \longrightarrow & S & \longrightarrow & 0
 \end{array}$$

We decompose  $Y' = \oplus_i Y'_i$  as the direct sum of indecomposable representations. Since the map  $X \rightarrow Y$  is the minimal sectional mono, we get  $Y'_i = X$  or  $Y'_i = Y$  or  $X \xrightarrow{0} Y'_i$  for each  $i$ . If there exists  $i$  such that  $Y'_i = X$ , then  $\iota^*\eta$  splits; if there exists  $i$  such that  $Y'_i = Y$ , then  $h$  is isomorphism, we get  $\iota$  is isomorphism; if for every  $i$  the map  $X \rightarrow Y'_i$  is 0, then the map  $X \rightarrow Y'$  is 0, we also get the contradiction.  $\square$


**Lemma 8.3.5** (Follows [12, Lemma 36], with the same proof). *Let  $E \rightarrow X$  be the minimal right almost split morphism ending in  $X$ , then we can decompose  $E$  as  $E = E' \oplus \tau X_1$ .<sup>5</sup> When  $Y$  is not projective,  $X_S$  is isomorphic to  $\ker(E \rightarrow \tau Y) \cong E' \oplus \ker(\tau X_1 \rightarrow \tau Y)$ ; when  $Y$  is projective,  $X_S \cong E$ .*

**Corollary 8.3.6.** *When  $X \rightarrow Y$  is irreducible monomorphism, the representation  $X_S$  is either 0 or an indecomposable representation with property that  $X_S \rightarrow X$  is also an irreducible monomorphism.*

<sup>5</sup>  $X_1$  is defined in Definition 7.5.5, where we take  $M = X$  and  $N = Y$ .

*Remark 8.3.7.* We can not copy everything in [12, Lemma 56], sometimes it would happen that  $X_S = F \oplus T$  with  $F$  and  $T$  indecomposable,  $F \hookrightarrow X$  is irreducible but  $T \rightarrow X/F$  is not a sectional mono. Even when  $X_S$  is indecomposable, the map  $X_S \rightarrow X$  can be not a sectional mono.

For example, take the quiver of type  $E_7$ :



take  $Y = {}_{111111}^0$ ,  $X = {}_{111110}^0$ , then  $X_S = {}_{111100}^0$ , the map  $X_S \rightarrow X$  is not a sectional mono.

Luckily, we can avoid this bad situation by carefully choosing the minimal sectional mono  $X \rightarrow Y$ . The minimal sectional monos I chose are presented in Figure 8.3. We will write down the induction process in detail for some examples.

For convenience, we simplify the notations: write  $\text{Gr}(M)$  as  $\text{Gr}_{\underline{f}}(\Phi(M))$ ,  $U(M, N)$  as  $\text{Gr}_{\underline{f}}(\Phi(M)) \setminus \text{Gr}_{\underline{f}}(\Phi(N))$ , where we omit subscripts which indicate the dimension vectors.

**Lemma 8.3.8** (Follows [12, Theorem 59]). *Let  $f : X \hookrightarrow Y$  be a minimal sectional mono and  $S := Y/X$  be the quotient. When  $X_S = F \oplus T$  with  $F$  and  $T$  indecomposable,  $F \hookrightarrow X$  irreducible and  $T \hookrightarrow X/F$  sectional mono, we have*

$$U(X, X_S) \longrightarrow \text{Gr}(F) \times U(X/F, T) \quad \text{or} \quad U(F, F_{X/F})$$

as a Zarisky-locally trivial affine bundle.

*Proof.* We have two short exact sequence satisfying the conditions in 8.1.3:

$$\begin{array}{lcl} \eta : & 0 \longrightarrow F \longrightarrow X \xrightarrow{\pi} X/F \longrightarrow 0 \\ \xi : & 0 \longrightarrow T \longrightarrow X/F \xrightarrow{\pi'} X/X_S \longrightarrow 0. \end{array}$$

Let  $N \in \text{Gr}(X)$  be a subrepresentation, it is obvious that  $N \in \text{Gr}(X_S) \iff \pi' \circ \pi(N) = 0$ , so

$$\begin{aligned} N \in U(X, X_S) &\iff \pi' \circ \pi(N) \neq 0 \\ &\iff \pi(N) \notin \text{Gr}(T) \\ &\iff \pi(N) \in U(X/F, T) \\ &\iff \Psi_\eta(N) \in \text{Gr}(F) \times U(X/F, T). \end{aligned}$$

Thus the Zarisky-locally trivial affine bundle map

$$U(X, F) \longrightarrow \text{Gr}(F) \times \text{Gr}(X/F)$$

restricted to the Zarisky-locally trivial affine bundle map

$$U(X, X_S) \longrightarrow \text{Gr}(F) \times U(X/F, T).$$

□

We conclude Theorem 8.1.3, Lemma 8.1.9, 8.3.3, 8.3.5, 8.3.8 in the following proposition.

**Proposition 8.3.9.** *Let  $f : X \hookrightarrow Y$  be a minimal sectional mono and  $S := Y/X$  be the quotient. One get Zariski-locally affine maps*

$$\begin{aligned} \mathrm{Gr}(Y) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(S) \quad \text{or} \quad U(X, X_S), \\ U(Y, X) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(S) \quad \text{or} \quad U(X, X_S), \end{aligned}$$

where

$$\begin{aligned} X_S &:= \max \{M \subseteq X \mid [S, X/M]^1 = 1\} \\ &= \ker(X \longrightarrow \tau S) \\ &= \begin{cases} E' \oplus \ker(\tau X_1 \longrightarrow \tau Y) & Y \text{ not projective,} \\ E & Y \text{ projective.} \end{cases} \end{aligned}$$

Moreover, when  $X_S = F \oplus T$  with  $F$  and  $T$  indecomposable,  $F \hookrightarrow X$  irreducible and  $T \hookrightarrow X/F$  sectional mono, one get a Zariski-locally affine map

$$U(X, X_S) \longrightarrow \mathrm{Gr}(F) \times U(X/F, T) \quad \text{or} \quad U(F, F_{X/F}).$$

With Proposition 8.3.9 in hand, we prove Claim 8.2.3 by cases, which turns out to be a pure combinatorial problem.

*Proof of Claim 8.2.3.* When the minimal sectional mono  $X \longrightarrow Y$  is irreducible, we use Theorem 8.1.3 to get morphism

$$\mathrm{Gr}(Y) \longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(S) \quad \text{or} \quad U(X, X_S).$$

By observation of Figure 8.3,  $\mathrm{ord}_e(S) = \mathrm{ord}_e(Y) - \mathrm{ord}_e(X)$  is smaller or equal to 2, so by Lemma 8.2.2  $\mathrm{Gr}(S)$  has the affine paving property. Let  $Y_1 := X$ ,  $X_1 := X_S$ ,  $S_1 := Y_1/X_1$ , we again use Theorem 8.1.3 to get Zariski-locally affine maps

$$\begin{aligned} \mathrm{Gr}(X) &\longrightarrow \mathrm{Gr}(X_1) \times \mathrm{Gr}(S_1) \quad \text{or} \quad U(X_1, X_{1S_1}) \\ U(X, X_S) &\longrightarrow \mathrm{Gr}(X_1) \times \mathrm{Gr}(S_1) \quad \text{or} \quad U(X_1, X_{1S_1}). \end{aligned}$$

Luckily  $\mathrm{ord}_e(S_1)$  is still smaller or equal to 2. We can continue this process until the order of representations are small enough.

The exceptional cases are similar, but the discussion is a bit more complicated. Let us look at some examples.

**Example 8.3.10.** *In the case of Figure 8.4(a), if  $X_1 \longrightarrow Y$  is injective, then we obtain some Zariski-locally affine maps*

$$\begin{aligned} \mathrm{Gr}(Y) &\longrightarrow \mathrm{Gr}(X_1) \times \mathrm{Gr}(Y/X_1) \quad \text{or} \quad U(X_1, X) \\ \mathrm{Gr}(X_1) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(X_1/X) \quad \text{or} \quad U(X, X_S) \\ U(X_1, X) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(X_1/X) \\ U(X, X_S) &\longrightarrow \mathrm{Gr}(X_S) \times \mathrm{Gr}(X/X_S). \end{aligned}$$

When  $X_1 \longrightarrow Y$  is not injective, we get

$$\mathrm{Gr}(Y) \longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(Y/X) \quad \text{or} \quad U(X, X_S).$$

Since the map  $\tau X_1 \longrightarrow \tau Y$  is injective, from Lemma 8.3.5 we get  $X_S \longrightarrow X$  is irreducible monomorphism. Thus

$$U(X, X_S) \longrightarrow \mathrm{Gr}(X_S) \times \mathrm{Gr}(X/X_S).$$

These maps give the variety  $\mathrm{Gr}(Y)$  an affine paving from bottom to top.






Figure 8.4: Special cases

**Example 8.3.11.** In Figure 8.4(b), we would like to prove that  $\text{Gr}(Y)$  has the affine paving property. We have

$$\text{Gr}(Y) \longrightarrow \text{Gr}(X) \times \text{Gr}(Y/X) \quad \text{or} \quad U(X, X_S).$$

When the map  $M \longrightarrow X$  is not monomorphism, we get

$$U(X, X_S) \longrightarrow \text{Gr}(X_S) \times \text{Gr}(X/X_S);$$

when the map  $M \longrightarrow X$  is monomorphism, we get

$$\begin{aligned} U(X, X_S) &= U(X, M) \bigsqcup U(M, X_S) \\ U(X, M) &\longrightarrow \text{Gr}(M) \times \text{Gr}(X/M) \\ U(M, X_S) &\longrightarrow \text{Gr}(X_S) \times \text{Gr}(M/X_S). \end{aligned}$$

Since the order of  $X$ ,  $Y/X$ ,  $X_S$ ,  $X/X_S$ ,  $M$ ,  $X/M$ ,  $M/X_S$  are smaller or equal to 2, the induction process stops, we get  $\text{Gr}(Y)$  has the affine paving property.

**Example 8.3.12.** In the case of Figure 8.4(c), we have

$$\text{Gr}(Y) \longrightarrow \text{Gr}(X) \times \text{Gr}(Y/X) \quad \text{or} \quad U(X, X_S)$$

where  $X_S = \ker(\tau X_1 \rightarrow \tau Y)$ . When  $X_S = 0$  we're done; if not, then  $A \neq 0$  and  $X_S = A$ , we decompose  $X_S \rightarrow Y$  as compositions of minimal sectional monos:

Case 1:  $M \rightarrow X$  is not injective, then

$$\begin{aligned} U(X, X_S) &= U(X, N) \bigsqcup U(N, X_S) \\ U(X, N) &\rightarrow \mathrm{Gr}(N) \times \mathrm{Gr}(X/N) \\ U(N, X_S) &\rightarrow \mathrm{Gr}(X_S) \times \mathrm{Gr}(N/X_S). \end{aligned}$$

Case 2:  $M \rightarrow X$  is injective, then

$$\begin{aligned} U(X, X_S) &= U(X, M) \bigsqcup U(M, N) \bigsqcup U(N, X_S) \\ U(X, M) &\rightarrow \mathrm{Gr}(M) \times \mathrm{Gr}(X/M) \\ U(M, N) &\rightarrow \mathrm{Gr}(N) \times \mathrm{Gr}(M/N) \\ U(N, X_S) &\rightarrow \mathrm{Gr}(X_S) \times \mathrm{Gr}(N/X_S). \end{aligned}$$

Since  $\mathrm{Gr}(X)$ ,  $\mathrm{Gr}(Y/X)$ ,  $\mathrm{Gr}(N)$ ,  $\dots$  have affine paving property, we conclude that  $\mathrm{Gr}(Y)$  has also the affine paving property.

**Example 8.3.13.** Finally we begin to tackle the most difficult case (Figure 8.4(d)). When  $X \rightarrow Y$  is not injective, we get

$$\mathrm{Gr}(Y) \rightarrow \mathrm{Gr}(F) \times \mathrm{Gr}(Y/F) \quad \text{or} \quad U(F, ?),$$

and then we get the result.<sup>6</sup>

When  $X \rightarrow Y$  is injective, we have

$$\mathrm{Gr}(Y) \rightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(Y/X) \quad \text{or} \quad U(X, X_S)$$

where  $X_S = F \oplus \ker(\tau X_1 \rightarrow \tau Y) = F \oplus T$  by Lemma 8.3.5. Since  $X \rightarrow Y$  is injective, we get  $A = 0$ , thus  $B = 0$  also, and then the sectional map  $T \rightarrow X/F$  is injective. We thus get two short exact sequence satisfying the conditions in 8.1.3:

$$\begin{aligned} \eta : \quad 0 &\longrightarrow F \longrightarrow X \xrightarrow{\pi} X/F \longrightarrow 0 \\ \xi : \quad 0 &\longrightarrow T \longrightarrow X/F \xrightarrow{\pi'} X/X_S \longrightarrow 0. \end{aligned}$$

Let  $N \in \mathrm{Gr}(X)$  be a subrepresentation, it is obvious that  $N \in \mathrm{Gr}(X_S) \iff \pi' \circ \pi(N) = 0$ , so

$$\begin{aligned} N \in U(X, X_S) &\iff \pi' \circ \pi(N) \neq 0 \\ &\iff \pi(N) \notin \mathrm{Gr}(T) \\ &\iff \pi(N) \in U(X/F, T) \\ &\iff \Psi_\eta(N) \in \mathrm{Gr}(F) \times U(X/F, T). \end{aligned}$$

Thus the Zarisky-locally trivial affine bundle map

$$U(X, F) \rightarrow \mathrm{Gr}(F) \times \mathrm{Gr}(X/F)$$

restricted to the Zarisky-locally trivial affine bundle map

$$U(X, X_S) \rightarrow \mathrm{Gr}(F) \times U(X/F, T).$$

---

<sup>6</sup> $\mathrm{Gr}(F)$  is empty or a singleton, so is  $U(F, ?)$ , no matter what representation is in the questionmark.

Finally, by applying the short exact sequence  $\xi$  to Theorem 8.1.3, we get the map

$$U(X/F, T) \longrightarrow \mathrm{Gr}(X/F) \times \mathrm{Gr}(T).$$

Since all the Grassmannians  $\mathrm{Gr}(X)$ ,  $\mathrm{Gr}(Y/X)$ ,  $\mathrm{Gr}(F)$ ,  $\mathrm{Gr}(X/F)$ ,  $\mathrm{Gr}(T)$  have the affine paving property, we conclude that  $\mathrm{Gr}(Y)$  has the affine paving property. □

## 8.4 Application: affine case

This section tries to explain the difficulty of the Conjecture 8.4.1.

**Conjecture 8.4.1.** *For any affine quiver  $Q$  and any indecomposable representation  $M \in \mathrm{rep}(Q)$ , the (strict) partial flag variety  $\mathrm{Flag}(M) \cong \mathrm{Gr}(\Phi(M))$  has an affine paving.*

Actually, if readers follow the proof in [12, Section 6], and change everything from  $\mathrm{Gr}(-)$  to  $\mathrm{Gr}(\Phi(-))$ , then there is no difference except the Proposition 48, in which the authors proved the affine paving properties of quasi-simple regular representations. So we reduced the question to the case of quasi-simple regular representation. Combined with Lemma 8.4.2, we've proved the affine paving properties for  $\tilde{A}$ ,  $\tilde{D}$  cases.

**Lemma 8.4.2.** *Assume that  $Q$  is an affine quiver of type  $A$  or  $D$ ,  $M \in \mathrm{rep}(Q)$  is the **regular quasi-simple** representation, then the Grassmannian  $\mathrm{Gr}(\Phi(M))$  has an affine paving.*

*Proof.* The concept “quasi-simple” is defined in [12, Definition 15]; the concepts “preprojective”, “preinjective” and “regular” are defined in [12, 2.1.1]. It's shown in [4, Section 9, Lemma 3] that the regular quasi-simple representation  $M$  have dimension vector smaller or equal to the minimal positive imaginary root, thus  $\mathrm{ord}_e(M) \leq 2$  for the quiver of type  $\tilde{D}$  and  $\mathrm{ord}_e(M) \leq 1$  for the quiver of type  $\tilde{A}$ . □

**Theorem 8.4.3.**

- (1) Assume that  $Q$  is an affine quiver of type  $A$  or  $D$ , then for any indecomposable representation  $M$ , the Grassmannian  $\mathrm{Gr}(\Phi(M))$  has an affine paving;
- (2) Assume that  $Q$  is an affine quiver of type  $E$ , and  $\mathrm{Gr}(\Phi(N))$  has an affine paving for any regular quasi-simple representation  $N \in \mathrm{rep}(Q)$ . The Grassmannian  $\mathrm{Gr}(\Phi(M))$  then has an affine paving for any indecomposable representation  $M$ .

For a regular quasi-simple representation  $Y$  of type  $\tilde{E}$ , it's possible that there's no short exact sequence

$$\eta : 0 \longrightarrow X \longrightarrow Y \longrightarrow S \longrightarrow 0$$

such that  $[S, X]^1 \leq 1$ . Then we can no longer use Theorem 8.1.2 or 8.1.3. Hence, the new methods are needed for this case.

# List of Tables

1	Current progress in affine pavings . . . . .	5
1.1	Orbit and fixed points of $\mathcal{F}$ and $\mathcal{F} \times \mathcal{F}$ . . . . .	12
1.2	Notations of the symmetric group calculus . . . . .	18
1.3	Different expressions of symmetric groups . . . . .	18
1.4	Lie algebras in different cases . . . . .	21
1.5	Base varieties and their preferred base point . . . . .	23
1.6	Stratifications of $\mathcal{F} \times \mathcal{F}$ . . . . .	27
1.7	Stratifications of $\mathcal{F}_{\mathbf{d}} \times \mathcal{F}_{\mathbf{d}}$ . . . . .	27
1.8	Stratifications of typical varieties . . . . .	29
3.1	Module structure of $K$ -theory: initial case . . . . .	47
3.2	Module structure of $K$ -theory: relative case . . . . .	47
3.3	Module structure of $K$ -theory: absolute case . . . . .	47
5.1	Lie algebras and corresponding Euler classes, $G_{\mathbf{d}}$ -action . . . . .	65
6.1	Lie algebras and corresponding Euler classes, $G_{\mathbf{d}} \times \mathbb{C}^{\times}$ -action . . . . .	68
7.1	Roots which control all other roots. . . . .	84
8.1	Dimension of Extension groups . . . . .	94

# List of Figures

5.1	The process of convolution product . . . . .	58
6.1	Generators of the $K$ -theory $K^{G_{\mathbf{d}}}(\mathcal{Z}_{\mathbf{d}})$ . . . . .	69
6.2	Relations of $K$ -theoretic KLR algebra, I . . . . .	71
6.3	Relations of $K$ -theoretic KLR algebra, II . . . . .	72
7.1	Realize flag as quiver Grassmannian . . . . .	80
7.2	The Auslander–Reiten quiver of type $E_6$ . . . . .	86
8.1	The quantity $\text{ord}_e$ for indecomposable representations in type $E$ . . . . .	91
8.2	The proof of affine paving in Dynkin case . . . . .	92
8.3	The chosen minimal sectional monos for complicated indecomposable representations . . . . .	93
8.4	A detailed discussion of the exceptional cases . . . . .	97

# Index

- affine paving, 5
- algebraic group
  - absolute algebraic group, 19
  - relative algebraic group, 19
- Atiyah–Segal completion theorem, 74
- augmentation ideal, 73
- Auslander–Reiten theory, 83
- Auslander–Reiten translation, 84
- Auslander–Reiten sequence, 85
- balanced product, 41
- canonical functor, 79
- cellular decomposition, 46
- cellular fibration structure, 46
- cellular fibration theorem, 46
- Characteristic class
  - Chern class, 74
  - Todd class, 74
- characteristic class
  - Euler class, 51
- Chevalley restriction theorem
  - for  $K$ -theory, 37
  - in cohomology theory, 44
- complete flag variety, 9
- convolution product
  - of  $K$ -theory, 58
  - of varieties, 57
- Demazure operator, 63
- diagram of strands, 69
- dimension vector
  - of  $Q$ -vector space, 14
  - of flag-type, 15
  - total dimension vector, 14
- equivariant cohomology, 43
- equivariant  $K$ -theory, 34
- equivariant sheaf, 33
- Euler form, 81
  - homological interpretation, 82
- excess base change formula, 52
- excess intersection formula, 59
- extended quiver, 77
- flag
  - of quiver representation, 14
  - of vector space, 9
  - standard flag, 22
- flag variety
  - absolute complete flag variety, 21
  - partial flag variety, 78
  - relative complete flag variety, 22
- Gabriel’s theorem, 83
- glueing theorem, 45
- Grassmannian, 78
- incidence variety, 23
  - for strict flag, 67
- indecomposable module, 83
- induction isomorphism, 42
- intersection product, *see* tensor product with supports
- irreducible morphism, 83
- KLR algebra, 70
- localization
  - fake localization formula, 53
  - localization formula, 53
  - localization theorem, 53
- minimal imaginary root, 83
- Nakayama functor, 84
- NilHecke algebra, 70
- order of representation, 91
- Poincaré–Hopf theorem, 51
- preinjective representation, 85
- preprojective representation, 85
- projection formula, 59

- $Q$ -vector space, 14
- quasi-simple representation, 98
- quiver, 13
- quiver representation, 13
- reduction isomorphism, 42
- $\text{Rep}_{\mathbf{d}}(Q)$ , 14
- $\text{rep}(Q)$ , 13
- representation ring, 34
- restriction with supports, 39
- right almost split, 85
- Schubert cell, 10
- Schubert variety, 11
- sectional morphism, 85
  - minimal sectional mono, 85
  - sectional mono, 85
- simple reflection, 16
- space of representations, *see*  $\text{Rep}_{\mathbf{d}}(Q)$
- stable module category, 84
- starting function, *see* order of representation
- Steinberg variety, 24
- stratification
  - of flag variety, 25
  - of incidence variety, 28
- tangent space
  - of  $T$ -fixed points, 31
- tensor product
  - external tensor product, 38
  - with supports, 40
- Thom isomorphism, 41
- twisted action, 25
- vector space with quiver partition, *see*  $Q$ -vector space
- virtual arrow, 80

# Bibliography

- [1] Klaus Bongartz. Critical simply connected algebras. *manuscripta mathematica*, 46(1):117–136, 1984.
- [2] Justin Brown. Flag varieties: An interplay of geometry, combinatorics, and representation theory. 2018.
- [3] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Boston, MA: Birkhäuser, reprint of the 1997 original edition, 2010.
- [4] William Crawley-Boevey. Lectures on representations of quivers. *unpublished notes*, 1992.
- [5] C. de Concini, George Lusztig, and C. Procesi. Homology of the zero-set of a nilpotent vector field on a flag manifold. *J. Am. Math. Soc.*, 1(1):15–34, 1988.
- [6] Daniel Dugger. A geometric introduction to  $K$ -theory. Available to download at <https://pages.uoregon.edu/ddugger/kgeom.pdf>, 2014.
- [7] Jens Niklas Eberhardt.  $K$ -motives and Koszul duality. *Bulletin of the London Mathematical Society*, 54(6):2232–2253, 2022.
- [8] Jens Niklas Eberhardt.  $K$ -theory Soergel bimodules, 2022.
- [9] Jens Niklas Eberhardt and Catharina Stroppel. Motivic Springer theory. *Indag. Math., New Ser.*, 33(1):190–217, 2022.
- [10] Dan Edidin and William Graham. Localization in equivariant intersection theory and the Bott residue formula. *Am. J. Math.*, 120(3):619–636, 1998.
- [11] D. Faenzi. A one-day tour of representations and invariants of quivers. *Rend. Semin. Mat., Univ. Politec. Torino*, 71(1):3–34, 2013.
- [12] Giovanni Cerulli Irelli, Francesco Esposito, Hans Franzen, and Markus Reineke. Cell decompositions and algebraicity of cohomology for quiver Grassmannians. *Adv. Math.*, 379:48, 2021. Id/No 107544.
- [13] David Kazhdan and George Lusztig. A topological approach to Springer’s representations. *Adv. Math.*, 38:222–228, 1980.
- [14] Mikhail Khovanov and Aaron D. Lauda. A diagrammatic approach to categorification of quantum groups. I. *Represent. Theory*, 13:309–347, 2009.
- [15] George Lusztig. Equivariant  $K$ -theory and representations of Hecke algebras. *Proc. Am. Math. Soc.*, 94:337–342, 1985.

- [16] Ruslan Maksimau. Flag versions of quiver Grassmannians for Dynkin quivers have no odd cohomology over  $\mathbb{Z}$ , 2019.
- [17] Tomasz Przezdziecki. Geometric approach to KLR algebras and their representation theory, 2015.
- [18] Raphael Rouquier. 2-Kac-Moody algebras, 2008.
- [19] Julia Sauter. Cell decompositions of quiver flag varieties for nilpotent representations of the oriented cycle. *arXiv preprint arXiv:1509.08026*, 2015.
- [20] Florian Seifarth. On KLR and quiver Schur algebras, 2017.
- [21] Robert Steinberg. On the desingularization of the unipotent variety. *Invent. Math.*, 36:209–224, 1976.
- [22] Catharina Stroppel and Ben Webster. Quiver Schur algebras and  $q$ -Fock space, 2011.
- [23] Robert W Thomason. Les  $K$ -groupes d’un schéma éclaté et une formule d’intersection excédentaire. *Inventiones mathematicae*, 112(1):195–215, 1993.
- [24] Ravi Vakil. The rising sea: Foundations of algebraic geometry. *preprint*, 2017.
- [25] M. Varagnolo and E. Vasserot. Canonical bases and KLR-algebras. *J. Reine Angew. Math.*, 659:67–100, 2011.
- [26] Xiaoxiang Zhou. Affine pavings of quiver flag varieties. *arXiv preprint arXiv:2206.00444*, 2022.