

Eine Woche, ein Beispiel

3.23: Schubert calculus: Chern class over Grassmannian

This is a follow up of [2025.02.23], [2025.03.16].

1. Formulas for tautological bundle
2. Homology class in $Gr(r,n)$

1. Formulas for tautological bundle

Chern class realized as pullback of σ_1 s

Prop. For those v.bs on $Gr(r, n)$, the Chern class is given by

$$\begin{aligned} c(\mathcal{S}) &= 1 - \sigma_1 + \dots + (-1)^r \sigma_r \\ c(\mathcal{Q}) &= 1 + \sigma_1 + \dots + \sigma_k + \dots + \sigma_{n-r} \\ c(\mathcal{S}^\vee) &= 1 + \sigma_1 + \dots + \sigma_r \\ c(\mathcal{Q}^\vee) &= 1 - \sigma_1 + \dots + (-1)^k \sigma_k + \dots + (-1)^{n-k} \sigma_{n-r} \end{aligned}$$

We omit the proof, as there are many equiv definition of Chern class, and I don't know which one to choose.

Cor If $f: X \rightarrow Gr(r, n)$ is induced by $(\mathcal{F}, s_1, \dots, s_n) = (\mathcal{O}_X^{\oplus n} \rightarrow \mathcal{F})$, then

$$\begin{aligned} c_s(\mathcal{F}) &= f^* c_s(\mathcal{S}^\vee) \\ &= f^* \sigma_1 \\ &= f^* \sum_1^s (\mathcal{V}^{st}) \\ &= f^* \{ \Delta \subset Gr(r, n) \mid \Delta + \mathcal{V}_{n-r+s-1}^{st} \subseteq H \} \\ &= \{ p \in X \mid (\mathcal{F}|_p)^* + \langle e_1^*, \dots, e_{n-r+s-1}^* \rangle \subseteq \mathcal{K}^{n-1} \} \\ &= \left\{ p \in X \mid \begin{array}{l} \exists (0, \dots, 0, k_{n-r+s}, \dots, k_n) \in \mathcal{K}^n - \{0\}, \text{ s.t.} \\ k_{n-r+s} s_{n-r+s}(p) + \dots + k_n s_n(p) = 0 \end{array} \right\} \\ &= \{ p \in X \mid \underbrace{s_{n-r+s}(p), \dots, s_n(p)}_{r-s+1 \text{ many}} \text{ are linear dependent} \} \end{aligned}$$

Especially,

$$c_r(\mathcal{F}) = \{ p \in X \mid s_n(p) = 0 \}$$

$$c_1(\mathcal{F}) = \{ p \in X \mid \underbrace{s_{n-r+1}(p), \dots, s_n(p)}_{r \text{ many}} \text{ are linear dependent} \}$$

$$= c_1(\Delta^r \mathcal{F})$$

$$= c_1(\det \mathcal{F})$$

Rmk. $c_s(\mathcal{F}) \neq c_{\text{top}}(\Delta^{r-s+1} \mathcal{F})$ since

$s_1 \wedge s_2$ (pure wedge) is not a general section in $\Delta^2 \mathcal{F}$!

Nevertheless, when $s=1$ or r , pure wedge is a general section, so
 $c_1(\mathcal{F}) = c_1(\det \mathcal{F})$ $c_r(\mathcal{F}) = c_r(\mathcal{F})$.

Porteous' formula

Thm [3264, Thm 12.4]

Let X/\mathbb{C} sm $k \in \mathbb{Z}_{\geq 0}$,
 \mathcal{E}, \mathcal{F} : v.b. over X of rank e, f ,
 $\varphi: \mathcal{E} \rightarrow \mathcal{F}$ map of v.b. (fiberwise linear).

$$M_k(\varphi) := \{x \in X \mid \text{rank } \varphi_x \leq k\}$$

remember multiplicity
 $\varphi_x: \mathcal{E}|_x \rightarrow \mathcal{F}|_x$

If $M_k(\varphi) \subset X$ has codim $(e-k)(f-k)$, then

$$[M_k(\varphi)] = \Delta_{f-k}^{e-k} \left[\frac{c(\mathcal{F})}{c(\mathcal{E})} \right] = (-1)^{(e-k)(f-k)} \Delta_{e-k}^{f-k} \left[\frac{c(\mathcal{E})}{c(\mathcal{F})} \right]$$

where

$$\Delta_{f-k}^{e-k}(\gamma) = \begin{vmatrix} \gamma_{f-k} & \cdots & \gamma_{e+f-2k-1} \\ \vdots & \ddots & \vdots \\ \gamma_{f-e+1} & \cdots & \gamma_{f-k} \end{vmatrix}_{(e-k-1) \times (e-k-1)}$$

2. Homology class in $Gr(r,n)$

Lines passing planes

E.g. 1. [3264, p131, Question(a)]

For 4 general lines l_1, l_2, l_3, l_4 in \mathbb{P}^3 , there are 2 lines meet all four.

Reason: In $Gr(2,4)$,

$$\begin{aligned} & \# \{l \in Gr(2,4) \mid l \cap l_i \neq \emptyset, \forall i\} \\ &= \deg \sigma_{\square}^4 \\ &= 2 \end{aligned}$$

E.g. 2. For 3 general lines l_1, l_2, l_3 in \mathbb{P}^4 , there is 1 line meet all three.

Reason: In $Gr(2,5)$,

$$\begin{aligned} & \# \{l \in Gr(2,5) \mid l \cap l_i \neq \emptyset, \forall i\} \\ &= \deg \sigma_{\square}^3 \\ &= 1 \end{aligned}$$

One can get further that no line in \mathbb{P}^5 passing 3 general lines.

E.g. 3.

For 6 general planes e_1, \dots, e_6 in \mathbb{P}^5 , there are 5 lines passing all these planes.

Reason: In $Gr(2,5)$,

$$\begin{aligned} & \# \{l \in Gr(2,5) \mid l \cap e_i \neq \emptyset, \forall i\} \\ &= \deg \sigma_{\square}^6 \\ &= 5 \end{aligned}$$

E.g. 4. [3264, p131, Question(a)]

For 4 general $(k-1)$ -planes $e_1, e_2, e_3, e_4 \cong \mathbb{P}^{k-1}$ in \mathbb{P}^{2k-1} , there are k lines passing all these planes.

Reason: In $Gr(2,2k)$,

$$\begin{aligned} & \# \{l \in Gr(2,2k) \mid l \cap e_i \neq \emptyset, \forall i\} \\ &= \deg \sigma_{\underbrace{\square}_{k-1}}^4 \\ &= k \end{aligned}$$