

# $SL_2(\mathbb{R})$ : Principal Series, Discrete Series and Modular Forms

Last time	$SL_2(\mathbb{R}) \curvearrowright IP'$	group action
$\rightsquigarrow$	$SL_2(\mathbb{R}) \curvearrowright L^2(IP')$	group representation
$\rightsquigarrow$	$(\mathfrak{g}, K) \curvearrowright L^2(IP')_{(K)}$	$(\mathfrak{g}, K)$ -module

We have also talked about the classification of admissible irreducible  $(\mathfrak{g}, K)$ -modules  $U$ :


Suppose  $Kv_n = \varepsilon^n v_n$ ,  $\Omega \cdot v \equiv \gamma v$  ( $\exists v_n \in U, \forall v \in U$ )

(1)  $\gamma \neq \frac{n^2-1}{4}$  for any  $m \in \mathbb{Z}_{\geq 0}$  of opposite parity to  $n$ :



(2)  $\gamma = \frac{m^2-1}{4}$ ,  $m \in \mathbb{Z}_{\geq 0}$ ,  $m \equiv n+1 \pmod{2}$ ,

$|n| < m$ :  $FD_{m-1}[-(m-1), m-1]$



$n > m$ :  $DS_{m+1}^+[m+1, +\infty)$



$n < -m$ :  $DS_{m+1}^-(-\infty, -(m+1)]$

We've also concluded which of the representations are **unitary**:

$PS_{\gamma, n}$ :  $\begin{cases} \textcircled{1} n \text{ is odd \& } \gamma \in \mathbb{R}, \gamma < \frac{1}{4} \\ \textcircled{2} n \text{ is even \& } \gamma \in \mathbb{R}, \gamma < 0 \end{cases}$

$FD_{m-1}$ :  $m=1$

$DS_{m+1}^+$  &  $DS_{m+1}^-$ : always unitary

## Notations:

the split basis

$$h = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad v_+ = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad v_- = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

the compact basis  
of  $sl_2(\mathbb{C})$

$$K = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \quad x_+ = \frac{1}{2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \quad x_- = \frac{1}{2} \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix}$$

the Casimir element

$$\Omega = -\frac{K^2}{4} + \frac{x_+ x_-}{2} + \frac{x_- x_+}{2}$$

Today: principal series & modular forms.

Q: Which  $\chi(g, k)$ -module comes from representations of  $G$ ?

A: For  $G = SL_2(\mathbb{R})$ , EVERY  $(g, k)$ -module

Idea: by direct construction.

$$\begin{aligned}
 &SL_2(\mathbb{R}) \curvearrowright IP' = p \backslash G \\
 &\leadsto SL_2(\mathbb{R}) \curvearrowright \Gamma(IP', E) \quad \text{where } E \text{ is a } G\text{-equivariant line bundle of } IP' \\
 &\quad \parallel \quad \begin{array}{l} \bar{E} = IP' \times \mathbb{R} \\ \Gamma(IP', E) = C^\infty(IP') \end{array} \quad \begin{array}{l} TIP' \\ \nu\text{-field} \end{array} \quad \begin{array}{l} T^*IP' \\ 1\text{-form} \end{array}
 \end{aligned}$$

We will realize  $\Gamma(IP', \bar{E})$  as subspace of  $C^\infty(G)$ .

## Principal Series

- Characters
- Definition of principal series
- Basis of principal series
- Actions on the basis
- Properties: irr & unitary

Discrete Series

Modular forms

## Character

Def: A **character** of a locally cpt group  $H$  is a 1-dim representation of  $H$ :

$$\chi: H \longrightarrow \mathbb{C}^\times$$

$\chi$  is called a **unitary character** when  $\text{Im } \chi \subseteq \{z \mid |z|=1\}$

E.g. 1. the character of  $\mathbb{R}_{>0}^\times \cong \mathbb{R}$  is of the form

$$\chi_s: x \longmapsto x^s \quad s \in \mathbb{C}$$

and the character of  $\mathbb{R}^\times$  is of the form

$$\chi_s \text{sgn}^n: x \longmapsto |x|^s \text{sgn}^n(x) \quad s \in \mathbb{C}, n \in \mathbb{Z}/2\mathbb{Z}$$

E.g. 2. the characters of  $K = \text{SO}_2(\mathbb{R}) \cong S^1$  is of the form

$$\chi^n: \begin{bmatrix} c & -s \\ s & c \end{bmatrix} \longmapsto (c+si)^n \quad n \in \mathbb{Z}$$

E.g. 3. the character of  $P = \left\{ \begin{bmatrix} * & * \\ 0 & * \end{bmatrix} \in \text{SL}_2(\mathbb{R}) \right\}$  is of the form

$$\chi_s \text{sgn}^n: \begin{bmatrix} t & x \\ 0 & \frac{1}{t} \end{bmatrix} \longmapsto |t|^s \text{sgn}^n(t) \quad s \in \mathbb{C}, n \in \mathbb{Z}/2\mathbb{Z}$$

[These are all char of  $P$ , since  $[PP] = N$   
 $\Rightarrow \chi$  is trivial on  $N$   
 $\Rightarrow \chi$  is lifted from  $P/N \cong \mathbb{R}^\times$ ]

We're specially interested in the modulus character

$$\delta = \chi_2 \text{sgn}^0: \begin{bmatrix} t & x \\ 0 & \frac{1}{t} \end{bmatrix} \longmapsto t^2$$

[Check:  $R_p: \mathbb{P}^1 \rightarrow \mathbb{P}^1, [x:y] \mapsto [x:y]_p$   
 $\rightsquigarrow (R_p)_*^*: T_0^* \mathbb{P}^1 \rightarrow T_0^* \mathbb{P}^1$  is multiplication by  $\delta(p)$ ]

"Principal": Characters give us information about the degree of twist of a line bundle.

E.g. 4 For the mobius strip over  $S^1$ , the sections can be viewed as a fct  $f$  on  $\mathbb{R}$  s.t  $f(z+n) = (-1)^n f(z)$ .  $\forall z \in \mathbb{R}$

E.g. 5 we know that  $P \backslash G \cong P'$ , then we have the correspondence:  $(a = [0:1])$

$$\Gamma(T^*P') \longleftrightarrow \{f: G \rightarrow T_a^*P' \mid f(pg) = (R_p)_a^*(f(g))\} \quad \forall p \in P, g \in G$$

$$X \rightsquigarrow \textcircled{H}_X: g \mapsto (R_g)^* X_{R_g(a)}$$

after calculation, we get

$$(R_p)_a^*: T_a^*P' \rightarrow T_a^*P'$$

is just multiplication by  $\delta(p)$ , so

$$\Gamma(T^*P') \longleftrightarrow \{f \in C^\infty(G) \mid f(px) = \delta(p) f(x) \quad \forall p \in P, x \in G\}$$

In general, we can construct the isomorphism between

$$\Gamma(E) \longleftrightarrow \{f \in C^\infty(G) \mid f(px) = \chi(p) f(x) \quad \forall p \in P, x \in X\}$$

where  $E$ : a  $G$ -equivariant line bundle

$\chi$ : a char of  $P$

Definition of  $\text{Ind}^\infty(\chi)$

Def. the principal series  $\text{Ind}^\infty(\chi)$  for a char  $\chi: P \rightarrow \mathbb{C}^\times$  is defined as

$$\text{Ind}^\infty(\chi) := \{f \in C^\infty(G) \mid f(pg) = \chi(p) \underbrace{\delta^{\frac{1}{2}}(p)}_{\text{for the bilinear pairing}} f(g) \quad \forall p \in P, g \in G\}$$

$$\text{e.g. } \text{Ind}^\infty(\delta^{-\frac{1}{2}}) = \{f \in C^\infty(G) \mid f(pg) = f(g) \quad \forall p \in P, g \in G\}$$

$$= C^\infty(P \backslash G) \cong C^\infty(P')$$

$$\text{Ind}^\infty(\delta^{\frac{1}{2}}) = \{f \in C^\infty(G) \mid f(pg) = \delta(p) f(g) \quad \forall p \in P, g \in G\}$$

$$= \Omega^1(P')$$

for  $f \in \text{Ind}^\infty(\delta^{\frac{1}{2}})$ , we can define the integral on  $P \backslash G$

$$\int_{P \backslash G} f(x) dx = \int_K f(k) dk$$

For  $f \in \text{Ind}^\infty(\chi)$ ,  $\varphi \in \text{Ind}^\infty(\chi^{-1})$ , we can define the bilinear pairing

$$\langle -, - \rangle: \text{Ind}^\infty(\chi) \times \text{Ind}^\infty(\chi^{-1}) \longrightarrow \mathbb{R}$$

$$\langle f, \varphi \rangle = \int_{P \backslash G} f \varphi(x) dx$$

$$\begin{aligned}
& G \subset G C^\infty(G) \quad f(-) \mapsto f(-g) \\
& \mapsto G \subset G \text{Ind}^\infty(X) \\
& \mapsto (g, k) \in G \text{Ind}(X), \text{ where} \\
& \text{Ind}(X) = \text{the Harish-Chandra modules of } \text{Ind}^\infty(X) \\
& \quad = \text{subspace of } K\text{-finite vectors.}
\end{aligned}$$

### The basis of $\text{Ind}(X)$

Now suppose  $f \in \text{Ind}(X)$  is an eigenvector of  $K(k)$  with  $f(I)=1$   
 $k.f = \varepsilon^m(k)f$

$$\begin{aligned}
f(k) &= (k.f)(I) = \varepsilon^m(k)f(I) = \varepsilon^m(k) \\
G=PK \quad f(pk) &= \chi(p) \delta^{\frac{1}{2}}(p) f(k) = \chi(p) \delta^{\frac{1}{2}}(p) \varepsilon^m(k) \\
P \cap K = \{\pm I\}
\end{aligned}$$

$$\begin{aligned}
& \text{Condition: } f((-I)(-I)) = f(I \cdot I) = 1 \\
& \Leftrightarrow \chi(-I) \text{sgn}^m(-I) = 1 \\
& \Leftrightarrow (\text{when } \chi = \chi_s \text{sgn}^n) \quad m \equiv n \pmod{2}
\end{aligned}$$

Conclusion: the basis of  $\text{Ind}(\chi_s \text{sgn}^n)$  are  
 $\varepsilon_s^m(q) = (\chi_s \text{sgn}^n)(p) \delta^{\frac{1}{2}}(p) \varepsilon^m(k)$

$$g = pk, m \equiv n \pmod{2}$$

### Actions on the basis: $K, q$ & $\Omega$

$$\begin{aligned}
K: \quad k'. \varepsilon_s^m(q) &= \varepsilon_s^m(pk k') \\
&= (\chi_s \text{sgn}^n)(p) \delta^{\frac{1}{2}}(p) \varepsilon^m(k) \varepsilon^m(k') \\
&= \varepsilon^m(k') \varepsilon_s^m(q)
\end{aligned}$$

$$\begin{aligned}
q: \quad k. \varepsilon_s^m(q) &= \frac{d}{dt} \Big|_{t=0} \exp(tk) \cdot \varepsilon_s^m(q) \\
&= \frac{d}{dt} \Big|_{t=0} \varepsilon^m(\exp(tk)) \cdot \varepsilon_s^m(q) \\
&= \frac{d}{dt} \Big|_{t=0} (\cos t + i \sin t)^m \varepsilon_s^m(q) \\
&= m i \varepsilon_s^m(q)
\end{aligned}$$

$$\begin{aligned}
(k. x_+ \cdot \varepsilon_s^m)(q) &= (zi + mi)(x_+ \cdot \varepsilon_s^m(q)) = (m+2)i(x_+ \cdot \varepsilon_s^m(q)) \\
\Rightarrow (x_+ \cdot \varepsilon_s^m)(q) &= C \varepsilon_s^{m+2}(q) \quad C \text{ to be determined} \\
\Rightarrow (x_+ \cdot \varepsilon_s^m)(1) &= C \quad 2x_+ = h - iK - 2iv_+
\end{aligned}$$

$$\begin{aligned}
C &= (x_+ \cdot \varepsilon_s^m)(1) = \frac{1}{2} [(h \cdot \varepsilon_s^m)(1) - i(k \cdot \varepsilon_s^m)(1) - 2i(v_+ \cdot \varepsilon_s^m)(1)] \\
&= \frac{1}{2} [(s+1) - i(mi) - 2i \cdot 0] \\
&= \frac{1}{2} (s+1+m)
\end{aligned}$$

$$\begin{aligned}
\therefore x_+ \cdot \varepsilon_s^m(q) &= \frac{1}{2} (s+1+m) \varepsilon_s^{m+2}(q) \\
\text{Similarly, } x_- \cdot \varepsilon_s^m(q) &= \frac{1}{2} (s+1-m) \varepsilon_s^{m-2}(q).
\end{aligned}$$

$\Omega$ : By direct calculation, we get  $\Omega = -\frac{k^2}{4} + \frac{x_- x_+}{2} + \frac{x_+ x_-}{2}$

$$\begin{aligned}\Omega \cdot \varepsilon_s^m(q) &= \left[ -\frac{1}{4}(mi)^2 + \frac{1}{2} \cdot \frac{1}{2}(s+1+m) \frac{1}{2}(s+1-m-2) \right. \\ &\quad \left. + \frac{1}{2} \cdot \frac{1}{2}(s+1-m) \frac{1}{2}(s+1+m-2) \right] \varepsilon_s^m(q) \\ &= \frac{1}{4}m^2 + \frac{1}{4}[(s+1)(s+1-2) - m^2] \varepsilon_s^m(q) \\ &= \frac{1}{4}(s^2-1) \varepsilon_s^m(q)\end{aligned}$$

As a conclusion:

$$\begin{aligned}K: \quad k \cdot \varepsilon_s^m(q) &= \varepsilon^m(k') \varepsilon_s^m(q) \\ \Omega: \quad \Omega \cdot \varepsilon_s^m(q) &= \frac{1}{4}(s^2-1) \varepsilon_s^m(q) \\ q: \quad k \cdot \varepsilon_s^m(q) &= m i \varepsilon_s^m(q) \\ x_+ \cdot \varepsilon_s^m(q) &= \frac{1}{2}(s+1+m) \varepsilon_s^{m+1}(q) \\ x_- \cdot \varepsilon_s^m(q) &= \frac{1}{2}(s+1-m) \varepsilon_s^{m-1}(q)\end{aligned}$$

Properties: irr & unitary

(Ir)reducibility of  $\text{Ind}(\chi_s \text{sgn}^n) = \langle \varepsilon_s^m \rangle_{\mathbb{C}\text{-basis}}$   $m \equiv n \pmod{2}$   
Normally: irreducible  $s \notin \mathbb{Z}$  or  $s \equiv n \pmod{2}$



$$\text{Ind}(\chi_s \text{sgn}^n) \cong \text{PS}_{\frac{1}{4}(s^2-1), n}$$

Special situation: reducible  $s \in \mathbb{Z}, s \equiv n+1 \pmod{2}$

	submodule	quotient module
i) $s > 0$	$DS_{s+1}^+ \oplus DS_{s+1}^-$	$FD_{s-1}$
ii) $s = 0$	$DS_1^+ \oplus DS_1^-$	$0$
iii) $s < 0$	$FD_{-s-1}$	$DS_{-s+1}^+ \oplus DS_{-s+1}^-$

Unitary: The irr rep  $\text{PS}_{\frac{1}{4}(s^2-1), n}$  is unitary iff

$$\begin{cases} n=0: & \frac{1}{4}(s^2-1) \in \mathbb{R}, \frac{1}{4}(s^2-1) < 0 & \Leftrightarrow s^2 \in \mathbb{R}, s^2 < 1 \\ n=1: & \frac{1}{4}(s^2-1) \in \mathbb{R}, \frac{1}{4}(s^2-1) < -\frac{1}{4} & \Leftrightarrow s^2 \in \mathbb{R}, s^2 < 0 \end{cases}$$



• reducible  
— unitary irr

Points of unitarity and reducibility of the principal series

The duality between  $\text{Ind}(\chi)$  &  $\text{Ind}(\chi^{-1})$  can explain the situations where  $\text{Re } s = 0$

$\text{Re } s = 0 \Rightarrow \chi = \chi_s \text{sgn}^n$  is unitary,  $\bar{\chi} = \chi^{-1}$   
induce an Hermite inner product of  $\text{Ind}(\chi)$ :

$$\langle f, \phi \rangle := \int_{p \backslash G} f(x) \bar{\phi}(x) dx$$

$$\bar{\phi} \in \text{Ind}(\bar{\chi}) = \text{Ind}(\chi^{-1})$$

## Discrete Series

$$SL_2(\mathbb{R}) \curvearrowright \mathbb{R}P^1$$

$$SL_2(\mathbb{R}) \curvearrowright \mathbb{C}P^1 = \mathbb{R}P^1 \sqcup \mathcal{H}^+ \sqcup \mathcal{H}^-$$

$$\leadsto SL_2(\mathbb{R}) \curvearrowright \Gamma(\mathcal{O}(m)) = \{\text{homogeneous poly of degree } m\}$$

$$m \geq 0: \dim \Gamma(\mathcal{O}(m)) = m+1$$

$$v_{m-2k} = (z+iw)^k (z-iw)^{m-k}$$

$$k, v_{m-2k} = \varepsilon^{m-2k}(k) v_{m-2k}$$

$$k, (z \pm iw) = \varepsilon^{\mp 1}(k) (z \pm iw)$$

$$\left[ \overset{\curvearrowright}{\underset{\curvearrowleft}{v_{-m}}}, \dots, \overset{\curvearrowright}{\underset{\curvearrowleft}{v_m}} \right]_{FD_m}$$

$$m < 0: \Gamma(\mathcal{O}(m)) = 0$$

$$\Gamma(\mathcal{O}(m)|_{\mathcal{H}^+}) \cong \langle v_{p-m} = \frac{(z-iw)^p}{(z+iw)^{p-m}} \mid p \geq 0 \rangle$$

$$\left[ \overset{\curvearrowright}{\underset{\curvearrowleft}{v_{-m}}}, \overset{\curvearrowright}{\underset{\curvearrowleft}{\dots}}, \dots, +\infty \right)_{DS_{|m|}^+}$$

$$\Gamma(\mathcal{O}(m)|_{\mathcal{H}^-}) \cong \langle v_{m-p} = \frac{(z+iw)^p}{(z-iw)^{p-m}} \mid p \geq 0 \rangle$$

$$\left( -\infty, \dots, \overset{\curvearrowright}{\underset{\curvearrowleft}{v_m}} \right]_{DS_{|m|}^-}$$



# Modular forms

Def. A modular form of weight  $k \in \mathbb{Z}$  for the group  $SL_2(\mathbb{Z})$  is a holomorphic fct on  $\mathcal{H}$ , satisfying the following conditions:

$$1) \quad f(\gamma\tau) = (C\tau + d)^k f(\tau)$$

for  $\forall \tau \in \mathcal{H}, \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$

$$j(\gamma, \tau) := C\tau + d$$

$$f(\gamma\tau) = j(\gamma, \tau)^k f(\tau)$$

$$j(\gamma_1 \gamma_2, \tau) = j(\gamma_1, \gamma_2 \tau) j(\gamma_2, \tau)$$

$$\chi(p): P \rightarrow \mathbb{C}^\times$$

$$f(pg) = \chi(p) f(g)$$

$$\chi(p_1 p_2) = \chi(p_1) \chi(p_2)$$

$$2) \quad f \text{ is bounded on } \{z \in \mathcal{H} \mid \text{Im } z > 1\}$$

Remark. We consider modular form as

- fct on  $\mathcal{H}$

- sections in the v.b. of  $SL_2(\mathbb{Z}) \backslash \mathcal{H}$

- a highest weight vector of  $C^\infty(SL_2(\mathbb{Z}) \backslash SL_2(\mathbb{R})) \oplus \mathfrak{g}$

$$\{\text{modular forms of weight } m\} \longrightarrow C^\infty(SL_2(\mathbb{R}))$$

$$f \longmapsto [\Phi_f: g \mapsto f(g(i)) j(g, i)^{-m}]$$

In general, we define

$$C^\infty(\mathcal{H}) \longrightarrow C^\infty(SL_2(\mathbb{R}))$$

$$f \longmapsto \Phi_f: g \mapsto f(g(i)) j(g, i)^{-m}$$


then for  $\forall f \in C^\infty(\mathcal{H}), g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R}),$

$$K: k \cdot \Phi_f(g) = \varepsilon^{-m}(k) \Phi_f(g)$$

$$g: k \cdot \Phi_f(g) = -mi \Phi_f(g)$$

$$x_+ \cdot \Phi_f(g) = -\frac{2i}{(c-i+d)^2} \Phi_{\frac{\partial f}{\partial \bar{z}}}(g) \quad \underline{\text{f is hdo}} \quad 0$$

$$x_- \cdot \Phi_f(g) = \frac{2i}{(c+i+d)^2} \Phi_{\frac{\partial f}{\partial \bar{z}}}(g) - m \frac{c-i}{c+i+d} \Phi_f(g)$$



Proof. Recall the definition

$$\Phi_f(g) = f(g(i)) j(g, i)$$

$$\Phi_f \begin{pmatrix} a & b \\ c & d \end{pmatrix} = f\left(\frac{a+i}{c+i}\right) (c+i)^{-m}$$

We have

$$\begin{aligned} k. \Phi_f(g) &= \Phi_f(gk) \\ &= f(gk(i)) j(gk, i)^{-m} \\ &= f(g(i)) j(g, k(i))^{-m} j(k, i)^{-m} \\ &= \Phi_f(g) \cdot (i \sin t + \cos t)^{-m} \\ &= E^{-m}(k) \Phi_f(g) \end{aligned}$$

$$k. \Phi_f(g) = -mi \Phi_f(g)$$

$$\begin{aligned} x_+. \Phi \begin{pmatrix} a & b \\ c & d \end{pmatrix} &= \frac{d}{dt} \Big|_{t=0} \Phi \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix} \exp(tx_+) \right) \\ &= \frac{d}{dt} \Big|_{t=0} \Phi \begin{pmatrix} a + \frac{1}{2}at - \frac{i}{2}bt & b - \frac{i}{2}bt - \frac{1}{2}at \\ c + \frac{1}{2}ct - \frac{i}{2}dt & d - \frac{i}{2}dt - \frac{1}{2}ct \end{pmatrix} \\ &= \frac{1}{2} [(a-bi)\Phi_1 - (-b-ai)\Phi_2 + (c-di)\Phi_3 + (-d-ci)\Phi_4] \\ &= \frac{1}{2i} [(a+ib)(\Phi_1 - i\Phi_2) + (c+d)(\Phi_3 - i\Phi_4)] \end{aligned}$$

$$\begin{aligned} \Phi_{f,1} &= \frac{\partial}{\partial a} \Phi_f = \frac{\partial}{\partial a} \left[ f\left(\frac{a+i}{c+i}\right) (c+i)^{-m} \right] \\ &= (c+i)^{-m} \frac{\partial f}{\partial z} \cdot \frac{i}{c+i} + (c+i)^{-m} \frac{\partial f}{\partial \bar{z}} \cdot \frac{-i}{-c+i} \\ &= \frac{i}{c+i} \Phi_{f_z} - \frac{i}{-c+i} \Phi_{f_{\bar{z}}} \end{aligned}$$

$$f_z = \frac{\partial f}{\partial z} \quad f_{\bar{z}} = \frac{\partial f}{\partial \bar{z}}$$

Similarly,  $\Phi_{f,2} = \frac{1}{c+i} \Phi_{f_z} + \frac{1}{-c+i} \Phi_{f_{\bar{z}}}$

$$\Phi_{f,3} = -i \frac{a+ib}{(c+i)^2} \Phi_{f_z} + i \frac{-a+ib}{(-c+i)^2} \Phi_{f_{\bar{z}}} - \frac{mi}{c+i} \Phi_f$$

$$\Phi_{f,4} = -\frac{a+ib}{(c+i)^2} \Phi_{f_z} - \frac{-a+ib}{(-c+i)^2} \Phi_{f_{\bar{z}}} - \frac{m}{c+i} \Phi_f$$

then  $x_+. \Phi_f \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \frac{1}{2i} [(a+ib)(\Phi_{f,1} - i\Phi_{f,2}) + (c+d)(\Phi_{f,3} - i\Phi_{f,4})]$

$$= -\frac{2i}{(c-i)^2} \Phi_{f_{\bar{z}}}$$

similarly,  $x_-. \Phi_f \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \frac{1}{2i} [(a-i)(\Phi_{f,1} + i\Phi_{f,2}) + (c-d)(\Phi_{f,3} + i\Phi_{f,4})]$

$$= \frac{2i}{(c+i)^2} \Phi_{f_z} - m \frac{c-i}{c+i} \Phi_f$$