

Eine Woche, ein Beispiel. 5.7. Lie group

Temporary roadmap of Lie Group


Why is it so difficult to write examples for Lie group?

- Plenty of examples, even restricted to matrix Lie groups:

- List of examples. https://en.wikipedia.org/wiki/Table_of_Lie_groups

[Some important subgroups are not written in the list,
e.g. parabolic subgroups, Heisenberg group, etc.]

$G \rightarrow K$ maximal cpt connected subgroup



[For examples of non matrix Lie group $(\widetilde{SL(2, IR)}, M_p(2n, IR))$, see
<https://math.stackexchange.com/questions/206117/is-there-a-non-matrix-lie-group>]

[On the contrary, every f.d. Lie alg can be viewed as a matrix Lie algebra by Ado's theorem, see <http://www.math.ubc.ca/~reichst/Ado's-Theorem.pdf>.]

- These groups are closely connected to each other.

e.g. the Whitehead tower:

$$\rightarrow \text{Fivebrane}(n) \rightarrow \text{String}(n) \rightarrow \text{Spin}(n) \rightarrow \text{SO}(n) \rightarrow \text{O}(n)$$

- Plenty of special phenomena on low-dimensional case:

e.g. $GL(1, IR) \cong IR^*$ is not connected

e.g. $SU(2) \cong S^3$

e.g. $so(4, IR)$ is not simple

\downarrow
 $SU(2) \cong S^3$
 \downarrow
 $SO(3, IR) \cong RP^3$
diff
not as variety

e.g. <https://www.zhihu.com/question/47264301/answer/320472431>

- Fruitful structures on examples

- important subgroups

e.g. maximal torus, unipotent subgroup

Borel subgroup, parabolic subgroup,

discrete subgroup,

radical, unipotent radical, derived group

$$G = [G, G] R(G)$$

* centralizer & normalizer of subgroups

This article may give a classification of the cocompact subgroup:
<https://people.uleth.ca/~dave.morris/papers/cocompact.pdf>

One page for defining these groups by pasting from

Anthony W. Knapp - Representation Theory of Semisimple Groups An Overview Based on Examples

Complex groups:

$$\mathrm{GL}(n, \mathbb{C}) = \{\text{nonsingular } n\text{-by-}n \text{ complex matrices}\}$$

$$\mathrm{SL}(n, \mathbb{C}) = \{g \in \mathrm{GL}(n, \mathbb{C}) \mid \det g = 1\}$$

$$\mathrm{SO}(n, \mathbb{C}) = \{g \in \mathrm{SL}(n, \mathbb{C}) \mid gg^{\mathrm{tr}} = 1\}$$

$$\mathrm{Sp}(n, \mathbb{C}) = \left\{ g \in \mathrm{SL}(2n, \mathbb{C}) \mid g^{\mathrm{tr}} J g = J \text{ for } J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} \right\}.$$

Compact groups:

~~$$\mathrm{SO}(n, \mathbb{R}) = \mathrm{SO}(n) = \{g \in \mathrm{SL}(n, \mathbb{C}) \mid g^{\mathrm{tr}} g = 1 \text{ and } g \text{ has real entries}\}$$~~

~~$$\mathrm{U}(n) = \{g \in \mathrm{GL}(n, \mathbb{C}) \mid \bar{g}^{\mathrm{tr}} g = 1\}$$~~

~~$$\mathrm{SU}(n) = \{g \in \mathrm{U}(n) \mid \det g = 1\}$$~~

~~$$\mathrm{Sp}(n) = \left\{ g \in \mathrm{U}(2n) \mid g^{\mathrm{tr}} J g = J \text{ for } J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} \right\}.$$~~

Real noncompact groups: We list G , \mathfrak{g} , K , and \mathfrak{k} for the noncomplex non-compact classical groups.

G	\mathfrak{g}	K	\mathfrak{k}
$\mathrm{SL}(n, \mathbb{R})$	$\mathfrak{sl}(n, \mathbb{R})$	$\mathrm{SO}(n)$	$\mathfrak{so}(n)$
$\mathrm{SL}(n, \mathbb{H})$	$\mathfrak{sl}(n, \mathbb{H})$	$\mathrm{Sp}(n)$	$\mathfrak{sp}(n)$
$\mathrm{SO}_0(m, n)$	$\mathfrak{so}(m, n)$	$\mathrm{SO}(m) \times \mathrm{SO}(n)$	$\mathfrak{so}(m) \oplus \mathfrak{so}(n)$
$\mathrm{SU}(m, n)$	$\mathfrak{su}(m, n)$	$\mathrm{S}(\mathrm{U}(m) \times \mathrm{U}(n))$	$\mathfrak{s}(\mathrm{u}(m) \oplus \mathrm{u}(n))$
$\mathrm{Sp}(m, n)$	$\mathfrak{sp}(m, n)$	$\mathrm{Sp}(m) \times \mathrm{Sp}(n)$	$\mathfrak{sp}(m) \oplus \mathfrak{sp}(n)$
$\mathrm{Sp}(n, \mathbb{R})$	$\mathfrak{sp}(n, \mathbb{R})$	$\mathrm{U}(n)$	$\mathfrak{u}(n)$
$\mathrm{SO}^*(2n)$	$\mathfrak{so}^*(2n)$	$\mathrm{U}(n)$	$\mathfrak{u}(n)$.

$\mathrm{SL}(n, \mathbb{R})$ and $\mathrm{SL}(n, \mathbb{H})$ refer to matrices of determinant one with real and quaternion entries, respectively. $\mathrm{SO}_0(m, n)$, $\mathrm{SU}(m, n)$, and $\mathrm{Sp}(m, n)$ are the linear isometry groups for the Hermitian form

$$|z_1|^2 + \cdots + |z_m|^2 - |z_{m+1}|^2 - \cdots - |z_{m+n}|^2$$

defined over \mathbb{R} , \mathbb{C} , and \mathbb{H} , respectively, with the subscript “0” referring to the identity component. The group $K = \mathrm{S}(\mathrm{U}(m) \times \mathrm{U}(n))$ for $\mathrm{SU}(m, n)$ is the subgroup of $\mathrm{U}(m) \times \mathrm{U}(n)$ of matrices of determinant one.

The group $\mathrm{Sp}(n, \mathbb{R})$ is the subgroup of real matrices in $\mathrm{Sp}(n, \mathbb{C})$ and can be conjugated by a unitary matrix so as to become

$$\{g \in \mathrm{SU}(n, n) \mid g^{\mathrm{tr}} J g = J\};$$

then $\mathrm{SO}^*(2n)$ is the analogous group

$$\left\{ g \in \mathrm{SU}(n, n) \mid g^{\mathrm{tr}} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} g = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \right\}.$$

- decomposition: not easy to prove, but still of great significant.

https://en.wikipedia.org/wiki/Lie_group_decomposition

e.g. Cartan decomposition

Jordan - Chevalley decomposition

Bruhat decomposition

:


- group, mfld, variety, etc. Common questions.

- Hopf algebra structure of homology & cohomology group under mild requirements.

- A large amount of related theories.

- structure theory of Lie group & Lie algebras.

" \rightarrow ": represent unfamiliar objects with simpler objects.



- representation theory

- * group / alg, f.d / inf.d

- * highest weight theory

- * Schul - Weyl duality

- * Borel - Weil - Bott theorem

- * geometric representation theory

- * Langlands program

<https://mathoverflow.net/questions/235917/roadmap-to-geometric-representation-theory-leading-to-langlands>

- algebraic topology, Bott periodicity

- algebra groups.


- ... see https://en.wikipedia.org/wiki/List_of_Lie_groups_topics

<http://staff.ustc.edu.cn/~wangzuoq/Courses/13F-Lie/Lie.html>

Basic knowledge which has been well-written in [Naive Lie Theory].

- canonical maximal tori
- center

- (1) $Z(\mathrm{SO}(2m)) = \{\pm 1\}$.
- (2) $Z(\mathrm{SO}(2m+1)) = \{1\}$.
- (3) $Z(\mathrm{U}(n)) = \{\omega \mathbf{1} : |\omega| = 1\}$.
- (4) $Z(\mathrm{SU}(n)) = \{\omega \mathbf{1} : \omega^n = 1\}$.
- (5) $Z(\mathrm{Sp}(n)) = \{\pm 1\}$.



In alg top: homotopy groups:

$\text{typo: } \pi_*(U(n)) = \mathbb{Z}$

see

<http://felix.physics.sunysb.edu/~abanov/Teaching/Spring2009/Notes/abanov-cpA1-upload.pdf>

more about $\pi_i(SO(n, \mathbb{R}))$:

<https://ncatlab.org/nlab/show/orthogonal+group#HomotopyGroups>

more about $\pi_i(G)$ (in Chinese):

<https://www.zhihu.com/question/443652195>

more about $\pi_n(G)$:

<https://mathoverflow.net/questions/8957/homotopy-groups-of-lie-groups>

$\pi_i(SO(m, n))$:

<https://math.stackexchange.com/questions/4126129/homotopy-group-of-op-q>

my try: Cartan decomposition \Rightarrow only consider cpt Lie group.

fibration + low-dimensional understanding, + Bott periodicity

e.g. • $O(1) = S^0$, a two-point discrete space

$U(n) \cong SU(n) \times S^1$

• $SO(1) = \{1\}$

• $SO(2)$ is S^1

• $SO(3)$ is RP^3 [3]

• $SO(4)$ is doubly covered by $SU(2) \times SU(2) = S^3 \times S^3$.

https://en.wikipedia.org/wiki/Orthogonal_group


Homotopy groups of orthogonal groups								
π_1	π_2	π_3	π_4	π_5	π_6	π_7	π_8	π_9
$SO(2)$	Z	0	0	0	0	0	0	0
$SO(3)$	$[Z_2]$	0	Z	Z_2	Z_2	Z_{12}	Z_2	Z_2
$SO(4)$	Z_2	$[0]$	$(Z)^{\times 2}$	$(Z_2)^{\times 2}$	$(Z_2)^{\times 2}$	$(Z_{12})^{\times 2}$	$(Z_2)^{\times 2}$	$(Z_2)^{\times 2}$
$SO(5)$	Z_2	0	$[Z]$	Z_2	Z_2	0	Z	0
$SO(6)$	Z_2	0	Z	$[0]$	Z	0	Z	Z_{24}
$SO(n), n > 6$	Z_2	0	Z	0	0	0	0	0

Activate Windows

my understanding

fibration: [Hatcher Example 4.54, 4.55]

space of n -frames in \mathbb{R}^k



Structure theory of Lie group

Ref:

<http://www.math.columbia.edu/~maki/sumi/cld/reductivegroups.pdf>

← notations, results, examples, pictures.

Today:

$$G = GL(2, \mathbb{C})$$

center

$$Z(G) = \mathbb{C}^{\times} \quad \pi_1(G) = \mathbb{Z}$$

radical

$$\text{rad}(G) := R(G) = \mathbb{C}^{\times}$$

reductive

$$\text{Unipotent rad} \quad \text{rad}_u(G) := R_u(G) = \{I\}$$

$$\text{rank } G = 2$$

canonical maximal torus $T = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix}$

$$C_G(T) = T$$

$$N_G(T) = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \cup \begin{pmatrix} * & * \\ 0 & * \end{pmatrix}$$

$$W(G) = S_2 \quad \text{Weyl group} \\ = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \cup \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$


$$X^* := X^*(T) = \langle \varepsilon_i : (t_1 t_2) \mapsto t_i \rangle_{\mathbb{Z}} \quad \hookrightarrow W(G) \quad \alpha \mapsto \alpha(g^{-1}g)$$

$$X_{\neq}^* := X_{\neq}(T) = \left\langle \begin{array}{c} \varepsilon_1^* : t \mapsto (t_1) \\ \varepsilon_2^* : t \mapsto (1_t) \end{array} \right\rangle_{\mathbb{Z}} \quad \hookrightarrow W(G) \quad \lambda \mapsto g \lambda(-)g^{-1}$$

$$\Phi = \{\varepsilon_1 - \varepsilon_2, -(\varepsilon_1 + \varepsilon_2)\} := \{\alpha_{12}, -\alpha_{12}\}$$

$$\Phi^V = \{\alpha_{12}^V, -\alpha_{12}^V\} = \{\varepsilon_1^* - \varepsilon_2^*, -(\varepsilon_1^* + \varepsilon_2^*)\}$$

$$\boxed{r_? = \text{action of } \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} : \begin{array}{ccc} X^* & \longrightarrow & X^* \\ \varepsilon_1 & \longmapsto & \varepsilon_2 \\ \varepsilon_2 & \longmapsto & \varepsilon_1 \end{array} \Rightarrow r_? \text{ is reflection and } r_? = r_{\alpha_{12}} \quad X := x_1 \varepsilon_1 + x_2 \varepsilon_2} \\ X - r_?(X) = x_1 \varepsilon_1 + x_2 \varepsilon_2 - x_1 \varepsilon_2 - x_2 \varepsilon_1 = (x_1 - x_2) \alpha_{12} \quad x := x_1 \varepsilon_1 + x_2 \varepsilon_2 \\ \langle -, \alpha_{12}^V \rangle : x_1 \varepsilon_1 + x_2 \varepsilon_2 \mapsto x_1 - x_2 \quad \Rightarrow \alpha_{12}^V = \varepsilon_1^* - \varepsilon_2^*}$$



$$G = SL(2, \mathbb{C})$$

$$Z(G) = \{\pm 1\} \quad \pi_1(G) = \{*\}$$

$$\text{rad}(G) = \{I\} \Rightarrow \text{semisimple}$$

$$\text{rad}_u(G) = \{I\}$$

$$\text{rank } G = 1$$

canonical maximal torus $T = \begin{pmatrix} t & \\ & t^{-1} \end{pmatrix}$

$$C_G(T) = T$$

$$N_G(T) = \{t' \mid t' \sim t\}$$

$$W(G) = S_2 = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \cup \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$$

$$X^* = \langle \varepsilon_i : (t_{-t}) \mapsto t \rangle_{\mathbb{Z}} \quad \text{with } \alpha \mapsto \alpha(g^{-1} - g) \quad \alpha_1 = 2\varepsilon_1!$$

$$X_t = \langle \varepsilon_i^t : t \mapsto (t_{-t}) \rangle_{\mathbb{Z}} \quad \text{with } \lambda \mapsto g \cdot \lambda(-) \cdot g^{-1}$$

$$\Phi = \{\alpha_{12}, -\alpha_{12}\} = \{2\varepsilon_1, -2\varepsilon_1\}$$

$$\vec{\Phi}^V = \{d_{12}^V, -d_{12}^V\} = \{\varepsilon_1^+, -\varepsilon_1^*\}$$

$$\left[\begin{array}{l} r_? = \text{action of } (\begin{smallmatrix} 1 & 1 \\ 1 & 0 \end{smallmatrix}): X^* \rightarrow X^* \quad \varepsilon_i \mapsto -\varepsilon_i \\ x - r_?(x) = x_0 \varepsilon_i - x_0 (-\varepsilon_i) = x_0 \alpha_{12} \stackrel{r_? \text{ ref}}{\substack{+ r_2 = r_{12}}} x := x_0 \varepsilon_i \\ \langle -, \alpha_{12}^\vee \rangle: x_0 \varepsilon_i \mapsto x_0 \quad \Rightarrow \alpha_{12}^\vee = \varepsilon_i^* \end{array} \right]$$

A horizontal number line starting at -3 and ending at 3. It has major tick marks at integer intervals (-3, -2, -1, 0, 1, 2, 3). Between each integer, there are four smaller tick marks, dividing the interval into four equal segments of length 1/4. The first tick mark after -3 is labeled $-\frac{3}{4}$, the second $-\frac{1}{2}$, the third $-\frac{1}{4}$, and the fourth $\frac{1}{4}$. The last tick mark before 3 is labeled $\frac{7}{4}$.

ω_1, ω_2

$$\bullet \quad C^* = P$$

二

$G = GL(3, \mathbb{C})$

$$\begin{aligned} Z(G) &= \mathbb{C}^\times & \pi_1(G) &= \mathbb{Z} \\ \text{rad}(G) &= \mathbb{C}^\times & \left. \begin{aligned} \text{rad}(G) &= \{I\} \end{aligned} \right\} &\Rightarrow \text{reductive} \\ \text{rank } G &= 3 & &\text{but not semi} \end{aligned}$$

canonical maximal torus $T = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix}$
 $C_G(T) = T$
 $N_G(T) = \text{monomial matrixs}$
 $W(G) = S_3$

$$X^* = \langle \varepsilon_i, \begin{pmatrix} t_1 & & \\ & t_2 & \\ & & t_3 \end{pmatrix} \mapsto t_i \rangle_{\mathbb{Z}}$$

$$X_* = \langle \varepsilon_i^*, t \mapsto (t, t, t) \rangle_{\mathbb{Z}}$$

$$\Phi = \{\alpha_{12}, \alpha_{23}, \alpha_{13}, -\alpha_{12}, -\alpha_{23}, -\alpha_{13}\}$$

$$\Phi^\vee = \{\alpha_{12}^\vee, \alpha_{23}^\vee, \alpha_{13}^\vee, -\alpha_{12}^\vee, -\alpha_{23}^\vee, -\alpha_{13}^\vee\}$$

$$\left[\begin{array}{l} r_{\alpha_{12}} = \text{action of } \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}, X^* \rightarrow X^* \\ \begin{array}{l} \varepsilon_1 \mapsto \varepsilon_2 \\ \varepsilon_2 \mapsto \varepsilon_1 \\ \varepsilon_3 \mapsto \varepsilon_3 \end{array} \end{array} \right]$$

$$x - r_{\alpha_{12}}(x) = x_1 \varepsilon_1 + x_2 \varepsilon_2 + x_3 \varepsilon_3 - x_1 \varepsilon_2 - x_2 \varepsilon_1 - x_3 \varepsilon_3 \\ = (x_1 - x_2) \alpha_{12}$$


$$\langle -, \alpha_{12}^\vee \rangle : x_1 \varepsilon_1 + x_2 \varepsilon_2 + x_3 \varepsilon_3 \mapsto x_1 - x_2 \quad \Rightarrow \alpha_{12}^\vee = \varepsilon_1^* - \varepsilon_2^*$$

$$\text{Similarly, } \alpha_{23}^\vee = \varepsilon_2^* - \varepsilon_3^*$$

$$\alpha_{ij} := \varepsilon_i - \varepsilon_j$$

$$\text{where } \alpha_{ij}^\vee = \varepsilon_i^* - \varepsilon_j^*$$

$$\begin{aligned} \varepsilon_1 &\mapsto \varepsilon_1 \left(\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} t_1 & & \\ & t_2 & \\ & & t_3 \end{pmatrix} \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \right) \\ &= \varepsilon_1 \begin{pmatrix} t_2 & & \\ & t_1 & \\ & & t_3 \end{pmatrix} = \varepsilon_2 \end{aligned}$$



$$x \subseteq \cdot \subseteq /$$

$$Q \subseteq X^* \subseteq P$$

$G = SL(3, \mathbb{C})$

$$Z(G) = \mathbb{Z}/3\mathbb{Z} \quad \pi_1(G) = \langle \gamma \rangle$$

$\text{rad}(G) = \{I\} \Rightarrow \text{semisimple}$

$$\text{rad}_u(G) = \{I\}$$

rank $G = 2$

canonical maximal torus $T = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix}$

$$C_G(T) = T$$

$N_G(T) = \text{monomial matrixs}$

$$W(G) = S_3$$

$$X^* = \langle \varepsilon_i : \begin{pmatrix} t_1 & t_2 & t_3 \end{pmatrix} \mapsto t_i \rangle_{\mathbb{Z}} / \mathbb{Z}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3)$$

$$X_\gamma = \langle \varepsilon_1^* - \varepsilon_2^*, \varepsilon_2^* - \varepsilon_3^* \rangle_{\mathbb{Z}}$$

$\varepsilon_1^*, \varepsilon_2^*, \varepsilon_3^*$: dual basis of $\varepsilon_1, \varepsilon_2, \varepsilon_3$

$$\Phi = \{\alpha_{12}, \alpha_{23}, \alpha_{13}, -\alpha_{12}, -\alpha_{23}, -\alpha_{13}\}$$

$$\Phi^\vee = \{\alpha_{12}^\vee, \alpha_{23}^\vee, \alpha_{13}^\vee, -\alpha_{12}^\vee, -\alpha_{23}^\vee, -\alpha_{13}^\vee\}$$

$r_{\alpha_{12}}$ = action of $\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}$, $X^* \rightarrow X^*$

$$\begin{aligned} \varepsilon_1 &\mapsto \varepsilon_2 \\ \varepsilon_2 &\mapsto \varepsilon_1 \\ \varepsilon_3 &\mapsto \varepsilon_3 \end{aligned}$$

$$\alpha_{ij} := \varepsilon_i - \varepsilon_j$$

$$\text{where } \alpha_{ij}^\vee = \varepsilon_i^* - \varepsilon_j^*$$


$$\begin{aligned} \varepsilon_1 &\mapsto \varepsilon_1 \left(\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} t_1 & t_2 & t_3 \end{pmatrix} \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \right) \\ &= \varepsilon_1 \begin{pmatrix} t_2 & t_1 & t_3 \end{pmatrix} = \varepsilon_2 \end{aligned}$$

$$x - r_{\alpha_{12}}(x) = x_1 \varepsilon_1 + x_2 \varepsilon_2 + x_3 \varepsilon_3 - x_1 \varepsilon_2 - x_2 \varepsilon_1 - x_3 \varepsilon_3 \quad x = x_1 \varepsilon_1 + x_2 \varepsilon_2 + x_3 \varepsilon_3$$

$$= (x_1 - x_2) \alpha_{12}$$

$$\langle -, \alpha_{12}^\vee \rangle : x_1 \varepsilon_1 + x_2 \varepsilon_2 + x_3 \varepsilon_3 \mapsto x_1 - x_2 \quad \Rightarrow \alpha_{12}^\vee = \varepsilon_1^* - \varepsilon_2^*$$

$$\text{Similarly, } \alpha_{23}^\vee = \varepsilon_2^* - \varepsilon_3^*$$



$$\begin{array}{c} \subseteq \\ Q \subseteq X^* = P \\ \mathbb{Z}/3\mathbb{Z} \end{array}$$

$$G = \mathrm{PGL}(3, \mathbb{C}) = \mathrm{SL}(3, \mathbb{C}) / \mu_3$$

$$Z(G) = \{\text{I}\} \quad \pi_1(G) = \mathbb{Z}/3\mathbb{Z}$$

$\mathrm{rad}(G) = \{\text{I}\} \Rightarrow$ semisimple

$$\mathrm{rad}_u(G) = \{\text{I}\}$$

rank $G = 2$

$$\mathbb{C}^* \cong \mathbb{C}_{\mu_3}^* \times \mathbb{C}^*$$

$$t_i^3 \leftrightarrow \begin{pmatrix} t_1 & t_2 \\ t_1 & t_2 \\ t_1 & t_2 \end{pmatrix} \times \begin{pmatrix} t_1 & t_2 \\ t_1 & t_2 \\ t_1 & t_2 \end{pmatrix}$$

canonical maximal torus $T = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} / \mu_3$

$$C_G(T) = T$$

$N_G(T)$ = monomial matrixs / μ_3

$$W(G) = S_3$$

$$X^* = \{n_1\epsilon_1 + n_2\epsilon_2 + n_3\epsilon_3 \mid n_i \in \mathbb{Z}, 3 \mid \sum n_i\} /_{(\epsilon_1 + \epsilon_2 + \epsilon_3)\mathbb{Z}}$$

$$= \langle \alpha_{12}, \alpha_{23} \rangle_{\mathbb{Z}}$$

$\epsilon_i : \begin{pmatrix} t_1 & t_2 & t_3 \end{pmatrix} \rightarrow t_i$ is no longer well-defined

$$\alpha_{ij} := \epsilon_i - \epsilon_j$$

$$X^* = \{n_1\epsilon_1^* + n_2\epsilon_2^* + n_3\epsilon_3^* \mid n_i \in \frac{1}{3}\mathbb{Z}, \sum n_i = 0\}$$

$$= \left\langle \frac{1}{3}\epsilon_1^* + \frac{1}{3}\epsilon_2^* - \frac{2}{3}\epsilon_3^*, \epsilon_2^* - \epsilon_3^* \right\rangle_{\mathbb{Z}}$$


$$\Phi = \{\alpha_{12}, \alpha_{23}, \alpha_{13}, -\alpha_{12}, -\alpha_{23}, -\alpha_{13}\}$$

$$\Phi^\vee = \{\alpha_{12}^\vee, \alpha_{23}^\vee, \alpha_{13}^\vee, -\alpha_{12}^\vee, -\alpha_{23}^\vee, -\alpha_{13}^\vee\}$$


$\epsilon_1^*, \epsilon_2^*, \epsilon_3^*$: dual basis of $\epsilon_1, \epsilon_2, \epsilon_3$

$$\alpha_{ij} := \epsilon_i - \epsilon_j$$

where $\alpha_{ij}^\vee = \epsilon_i^* - \epsilon_j^*$



$$Q = X^* \subseteq P$$



fundamental domain of

Weyl group []
affine Weyl group []
extended affine Weyl group []

$$G = \mathrm{Sp}(4, \mathbb{C}) = \{ A \in GL(4, \mathbb{C}) \mid A^T M A = M \}$$

$Z(G) = \{\pm I\}$ $\pi_1(G) = \mathbb{Z}_2$
 $\mathrm{rad}(G) = \{I\} \Rightarrow \text{semisimple}$
 $\mathrm{rad}_u(G) = \{I\}$
 $\mathrm{rank } G = 2$

$M = \begin{pmatrix} & & -1 & 1 \\ & -1 & & \\ -1 & & & \\ & & 1 & -1 \end{pmatrix}$
 canonical maximal torus $T = \begin{pmatrix} t_1 & & & \\ & t_2 & & \\ & & t_3^{-1} & \\ & & & t_4^{-1} \end{pmatrix}$
 $C_G(T) = T$
 $N_G(T) = \text{monomial matrixs}$
 $W(G) = D_4 \subseteq S_4$

$$\mathrm{sp}(4, \mathbb{C}) = \{ A \in M^{4 \times 4}(\mathbb{C}) \mid A^T M + M A = 0 \}$$



$$= \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ & -a_{13} & & \\ a_{21} & a_{22} & a_{23} & \\ & & -a_{31} & a_{31} \end{pmatrix} \in M^{4 \times 4}(\mathbb{C}) \right\}$$

$$X^* = \mathbb{Z}\varepsilon_i / (\varepsilon_1 + \varepsilon_4, \varepsilon_2 + \varepsilon_3) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$$

$$X_* = \mathbb{Z}\lambda_{14} \oplus \mathbb{Z}\lambda_{23}$$

$$\Phi = \{\pm \alpha_{12}, \pm \alpha_{13}, \pm \alpha_{14}, \pm \alpha_{23}\} = \{\pm (\varepsilon_1 - \varepsilon_2), \pm (\varepsilon_1 + \varepsilon_2), \pm 2\varepsilon_1, \pm 2\varepsilon_2\}$$

$$\Phi^\vee = \{\pm \alpha_{12}^\vee, \pm \alpha_{13}^\vee, \pm \alpha_{14}^\vee, \pm \alpha_{23}^\vee\} = \{\pm (\lambda_{14} - \lambda_{23}), \pm (\lambda_{14} + \lambda_{23}), \pm \lambda_{14}, \pm \lambda_{23}\}$$



$$\Phi \subseteq Q \subseteq X^* = P$$

$$\Phi^\vee$$

$\mathbb{Z}/2\mathbb{Z}$

$$G = SO(5, \mathbb{C}) \cong \{ A \in SL(5, \mathbb{C}) \mid A^T M A = M \}$$

$$Z(G) = \{ Id \} \quad \pi_1(G) = \mathbb{Z}/2\mathbb{Z}$$

$\text{rad}(G) = \{ I \} \Rightarrow \text{semisimple}$

$$\text{rad}_u(G) = \{ I \}$$

$$\text{rank } G = 2$$

$$M = \begin{pmatrix} & & & & \\ & -1 & & & \\ & & 1 & & \\ & & & -1 & \\ & & & & 1 \end{pmatrix}$$

canonical maximal torus

$$C_G(T) = T$$

$N_G(T) = \text{monomial matrixs}$

$$W(G) = D_4$$

$$T = \begin{pmatrix} t_1 & & & \\ & t_2 & & \\ & & t_3 & \\ & & & t_4^{-1} \end{pmatrix}$$

$$SO(5, \mathbb{C}) = \{ A \in M^{5 \times 5}(\mathbb{C}) \mid A^T M + M A = 0 \}$$



$$= \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & 0 \\ & 0 & a_{14} & & \\ & & 0 & a_{23} - a_{13} & \\ & & & a_{32} - a_{22} & a_{12} \\ 0 & & & a_{41} - a_{31} & a_{21} - a_{11} \end{pmatrix} \in M^{5 \times 5}(\mathbb{C}) \right\}$$

$$X^* = \mathbb{Z}\varepsilon_i / (\varepsilon_1 + \varepsilon_5, \varepsilon_2 + \varepsilon_4, \varepsilon_3) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$$

$$X_* = \mathbb{Z}\lambda_{15} \oplus \mathbb{Z}\lambda_{24}$$

$$\Phi = \{ \pm \alpha_{12}, \pm \alpha_{13}, \pm \alpha_{14}, \pm \alpha_{23} \} = \{ \pm (\varepsilon_1 - \varepsilon_2), \pm \varepsilon_1, \pm (\varepsilon_1 + \varepsilon_2), \pm \varepsilon_2 \}$$

$$\Phi^\vee = \{ \pm \alpha_{12}^\vee, \pm \alpha_{13}^\vee, \pm \alpha_{14}^\vee, \pm \alpha_{23}^\vee \} = \{ \pm (\lambda_{15} - \lambda_{24}), \pm 2\lambda_{15}, \pm (\lambda_{15} + \lambda_{24}), \pm 2\lambda_{24} \}$$




$$\Phi \subseteq Q = X^* \subseteq P$$

$\mathbb{Z}/2\mathbb{Z}$

$$\Phi^\vee$$

Now let's enjoy the beauty of low-dim root system!



Coxeter number

The Weyl groups for classical groups:

$$W(\mathrm{U}(n)) = S(n).$$

- The Weyl group of $SU(n)$ is still $S(n)$.


- The Weyl group of $SO(2l+1)$ is $G(l)$, the group of permutations φ of the set $\{-l, \dots, -1, 1, \dots, l\}$ with $\varphi(-k) = -\varphi(k)$ for all $1 \leq k \leq l$.

- The Weyl group of $SO(2l)$ is the subgroup $SG(l)$ of $G(l)$ that consists of even permutations.


- The Weyl group of $Sp(n)$ is still $G(n)$.

<http://staff.ustc.edu.cn/~wangzuoq/Courses/13F-Lie/Notes/Lec%2027.pdf>

https://en.wikipedia.org/wiki/Root_system



$k = \overline{k}$, $\text{char } k = 0$.



The following table comes from [linear algebraic groups and finite groups of Lie type, p72].
 Errata for this book: https://www.mathematik.uni-kl.de/~malle/download/MT_errata.pdf

Table 9.2 *Isogeny types of simple algebraic groups*

Φ	$\Lambda(\Phi)$	G_{sc}	G_{ad}	in between
$A_{n-1}, n \geq 2$	Z_n	SL_n	PGL_n	$SL_n/Z_d \ (d n)$
$B_n, n \geq 2$	Z_2	$Spin_{2n+1}$	SO_{2n+1}	—
$C_n, n \geq 2$	Z_2	Sp_{2n}	$PCSp_{2n}$	—
$D_n, n \geq 3$ odd	Z_4	$Spin_{2n}$	PCO_{2n}°	SO_{2n}
$D_n, n \geq 4$ even	$Z_2 \times Z_2$	$Spin_{2n}$	PCO_{2n}°	$SO_{2n}, HSpin_{2n}$
G_2	1	G_2		—
F_4	1	F_4		—
E_6	Z_3	$(E_6)_{sc}$	$(E_6)_{ad}$	—
E_7	Z_2	$(E_7)_{sc}$	$(E_7)_{ad}$	—
E_8	1	E_8		—