Eine Woche, ein Beispiel 9.5 vector bundle v.s. Local system

Key objects in Geometry & Algebra.

vector bundle over manifoldmodule over ring

There are hundreds of different versions of it.

— vector bundle over manifold not not to the differential v.b. over (real) differential mfld Riemann surface · cplx (analytic) line bundle over Riemann surface

— sheaf over space 代数几何

scheme theory · locally free sheaf on scheme

· coherent sheaf on scheme

geo rep theory · local system over (real/cplx) mfld

· perverse sheaf over Riemann surface (derived)

— module over ring n数

comm alg . f.g module over Noetherian commutative ring (with 1)

rep of grp · group representation over group (~> group algebra)

quiver theory · quiver representation over quiver (~> path algebra, bound quiver algebra)

Lie algebra · Lie alg representation over Lie alg (~> universal enveloping algebra)

Goal - structures & invariants

- classifications of special v.b, mfld, subv.b, submfld

- symmetry & quotient

- special functors

- homological algebra, derived version

Today we will focus on the comparison between v.b. and local system.

1. classifications of real/cplx v.b. on S?.

(by homotopy group! ~> generalized Picard group?)

Q: Is this group structure natural?

ref: https://math.stackexchange.com/questions/1923402/understanding-vector-bundles-over-spheres

Frank m K-v.b. over S^n $\longleftrightarrow \pi_{n-1}(GL_m(K))$ Thm K=IR, C >6 5 7/27/ Z/2/2 7/2/2 2/2 2/2 7/27/ 2/12/ 2/2/2 74/2/ 7/27/ 7/27/ 2/22 \mathbb{Z} 0 0 0 Z Z \mathbb{Z} \mathbb{Z} (2/12)2 2/2/ 24/22 υ o 0 2/12 (2/12) 0 IRIP = K(2//2/1) Ta-(GLa(C)) rank >6 6 5 2 3 O 0 0 0 \mathbb{Z} \mathbb{Z} \mathbb{Z} \mathbb{Z} \mathbb{Z} \mathbb{Z} \mathbb{Z}

 $\mathbb{C}\mathbb{P}^{2} \cong \mathsf{K}(\mathbb{Z}_{2})$

Problems. Describe the special bundles, e.g. TS^n Describe the operations, e.g. dual, Θ , Q, Λ^k , Sym^k , Res, Ind

For the other spaces:

https://math.stackexchange.com/questions/383838/classifying-vector-bundles

http://www.ms.uky.edu/~guillou/F18/751Notes.pdf

It's still not so explicit.

Frank m K-v.b. over M ? (M, Grk(m, 00)] K=IR, C

K=IR, C

M: paracompact