

# Eine Woche, ein Beispiel

## 6.26. adic space

Ref: Berkeley notes by Peter Scholze.

Section 4, [http://www.math.uni-bonn.de/people/ja/lubintate/lecture\\_notes\\_lubin\\_tate.pdf](http://www.math.uni-bonn.de/people/ja/lubintate/lecture_notes_lubin_tate.pdf)

I just feel the need to record these results, so that I won't check them again as time went by.

Also I just learned a little about the discrete version. It is still not obvious for me to find out all valuations up to equivalence.

### 1. Discrete Huber pairs.

Set level

$(A, A^+)$ :  $A \in \text{CRing}$   $A^+ \leq$  integrally closed subring (containing 1)

$$\text{Spa}(A, A^+) = \left\{ v: A \longrightarrow \Gamma \cup \{+\infty\} \mid \begin{array}{l} v(f+g) \geq \min\{v(f), v(g)\} \\ v(fg) = v(f) + v(g) \\ v(A^+) \geq 0 \end{array} \right\} \sim$$

$$= \left\{ |\cdot|: A \longrightarrow \Gamma' \cup \{0\} \mid \begin{array}{l} |f+g| \leq \max\{|f|, |g|\} \\ |fg| = |f||g| \\ |A^+| \leq 1 \end{array} \right\} \sim$$

$$v \sim v' \Leftrightarrow [v(a) \geq v(b) \Leftrightarrow v'(a) \geq v'(b)]$$

$$|\cdot| \sim |\cdot|' \Leftrightarrow [|a| \leq |b| \Leftrightarrow |a|' \leq |b|']$$


$(\Gamma, +, 0)$ ,  $(\Gamma', \cdot, 1)$ : totally ordered abelian group

We use the same notations as in the document [Berkovich space, 2021.11.7].


E.g.  $\text{Spa}(\mathbb{Z}, \mathbb{Z}) = \{|\cdot|_{\text{triv}}, |\cdot|_p, |\cdot|_{\mathbb{F}_p}\}$

$\text{Spa}(\mathbb{Q}, \mathbb{Z}) = \{|\cdot|_{\text{triv}}, |\cdot|_p\}$

$\text{Spa}(\mathbb{Q}, \mathbb{Q}) = \{|\cdot|_{\text{triv}}\}$



supp:  $\text{Spa}(\mathbb{Z}, \mathbb{Z}) \longrightarrow \text{Spec } \mathbb{Z}$   
 $k = \text{supp}$ : kernel map (contraction)



## Topology level.

Def (Rational open subsets of  $\text{Spa}(A, A^+)$ ) For  $f, g_1, \dots, g_n \in A$ ,

$$\begin{aligned} \mathcal{U}\left(\frac{g_1, \dots, g_n}{f}\right) &= \{v \mid v(g_i) \geq v(f) \neq +\infty \quad \forall i\} \\ &= \{v \mid |g_i| \leq |f| \neq 0 \quad \forall i\} \\ &= \mathcal{U}\left(\frac{g_1}{f}\right) \cap \mathcal{U}\left(\frac{g_2}{f}\right) \cap \dots \cap \mathcal{U}\left(\frac{g_n}{f}\right) \end{aligned}$$

E.g. For  $\text{Spa}(\mathbb{Z}, \mathbb{Z})$ , Yellow: remove

$$\mathcal{U}\left(\frac{0}{1}\right) = \mathcal{U}\left(\frac{9}{1}\right) = \mathcal{U}\left(\frac{3}{1}\right) = \mathcal{U}\left(\frac{1}{1}\right) = \text{Spa}(\mathbb{Z}, \mathbb{Z})$$

$$\mathcal{U}\left(\frac{0}{3}\right) = \mathcal{U}\left(\frac{9}{3}\right) = \mathcal{U}\left(\frac{3}{3}\right) = \{ \text{valuations with } v(3) < v(0) \} \quad \mathcal{U}\left(\frac{1}{3}\right) = \{ \text{valuations with } v(3) < v(1) \} = \mathcal{U}\left(\frac{1}{9}\right)$$

$$\mathcal{U}\left(\frac{45}{36}\right) = \{ \text{valuations with } v(45) \geq v(36) \} \quad \mathcal{U}\left(\frac{0}{0}\right) = \emptyset$$

For  $\text{Spa}(\mathbb{Q}, \mathbb{Z})$ ,

$$\mathcal{U}\left(\frac{0}{1}\right) = \mathcal{U}\left(\frac{9}{1}\right) = \mathcal{U}\left(\frac{3}{1}\right) = \mathcal{U}\left(\frac{1}{1}\right) = \text{Spa}(\mathbb{Q}, \mathbb{Z})$$

$$\mathcal{U}\left(\frac{0}{3}\right) = \mathcal{U}\left(\frac{9}{3}\right) = \mathcal{U}\left(\frac{3}{3}\right) = \text{Spa}(\mathbb{Q}, \mathbb{Z}) \quad \mathcal{U}\left(\frac{1}{3}\right) = \{ \text{valuations with } v(3) < v(1) \} = \mathcal{U}\left(\frac{1}{9}\right)$$

$$\mathcal{U}\left(\frac{45}{36}\right) = \{ \text{valuations with } v(45) \geq v(36) \} \quad \mathcal{U}\left(\frac{0}{0}\right) = \emptyset$$

$$\mathcal{U}\left(\frac{1}{9}\right) = \{ \text{valuations with } v(9) < v(1) \} \quad \mathcal{U}\left(\frac{1}{9}\right) = \text{Spa}(\mathbb{Q}, \mathbb{Z}) \quad \mathcal{U}\left(\frac{1}{9}\right) = \text{Spa}(\mathbb{Q}, \mathbb{Z})$$

It is actually easier in this case, we just take the reduction of a fraction, and remove valuations which correspond to primes on the denominators.

Rmk. Rational open subsets of  $\text{Spa}(A, A^+)$  is the basis of a topo, as

$$\mathcal{U}\left(\frac{g_1, \dots, g_n}{f}\right) \cap \mathcal{U}\left(\frac{g'_1, \dots, g'_m}{f'}\right) = \mathcal{U}\left(\frac{g_1 f', \dots, g_n f', g'_1 f, \dots, g'_m f}{ff'}\right)$$

Now we can do the same formalism as two years ago.

- Topology

• Closed subset of  $\text{Spa}(\mathbb{Z}, \mathbb{Z})$ :  $\bigcup_{p \in \Delta_1} \{1 \cdot |_p, 1 \cdot |_{\mathbb{F}_p}\} \cup \bigcup_{p \in \Delta_2} \{1 \cdot |_{\mathbb{F}_p}\}$ , or  $\text{Spa}(\mathbb{Z}, \mathbb{Z})$

$\Delta_1, \Delta_2 \subset \{\text{primes } p\}$  finite,  $\Delta_1 \cap \Delta_2 = \emptyset$

• Open subset of  $\text{Spa}(\mathbb{Z}, \mathbb{Z})$ :  $\text{Spa}(\mathbb{Z}, \mathbb{Z}) - \bigcup_{p \in \Delta_1} \{1 \cdot |_p, 1 \cdot |_{\mathbb{F}_p}\} \cup \bigcup_{p \in \Delta_2} \{1 \cdot |_{\mathbb{F}_p}\}$ , or  $\emptyset$

• closure of a pt:

$$\overline{\{1 \cdot |_{\text{triv}}\}} = \text{Spa}(\mathbb{Z}, \mathbb{Z}) \quad \overline{\{1 \cdot |_p\}} = \{1 \cdot |_p, 1 \cdot |_{\mathbb{F}_p}\} \quad \overline{\{1 \cdot |_{\mathbb{F}_p}\}} = \{1 \cdot |_{\mathbb{F}_p}\}$$

• Topo properties: connected  $\checkmark$  Hausdorff  $\times$  quasi-compact  $\checkmark$   
irreducible  $\checkmark$

$$A^{\circ\circ} \subset A^{\circ} \subset A$$

$$U \supset A^+ \subset U$$

$$I \subset A_{\circ}$$