

Eine Woche, ein Beispiel

5.12 sheaf version of \otimes & Hom

Ref: from [23.11.19]

1. def of sheaf Hom
2. def of sheaf \otimes

sheaf version of Tensor-Hom adjunction is left in the next document.

Compared with \otimes , Hom is more delicate, and it is harder than you expected.

1. def of sheaf Hom

$$\begin{array}{lcl}
 \text{Hom}_A(-, -): (A\text{-Mod})^{\text{op}} \times A\text{-Mod} & \longrightarrow & A\text{-Mod} \quad A: \text{comm ring} \\
 \downarrow \\
 \text{Hom}_{\text{Sh}(X)}(-, -): \text{Sh}(X)^{\text{op}} \times \text{Sh}(X) & \longrightarrow & \mathbb{Z}\text{-Mod} \\
 \downarrow \\
 \underline{\text{Hom}}_{\text{Sh}(X)}(-, -): \text{Sh}(X)^{\text{op}} \times \text{Sh}(X) & \longrightarrow & \text{Sh}(X) \\
 \downarrow \\
 R\underline{\text{Hom}}_{\mathcal{D}^+(X)}(-, -): \mathcal{D}^+(X)^{\text{op}} \times \mathcal{D}^+(X) & \longrightarrow & \mathcal{D}^+(X)
 \end{array}$$

Non-derived Hom

Def [FOAG, 2.3.1] For $\mathcal{F}, \mathcal{G} \in \text{Sh}(X)$, a morphism of sheave
 $\phi: \mathcal{F} \rightarrow \mathcal{G}$

is the data of maps

$\phi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ for all $U \subseteq X$ open.
 which is compatible with restriction.

We write

$$\phi \in \text{Hom}_{\text{Sh}(X)}(\mathcal{F}, \mathcal{G})$$

Similarly, one can define

$\text{Hom}_{\mathcal{D}(X)}(\mathcal{F}', \mathcal{G}')$
 as the set of morphisms in $\mathcal{D}(X)$.

Def [FOAG, 2.3.C] (Sheaf Hom / Internal Hom)

For $\mathcal{F}, \mathcal{G} \in \text{Sh}(X)$, one gets a sheaf $\underline{\text{Hom}}(\mathcal{F}, \mathcal{G}) \in \text{Sh}(X)$ given by

$$(\underline{\text{Hom}}(\mathcal{F}, \mathcal{G}))(U) = \text{Hom}_{\text{Sh}(U)}(\mathcal{F}|_U, \mathcal{G}|_U)$$

Cor.

$$\text{Hom} = \Gamma \circ \underline{\text{Hom}} : \text{Sh}(X)^{\text{op}} \times \text{Sh}(X) \xrightarrow{\underline{\text{Hom}}} \text{Sh}(X) \xrightarrow{\Gamma} \text{Abel}$$



Even though $(\mathcal{F} \otimes \mathcal{G})_p \cong \mathcal{F}_p \otimes \mathcal{G}_p$,

$\underline{\text{Hom}}$ does not commute with taking stalks.

$$(\underline{\text{Hom}}(\mathcal{F}, \mathcal{G}))_p \xrightarrow{\neq} \text{Hom}(\mathcal{F}_p, \mathcal{G}_p)$$

It's neither inj nor surj.

[Left adj comm with limit, $\otimes \dashv \underline{\text{Hom}}$].

Ex. Try to compute coefficient \mathbb{Q} .

$$\begin{aligned} \underline{\text{Hom}}_{\text{Sh}(X)}(\underline{\mathbb{Q}}_X, \mathcal{F}) &\cong \mathcal{F} \\ \underline{\text{Hom}}_{\text{Sh}(X)}(j_* \underline{\mathbb{Q}}_U, \mathcal{F}) &\cong j_* (\mathcal{F}|_U) \\ \underline{\text{Hom}}_{\text{Sh}(\mathbb{C})}(\text{sky}_0(\mathbb{Q}), \underline{\mathbb{Q}}_{\mathbb{C}}) &\cong 0 \end{aligned}$$