


Eine Woche, ein Beispiel
1.28 conormal bundle

Ref: from [23.11.19]

slogan:

SES	induces	LES,
filtration	induces	spectral sequence.

To expand a little bit,



Even though "filtration \Rightarrow spectral seq" is the most general statement, people start with "SES \Rightarrow LES" and "acyclic resolution \Rightarrow other coh \approx hyper coh". Let us leave spectral seq in other people's notes.

Methods to construct SES: $\left\{ \begin{array}{l} \text{check by stalks} \\ \text{filtration by } H^i(\mathcal{F}) \\ \text{filtration by } \mathcal{F}^i \end{array} \right.$

method	spectral seq	LES	cohomology/resolution
check by stalks	... for stratifications	relative coh seq	simplicial/cellular
	Čech-to-derived fctor	MV	Čech
filtration by $H^i(\mathcal{F})$	Grothendieck		
	Leray-Serre	Cysin	Euler class
			Hodge-Tate
filtration by \mathcal{F}^i need resolution to get "another" complex	Hodge-de Rham		de Rham, Hodge-de Rham
			Dolbeault $H^p(X, \Omega^q) = H^{p,q}(X)$
	Frölicher		$H^{p,q}(X) \Rightarrow H^{p+q}(X)$ "composition"
			singular
spectral sequences which I don't know	Adams Atiyah-Hirzebruch Bar Bockstein Cartan-Leray Eilenberg-Moore Green ⋮		for stable homotopy gp for top K-theory for group for group homology for Koszul cohomology ⋮

For more spectral sequences, see:

https://en.wikipedia.org/wiki/Spectral_sequence

<https://github.com/CubicBear/SpectralSequences/blob/main/SpectralSequences.pdf>

1. open-closed formalism
2. open cover
3. filtration by $H^i(\mathcal{F})$
4. Hodge related filtration

1. open-closed formalism

|| related: comparison of $j_!$ & j_*
one-point compactification.

Observe the following pictures:

$$\begin{array}{ccccc} Z & \xrightarrow{i} & X & \xleftarrow{j} & U \\ & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{D}(Z) & \xrightarrow{i_* = i_!} & \mathcal{D}(X) & \xrightarrow{j^* = j^!} & \mathcal{D}(U) \\ & \xleftarrow{i^!} & & \xleftarrow{Rj_*} & \end{array}$$

Black box:

0. We assume some nice conditions.

e.g. in the category $\text{Haus}^{\text{loc. cpt.}}$, and $Z \subset X$ is loc. contractible.

Under these conditions,

1. $i_* = i_!$, $j^* = j^!$
2. $j_!$, i^* , j^* , i_* are exact.

Ex. 1. Shows that

$$\underline{i^* i_*} = \underline{i^! i_*} = \text{Id}_{\mathcal{D}(Z)} \quad \underline{j^* j_!} = \underline{j^* Rj_*} = \text{Id}_{\mathcal{D}(U)}$$

$$\underline{i^* j_!} = 0, \quad \underline{j^* i_*} = 0, \quad \underline{i^! Rj_*} = 0$$

— : base change

~~~~~ : check stalkwise.

2. (for category fans)

$i_*$ ,  $j_*$ ,  $j_!$  are fully faithful, and  
 $i_*$ ,  $i^!$ ,  $j^*$ ,  $Rj_*$  preserve injectives.

3. One has SES

$$0 \longrightarrow j_! j^! \mathcal{F} \longrightarrow \mathcal{F} \longrightarrow i_* i^* \mathcal{F} \longrightarrow 0 \quad (1)$$

Ex for (1).

1. Apply the  $R\pi_{X,*}$  to (1), take  $\mathcal{F} = \underline{\mathbb{Q}}_X$ , what do we get?

In general, what do we get when applying  $R\pi_{X,*}$  &  $R\pi_{X,!}$ ?

Discuss 2 spectral cases  $\mathcal{F} = \underline{\mathbb{Q}}_X$   $\text{ID}_X := \pi_{X,!} \underline{\mathbb{Q}}_{f^{-1}Y} = \text{ID}_X(\underline{\mathbb{Q}}_X)$

2. Derive from (1) the SES

$$0 \longrightarrow j_! \mathcal{F} \longrightarrow Rj_* \mathcal{F} \longrightarrow i_* i^* Rj_* \mathcal{F} \longrightarrow 0$$

which measures the difference between  $j_! \mathcal{F}$  &  $j_* \mathcal{F}$ .

3. Shows that

$$H_c(X) \cong H(\bar{X}, \{\infty\}; \mathbb{Z})$$

for one pt compactification  $i: X \hookrightarrow \bar{X}$ .

Try to compute  $H_c(\mathbb{R}^n)$  in this way.

It seems that we get only half of the results.

### Verdier dual

Def. The Verdier dual / dualizing functor is defined as

$$ID_X: D^b(X; \mathbb{Q}) \longrightarrow D^b(X; \mathbb{Q}) \quad ID_X \mathcal{F}^\bullet := \underline{\text{Hom}}_{D^b(X; \mathbb{Q})}(\mathcal{F}^\bullet, \pi_X^! \underline{\mathbb{Q}}_{\{*\}})$$

We know that

$$ID_X \underline{\mathbb{Q}}_X = \pi_X^! \underline{\mathbb{Q}}_{\{*\}}$$

$$ID_X(\mathcal{F}[n]) = (ID_X \mathcal{F}^\bullet)[-n]$$

$$\mathcal{F}^\bullet \longrightarrow \mathcal{G}^\bullet \longrightarrow \mathcal{H}^\bullet \xrightarrow{+1} \rightsquigarrow ID \mathcal{H}^\bullet \longrightarrow ID \mathcal{G}^\bullet \longrightarrow ID \mathcal{F}^\bullet \xrightarrow{+1}$$

$$f^! ID_X = ID_Y f^*$$

$$Rf_* ID_Y = ID_X Rf_!$$

$$f: Y \longrightarrow X$$

When  $\mathcal{F}^\bullet \in D^b(X; \mathbb{Q})$  is constructible, then

$$ID_X^2 \mathcal{F}^\bullet \cong \mathcal{F}^\bullet$$

Therefore, in the constructible setting,

$$f^* ID_X = ID_Y f^!$$

$$Rf_! ID_Y = ID_X Rf_*$$

For exact statements about  $ID_X$ , see [MS21, Cor 2.11] [IHPS, Thm 5.3.9]


Ex. Derive from (1) the triangle

$$i_! i^! \mathcal{F} \longrightarrow \mathcal{F} \longrightarrow Rj_* j^* \mathcal{F} \xrightarrow{+1} \quad (2)$$

for  $\mathcal{F}^\bullet \in D^b(X; \mathbb{Q})$  constructible.

Ex for (2). Do the same arguments in "Ex for (1)".

E.g. For a finite graph (as a topo space)  $X$ .



$$0 \longrightarrow j_! j^! \underline{\mathbb{Q}}_X \longrightarrow \underline{\mathbb{Q}}_X \longrightarrow i_* i^* \underline{\mathbb{Q}}_X \longrightarrow 0$$

$$0 \longrightarrow j_! \underline{\mathbb{Q}}_{X - sk_0 X} \longrightarrow \underline{\mathbb{Q}}_X \longrightarrow i_! \underline{\mathbb{Q}}_{sk_0 X} \longrightarrow 0$$

Take  $R\pi_{X,!}$ :

$$\begin{array}{c}
 \xrightarrow{\quad} H_c^1(X - sk_0 X) \overset{= \oplus \mathbb{Q}}{\longrightarrow} H_c^1(X) \longrightarrow H_c^1(sk_0 X) \overset{= 0}{\xrightarrow{+1}} \\
 \searrow \hspace{10em} \nearrow \\
 0 \longrightarrow H_c^0(X - sk_0 X) \overset{= 0}{\longrightarrow} H_c^0(X) \longrightarrow H_c^0(sk_0 X) \overset{= \oplus \mathbb{Q}}{\longrightarrow}
 \end{array}$$

This calculates the sheaf cohomology as simplicial cohomology.

E.x. Shows that

$$H_c^i(\mathbb{R}) = \begin{cases} \mathbb{Q} & i=1 \\ 0 & \text{otherwise} \end{cases}$$

in a similar way.

Generalizing this argument, one can relate sheaf cohomology with simplicial/cellular cohomology, using the following filtration:

$$0 \subset sk^0 X \subset sk^1 X \subset \cdots \subset sk^n X = X$$

Ex. derive the Wang LES for the cpt supp version. over  $S'$

2. open cover

Ex. For an open cover  $X = U_1 \cup U_2$ , deduce the SES

$$\begin{array}{ccccccc} 0 & \longleftarrow & \mathcal{Q}_X & \longleftarrow & j_! \mathcal{Q}_{U_1} \oplus j_! \mathcal{Q}_{U_2} & \longleftarrow & j_! \mathcal{Q}_{U_1 \cap U_2} \longleftarrow 0 \\ & & \mathcal{Q}_X & \longrightarrow & Rj_* \mathcal{Q}_{U_1} \oplus Rj_* \mathcal{Q}_{U_2} & \longrightarrow & Rj_* \mathcal{Q}_{U_1 \cap U_2} \xrightarrow{+1} \end{array} \quad (3)$$

▽ We omit the derived symbol and some subscripts in this section.  $U_{12} = U_1 \cap U_2$

(3) works for general sheaf

and, induce from (3) the MV sequence:

$$\begin{array}{ccccccc} \xrightarrow{+1} & H_c^k(X) & \longleftarrow & H_c^k(U_1) \oplus H_c^k(U_2) & \longleftarrow & H_c^k(U_1 \cap U_2) \\ & H^k(X) & \longrightarrow & H^k(U_1) \oplus H^k(U_2) & \longrightarrow & H^k(U_1 \cap U_2) \xrightarrow{+1} \end{array}$$

Hint: Apply  $R\pi_{X,!}$  &  $R\pi_{X,*}$ , see [StackProject, 01E9]

Ex. Derived the Wang LES. over  $S^1$

Ex. For an open cover  $X = \bigcup_{i \in \Lambda} U_i$ ,  $\Lambda$  finite, deduce the exact seq

$$0 \longleftarrow \mathcal{Q}_X \longleftarrow \bigoplus_i j_! \mathcal{Q}_{U_i} \longleftarrow \bigoplus_{i < j} j_! \mathcal{Q}_{U_i \cap U_j} \longleftarrow \cdots \longleftarrow j_! \mathcal{Q}_{\bigcap_i U_i} \longleftarrow 0$$

and t-exact seq

$$0 \longrightarrow \mathcal{Q}_X \longrightarrow \bigoplus_i Rj_* \mathcal{Q}_{U_i} \longrightarrow \bigoplus_{i < j} Rj_* \mathcal{Q}_{U_i \cap U_j} \longrightarrow \cdots \longrightarrow Rj_* \mathcal{Q}_{\bigcap_i U_i} \longrightarrow 0$$

When  $\{U_i\}_{i \in \Lambda}$  is a good cover,  $H^i(U_{i_1, \dots, i_k}) = H^0(U_{i_1, \dots, i_k})$ ,  
 $\uparrow$  acyclic in AG

one can compute  $H^i(X)$  by applying  $R\pi_{X,*}$ :

$$\begin{array}{ccccccc} 0 \longrightarrow & \bigoplus_i \Gamma(U_i) & \xrightarrow{d^1} & \bigoplus_{i < j} \Gamma(U_i \cap U_j) & \xrightarrow{d^2} & \cdots & \Gamma(\bigcap_i U_i) \longrightarrow 0 \\ & & & \downarrow \text{Ker}/I_m & & & \\ & H^0(X) & & H^1(X) & & \cdots & H^{\#\Lambda-1}(X) \end{array}$$

Rmk. When  $X$  is paracompact & Hausdorff, "limited" Čech = sheaf  
 $\uparrow$  e.g. loc cpt Haus + second-countable, or CW cplx

compare the first step:

$$\mathcal{F} \longrightarrow \bigoplus_i Rj_* \mathcal{F}|_{U_i}$$

$$\mathcal{F} \longrightarrow \bigoplus_{x \in X} \mathcal{F}_x$$

If you haven't seen the acyclic resolution before, the following example may provide some intuition.

#  $\Delta = 3$  case:

$$\begin{array}{ccccccc}
 & & \begin{array}{c} 0 \searrow \\ \mathcal{F}_0 \\ 0 \nearrow \end{array} & & & & \begin{array}{c} 0 \searrow \\ \mathcal{F}_2 \\ 0 \nearrow \end{array} \\
 & & \nearrow & & & & \nearrow \\
 0 \longrightarrow \mathcal{Q}_X & \longrightarrow & \bigoplus_i R_{j*} \mathcal{Q}_{U_i} & \xrightarrow{d'} & \bigoplus_i R_{j*} \mathcal{Q}_{U_i \cap U_j} & \xrightarrow{d^2} & R_{j*} \mathcal{Q}_{\cap U_i} \xrightarrow{d^3} 0 \\
 & & \searrow & & & & \searrow \\
 & & \begin{array}{c} 0 \nearrow \\ \mathcal{F}_1 \\ 0 \searrow \end{array} & & & & 
 \end{array}$$

$$\mathcal{F}_2 = R_{j*} \mathcal{Q}_{\cap U_i} \Rightarrow H^i(\mathcal{F}_2) = \ker d^3$$

$$\begin{array}{c}
 \hookrightarrow H^1(\mathcal{F}_1) \longrightarrow 0 \longrightarrow H^1(\mathcal{F}_2) \xrightarrow{+1} \\
 \hookrightarrow H^0(\mathcal{F}_1) \longrightarrow \bigoplus_{i,j} \Gamma(U_i \cap U_j) \xrightarrow{d^2} H^0(\mathcal{F}_2)
 \end{array}$$

$$\Rightarrow H^i(\mathcal{F}_1) = \begin{cases} \ker d^3 / \text{Im } d^2, & i=1 \\ \ker d^2, & i=0 \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{array}{c}
 \hookrightarrow H^2(\mathcal{F}_0) \longrightarrow 0 \longrightarrow H^2(\mathcal{F}_1) \xrightarrow{+1} \\
 \hookrightarrow H^1(\mathcal{F}_0) \longrightarrow 0 \longrightarrow H^1(\mathcal{F}_1) \\
 \hookrightarrow H^0(\mathcal{F}_0) \longrightarrow \bigoplus_i \Gamma(U_i) \xrightarrow{d^1} H^0(\mathcal{F}_1)
 \end{array}$$

$$\Rightarrow H^i(X) = H^i(\mathcal{F}_0) = \begin{cases} \ker d^3 / \text{Im } d^2 & i=2 \\ \ker d^2 / \text{Im } d^1 & i=1 \\ \ker d^1 & i=0 \\ 0, & \text{otherwise} \end{cases}$$

Rmk. When  $\{U_i\}_{i \in \Delta}$  is not a good cover,  
one needs Čech-to-derived functor spectral seq to compute  $H^i(X)$ .



Rmk. stratification & open cover are two main tools to extract topological information.  
 They appear with different names in different fields.  
 Once you realize them, you can apply the six-functor machine to analyze structures.

stratification with extra properties  $\left\{ \begin{array}{l} \text{CW cplx} \\ \text{triangulization} \\ \text{dessin d'enfant} \\ \text{affine paving} \\ \text{Whitney stratification} \\ \vdots \end{array} \right.$

Q: How to construct stratifications?

A: For me, there are two efficient methods:  $\left\{ \begin{array}{l} \text{orbit of gp action} \\ \text{Morse theory} \end{array} \right.$

That's why some geometrical problems are finally reduced to combinatorial / analytic problems.  
 Other fields come to geometry by providing stratifications.

In fact, there is only one method:

find a fct  $f: X \rightarrow Y$ , and get stratification of  $X$  from  $Y$ .

- E.g.
1. Morse theory
  2. tessellation
  3. semi-continuous fct
  4. my thesis
  5. orbit of gp action

$$f: X \rightarrow \mathbb{R}$$

$$f: \mathcal{H} \rightarrow \mathcal{H}/\Gamma$$

$$f: X \rightarrow \mathbb{N}_{\geq 0} \quad \text{e.g. } f(p) = \dim T_p X$$

$$f: \text{Gr}(X) \rightarrow \text{Gr}(S) \times \text{Gr}(X/S)$$

$$f: X \rightarrow X/G$$