

Eine Woche, ein Beispiel
11.26 calculation of double point

Final goal: Fill in the tables in the next page.
(for first time, remove the $i!$ column)

Ref:

[Willians]: Langlands correspondence and Bezrukavnikov's equivalence

calculations from Lukas Bonfert's note (don't forward this to anyone else).

$$X = \mathbb{C} \cup_{\{0\}} \mathbb{C} = \{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 z_2 = 0\}, \quad Z = \{0\}, \quad U = \mathbb{C}^\times \cup \mathbb{C}^\times$$

$$i_* \mathbb{Q}_Z$$

(0, 1, 1, 1)

	n	-2	-1	0	1
\mathcal{U}	j^*	0	0	0	0
$\{0\}$	i^*	0	0	\mathbb{Q}	0
	$i'!$	0	0	\mathbb{Q}	0
	$R^n \Gamma$	0	0	\mathbb{Q}	0

$$\mathbb{Q}_X[1]$$

(-1, -1, -1, -1)

	n	-2	-1	0	1
\mathcal{U}	j^*	0	\mathbb{Q}	0	0
$\{0\}$	i^*	0	\mathbb{Q}	0	0
	$i'!$	0	0	\mathbb{Q}	\mathbb{Q}^2
	$R^n \Gamma$	0	\mathbb{Q}	0	0

perverse sheaf
IC sheaf

? by dim argument

	n	-2	-1	0	1
\mathcal{U}	j^*	X		X	X
$\{0\}$	i^*	—		X	X
	$i'!$	X	X	X	
	$R^n \Gamma$	—			

$$Rj_* \mathbb{Q}_U[1]$$

(-1, 0, 0, 0)

	n	-2	-1	0	1
\mathcal{U}	j^*	0	\mathbb{Q}	0	0
$\{0\}$	i^*	0	\mathbb{Q}^2	\mathbb{Q}^2	0
	$i'!$	0	0	0	0
	$R^n \Gamma$	0	\mathbb{Q}^2	\mathbb{Q}^2	0
	Γ	0	\mathbb{Q}^2	\mathbb{Q}	0

$$\Gamma^{(n)} = \Gamma(Rj_* \mathbb{Q}_U[1])$$

$$j! \mathbb{Q}_U[1]$$

(-1, 0, 0, 0)

	n	-2	-1	0	1
\mathcal{U}	j^*	0	\mathbb{Q}	0	0
$\{0\}$	i^*	0	0	0	0
	$i'!$	0	0	\mathbb{Q}^2	\mathbb{Q}^2
	$R^n \Gamma$	0	0	0	0

$$\pi^! \mathbb{Q}[-1]$$

(-1, -1, -1, -1)

	n	-2	-1	0	1
\mathcal{U}	j^*	0	\mathbb{Q}	0	0
$\{0\}$	i^*	0	\mathbb{Q}^2	\mathbb{Q}	0
	$i'!$	0	0	0	\mathbb{Q}
	$R^n \Gamma$	0	\mathbb{Q}	0	0

$$X = \mathbb{C}^2, \quad Z = \{0\}, \quad U = \mathbb{C}_3 - \{0\} \cong S^3 \times \mathbb{R}_{>0}$$

$$i_! \underline{\mathbb{Z}}_Z$$

(0, 1, 1, 1)

	n	-2	-1	0	1	2
\mathcal{U}	j^*	0	0	0	0	0
$\{0\}$	i^*	0	0	\mathbb{Z}	0	0
	$i^!$	0	0	\mathbb{Z}	0	0
	$R^n \Gamma$	0	0	\mathbb{Z}	0	0

$$\mathcal{H}^\circ(\mathcal{F}) \in \text{Perv}_\Lambda(X)$$

$$\underline{\mathbb{Z}}_X[2]$$

(1, 1, 1, 1)

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$Rj_* \underline{\mathbb{Z}}_U[2] \in {}^p\mathcal{D}^{\geq 0}(X) - {}^p\mathcal{D}^{\leq 0}(X)$$

(1, 0, 0, 0)

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	\mathbb{Z}	0
	$i^!$	0	0	0	0	0
	$R^n \Gamma$	\mathbb{Z}	0	0	\mathbb{Z}	0

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$j_! \underline{\mathbb{Z}}_U[2] \in {}^p\mathcal{D}^{\leq 0}(X) - {}^p\mathcal{D}^{\geq 0}(X)$$

(1, 0, 0, 0)

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	0	0	0	0	0
	$i^!$	0	\mathbb{Z}	0	0	\mathbb{Z}
	$R^n \Gamma$	0	0	0	0	0

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$\pi^! \underline{\mathbb{Z}}[-2]$$

(1, 1, 1, 1)

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$X = X_3 = \{(z_1, z_2, z_3) \in \mathbb{C}^3 \mid z_1^2 + z_2^2 + z_3^2 = 0\}, \quad Z = \{0\}, \quad \mathcal{U} = X_3 - \{0\} = \mathbb{C}^x\text{-bd over } \mathbb{C}P^1$$

twisted

$$i_! \mathbb{Z}_Z$$

$$(0, 1, 1, 1)$$

	n	-2	-1	0	1	2
\mathcal{U}	j^*	0	0	0	0	0
$\{0\}$	i^*	0	0	\mathbb{Z}	0	0
	$i^!$	0	0	\mathbb{Z}	0	0
	$R^n \Gamma$	0	0	\mathbb{Z}	0	0

$$\mathcal{H}^0(\mathcal{F}) \in \text{Perv}_\Lambda(X)$$

$$\mathbb{Z}_X[2]$$

$$(1, 1, 1, 1)$$

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	$\mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$Rj_* \mathbb{Z}_\mathcal{U}[2] \in {}^p\mathcal{D}^{\geq 0}(X) - {}^p\mathcal{D}^{\leq 0}(X)$$

$$(1, 0, 0, 0)$$

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}	0
	$i^!$	0	0	0	0	0
	$R^n \Gamma$	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}	0

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	0	0

$$j_! \mathbb{Z}_\mathcal{U}[2] \in {}^p\mathcal{D}^{\leq 0}(X) - {}^p\mathcal{D}^{\geq 0}(X)$$

$$(1, 0, 0, 0)$$

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	0	0	0	0	0
	$i^!$	0	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}
	$R^n \Gamma$	0	0	0	0	0

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	0	0	0
	$i^!$	0	0	0	$\mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	0	0	0

$$\pi^! \mathbb{Z}[-2]$$

$$(1, 1, 1, 1)$$

	n	-2	-1	0	1	2
\mathcal{U}	j^*	\mathbb{Z}	0	0	0	0
$\{0\}$	i^*	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	0	0
	$i^!$	0	0	0	0	\mathbb{Z}
	$R^n \Gamma$	\mathbb{Z}	0	$\mathbb{Z}/2\mathbb{Z}$	0	0

perverse sheaf
IC sheaf

	n	-2	-1	0	1	2
\mathcal{U}	j^*		x	x	x	x
$\{0\}$	i^*			x	x	x
	$i'!$	x	x	x		
	$R^n \Gamma$					

Conclusion. \subseteq IC \subseteq Perv \subseteq Constructable

Q-coefficient $i_+ \underline{\mathbb{Q}}_{\mathbb{Z}}$
 $\underline{\mathbb{Q}}_X[2] \cong \pi'_! \underline{\mathbb{Q}}[-2]$

$Rj_* \underline{\mathbb{Q}}_u[2], j'_! \underline{\mathbb{Q}}_u[2]$

Z-coefficient $i_+ \underline{\mathbb{Z}}_{\mathbb{Z}}$
 $\underline{\mathbb{Z}}_X[2]$

$Rj_* \underline{\mathbb{Z}}_u[2], j'_! \underline{\mathbb{Z}}_u[2]$
 $\pi'_! \underline{\mathbb{Z}}[-2]$