

Eine Woche, ein Beispiel

8.6. Kottwitz set

This document is a continuation of [23.08.06].
Reorganized from Luozi Shi (and his partners)'s talk.

Recall that $\widehat{\mathbb{Q}_p^{ur}} = \text{Frac}(\widehat{\mathbb{Z}_p^{ur}})$, $\widehat{\mathbb{Z}_p^{ur}} = W(\mathbb{F}_p)$. Here " $\widehat{}$ " is completion w.r.t. valuation.

Setting. In this document, F is a NA local field,

$$L := \widehat{F^{ur}} \stackrel{\text{if } F: p\text{-adic}}{=} \text{Frac}(W(\bar{\kappa}_F) \otimes_{W(\kappa_F)} \mathcal{O}_F).$$

Def For G/F reductive, the Kottwitz set $B(G)$ is defined as

$$\begin{aligned} B(G) &:= H^1(W_F, G_{\bar{F}}) \\ &\cong H^1(\langle \sigma \rangle, G_L) && \text{by Inf-Res seq \& } H^1(I_F, G_{\bar{F}}) = 0 \\ &\cong G(L)/\sigma\text{-twisted } G(L)\text{-conj} \\ &\stackrel{\text{when } G = GL_n}{\cong} \text{Isoc}/\sim \\ &\quad \quad \quad \uparrow \\ &\quad \quad \quad \text{Isocrystals.} \end{aligned}$$

E.g. $B(G_m) \cong \mathbb{Z}$

Proof. The map

$$\beta: \mathcal{O}_L^\times \rightarrow \mathcal{O}_L^\times \quad x \mapsto x \cdot \sigma(x)^{-1}$$

is surjective, since

$$\beta_1: \kappa_L^\times \rightarrow \kappa_L^\times \quad x \mapsto x^{1-q}$$

$$\beta_2: \kappa_L \rightarrow \kappa_L \quad x \mapsto x - x^q$$

are surjective. ($\kappa_L \cong \bar{\mathbb{F}_p}$ is alg closed)

Then the well-defined morphism

$$v: L^\times / \text{Im } \beta \rightarrow \mathbb{Z} \quad x \mapsto v(x)$$

is injective, thus an iso.

Ex. Check that the SES

$$1 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow G_m \xrightarrow{(-)^2} G_m \rightarrow 1$$

induce LES in gp cohomology:

$$\begin{array}{c} \hookrightarrow B(\mathbb{Z}/2\mathbb{Z}) \rightarrow B(G_m) \xrightarrow[\substack{x_1? \\ x_2?}]{x_1?} B(G_m) \xrightarrow{x_2?} B(G_m) \\ \searrow \hspace{10em} \nearrow \\ 0 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow F^\times \xrightarrow{(-)^2} F^\times \end{array}$$

where

$$\begin{aligned} B(\mathbb{Z}/2\mathbb{Z}) &\stackrel{\text{def}}{=} H^1(W_F, \mathbb{Z}/2\mathbb{Z}) & W_F \subset \mathbb{Z}/2\mathbb{Z} \text{ trivially} \\ &= \text{Hom}(W_F, \mathbb{Z}/2\mathbb{Z}) \\ &\cong \{ H \triangleleft W_F \text{ closed with index } 2 \} \cup \{0\} \\ &\cong \begin{cases} \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & p \neq 2 \\ \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & p = 2 \end{cases} \end{aligned}$$