

Eine Woche, ein Beispiel

12.1 weights of type E

It feels incomplete to discuss only the type E case without addressing the other classical cases.

Hence, this document serves as a complement to [2024.12.01].

▽ There are some new phenomena outside type E (which are not essential).

1. The formula becomes

$$2 \frac{\langle \varpi_i, \alpha_j \rangle}{\langle \alpha_j, \alpha_j \rangle} = \delta_{ij} \quad \text{i.e.,} \quad \langle \varpi_i \frac{2}{\langle \alpha_i, \alpha_i \rangle}, \alpha_j \rangle = \delta_{ij}$$

when $i \neq j$, $\frac{2}{\langle \alpha_i, \alpha_i \rangle}$ won't impact,
as $\delta_{ij} = 0$

$$s_k v = v - 2 \frac{\langle \alpha_k, v \rangle}{\langle \alpha_k, \alpha_k \rangle} \alpha_k$$

2. $A = (\langle \alpha_i, \alpha_j \rangle)_{i,j}$ is not Cartan matrix. It is $(2 \frac{\langle \alpha_i, \alpha_j \rangle}{\langle \alpha_i, \alpha_i \rangle})_{i,j}$.

3. The minuscule weight ^{orbit} may not generate the whole lattice
in type A, B, D

4. The minuscule weight ^{orbit} may not be the wts nearest to the origin
in type A, B, C, D

Since the coordinate itself already offers good symmetry (compared with type E case), we will omit many details.

1. A_n E.g. $n=5$

- Weights nearest to the origin

There are n many minuscule representations of A_n :

#	typical coordinates
$\binom{6}{1}$	$\frac{1}{6} (5, -1, -1, -1, -1)^T$
$\binom{6}{2}$	$\frac{1}{6} (4, 4, -2, -2, -2)^T$
$\binom{6}{3}$	$\frac{1}{6} (3, 3, 3, -3, -3)^T$
$\binom{6}{4}$	$\frac{1}{6} (2, 2, 2, 2, -4)^T$
$\binom{6}{5}$	$\frac{1}{6} (1, 1, 1, 1, -5)^T$

symbol

v_i	
v_{ij}	
v_{ijk}	
v_{ijkl}	
v_{ijklm}	

$$|v_i|^2 = \langle v_i, v_i \rangle \in \left\{ \frac{5}{6}, \frac{4}{3}, \frac{3}{2} \right\}$$

in general, in $\binom{n+1}{k}$, $\langle v_i, v_i \rangle = \frac{k(n+1-k)}{n+1}$


in $\left\{ \sum_{i=1}^{n+1} z_i = 0 \right\} \cong \mathbb{R}^n$

Restrict to the standard rep case,

$$\langle v_i, v_j \rangle \in \left\{ \frac{n}{n+1}, -1 \right\}.$$

The graph has no edges.

- Simple roots



$$\begin{aligned}
 & \{ \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5 \} \\
 &= \{ \nu_1 - \nu_2, \nu_2 - \nu_3, \nu_3 - \nu_4, \nu_4 - \nu_5, \nu_5 - \nu_6 \} \\
 &= \left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -1 \end{pmatrix} \right\}
 \end{aligned}$$

Ex. Verify that all the $z(\frac{n+1}{2}) = 30$ roots are given by

#	typical coordinates	symbol
30	$(1, -1, 0, 0, 0, 0)^T$	$\alpha_{1,2}$

- Fundamental weights

$$\begin{aligned}
 & \{ \omega_1, \omega_2, \omega_3, \omega_4, \omega_5 \} \\
 &= \{ \nu_1, \nu_{12}, \nu_{123}, \nu_{1234}, \nu_{12345} \} \\
 &= \left\{ \begin{pmatrix} 5 \\ -1 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 4 \\ 4 \\ -2 \\ -2 \\ -2 \\ -2 \end{pmatrix}, \begin{pmatrix} 3 \\ 3 \\ 3 \\ -3 \\ -3 \\ -3 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \\ 2 \\ -4 \\ -4 \\ -4 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ -5 \end{pmatrix} \right\} \\
 &= \left\{ \begin{pmatrix} 5 \\ -1 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ -2 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ -5 \end{pmatrix} \right\}
 \end{aligned}$$




- Weyl group action

S_{n+1} acts by permutations. Nothing special.

2. D_n E.g. $n=6$ $n > 4$ for avoiding special cases.

- Weights nearest to the origin

There are 3 minuscule representations of D_n :

	#	typical coordinates	symbol	
vector	$2 \cdot 6$	$(1, 0, 0, 0, 0, 0)^T$	v_i & $-v_i$	
	2^5	$\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1)^T$ odd (neg) sign	$v_{\pm \pm \pm \pm \pm}$	
half spin	2^5	$\frac{1}{2}(1, 1, 1, 1, 1, -1)^T$	v_-	
	2^5	$\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1)^T$ even (neg) sign	$v_{\pm \pm \pm \pm \pm \pm}$	
		$\frac{1}{2}(1, 1, 1, 1, 1, 1)^T$	v_+	

in general, $\langle v_i, v_i \rangle \in \{1, \frac{n}{4}\}$ in \mathbb{R}^n .


Restrict to the standard rep case,
 $\langle v_i, v_j \rangle \in \{1, 0, -1\}$.

The graph is \equiv .



Here, the weights corresponding to standard reps does not generate all other weights.

- Simple roots



$$\begin{aligned}
 & \{ \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \} \\
 &= \{ \nu_1 - \nu_2, \nu_2 - \nu_3, \nu_3 - \nu_4, \nu_4 - \nu_5, \nu_5 - \nu_6, \nu_5 + \nu_6 \} \\
 &= \left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 1 \end{pmatrix} \right\}
 \end{aligned}$$

Ex. Verify that all the $4 \binom{n}{2} = 60$ roots are given by

#	typical coordinates	symbol
60	$(\pm 1, \pm 1, 0, 0, 0, 0)^T$ $(-1, 1, 0, 0, 0, 0)^T$	$\alpha_{\pm i \pm j}$ α_{-1+2}

- Fundamental weights

$$\begin{aligned}
 & \{ \omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6 \} \\
 &= \{ \nu_1, \nu_1 + \nu_2, \sum_{i=1}^3 \nu_i, \sum_{i=1}^4 \nu_i, \nu_-, \nu_+ \} \\
 &= \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ -1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \right\}
 \end{aligned}$$

- Weyl group action

$$S_k = S_{(k, k+1)} \quad \text{for } k=1, \dots, n-1.$$




$$S_n = \begin{pmatrix} 1 & & & \\ & \ddots & & \\ & & 1 & \\ & & & -1 \end{pmatrix}$$

$$W(D_n) \cong (\mathbb{Z}/2\mathbb{Z})^{n-1} \rtimes S_n \subseteq (\mathbb{Z}/2\mathbb{Z})^n \rtimes S_n$$

2. D_4

- Weights nearest to the origin

D_4 is more symmetric.

#	typical coordinates	symbol	
$8 = 2 \cdot 4$	$(1, 0, 0, 0)^T$	v_i & $-v_i$	
$8 = 2^3$	$\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)^T$ odd sign	$v_{\pm\pm\pm\pm}$	
$8 = 2^3$	$\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)^T$ even sign	$v_{\pm\pm\pm\pm}$	
	$\frac{1}{2}(1, 1, 1, 1)^T$	v_+	

If not restricted to the standard representation case,

$$\langle v_i, v_j \rangle \in \{1, \frac{1}{2}, 0, -\frac{1}{2}, -1\}.$$