Eine Woche, ein Beispiel

12.21. Hodge structure

Ref:

[Car17]: James Carlson, Stefan Müller-Stach, and Chris Peters. Period Mappings and Period Domains. 2nd ed. Cambridge University Press, 2017. https://doi.org/10.1017/9781316995846.

X: cpt Kähler mfld of dim n.

Slogan: Hodge structure collects (nearly) all linear algebraic structures of X.

These linear algebraic structures are: $k = p+q$

① $H^k(X;\mathbb{Z})$

② $H^q(X;\Omega_X^p) \cong \{\text{harmonic } (p,q)\text{-forms}\}$

③ A comparison iso as $\mathbb{C}$-v.s. (J-action)

$$H^k(X;\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{C} \cong \bigoplus_{p+q=k} H^q(X;\Omega_X^p)$$

②′ A Hodge filtration

$$H^k(X;\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{C} \cong H^k(\Omega_X^\cdot) \supset \underset{H^{0,k}}{H^{k-1}(\Omega_X^{\geq 1})} \supset \underset{H^{1,k-1}}{H^{k-2}(\Omega_X^{\geq 2})} \supset \cdots \supset \underset{H^{k-1,1}}{H^0(\Omega_X^{\geq k})} \supset \underset{H^{k,0}}{0}$$

③′ Compatability with conj_J:

$$H^k(X,\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{C} = H^{k-p}(\Omega_X^{\geq p}) \oplus \overline{H^{p-1}(\Omega_X^{\geq k-p+1})}$$

Here, $\Omega_X^{\geq p} := \sigma_{\geq p}\Omega_X^\cdot$. We have

$$\begin{tikzcd}
 & & 0 \arrow[dr] & & 0 & & \\
 & & & \Omega_X^{\geq 1} \arrow[ur] \arrow[dr] & & & \\
0 \arrow[r] & \mathbb{C} \arrow[r] \arrow[dr] & \Omega_X^0 \arrow[ur] \arrow[r, "d^1"] & \Omega_X^1 \arrow[r, "d^2"] \arrow[dr] & \Omega_X^2 \arrow[ur] \arrow[r, "d^3"] & \cdots & \\
 & & \Omega_X^\cdot = \Omega_X^{\geq 0} \arrow[ur] & & \Omega_X^{\geq 2} \arrow[ur] \arrow[dr] & & \\
 & 0 \arrow[ur] & & 0 \arrow[ur] & & 0 &
\end{tikzcd}$$

④ (polarization)

A bilinear form

$$\Lambda : H^k(X;\mathbb{C}) \otimes H^k(X;\mathbb{C}) \xrightarrow{\Lambda} H^{2k}(X;\mathbb{C}) \xrightarrow{\wedge \omega^{n-k}} H^{2n}(X;\mathbb{C}) \cong \mathbb{C}$$

$$\Lambda(\xi,\eta) := \int_X \xi \wedge \eta \wedge \omega^{n-k}$$

with two compatability conditions:

Assume ω is integral, i.e., X is proj

$$\begin{array}{ccccccc}
\Lambda : H^k(X;\mathbb{Z}) & \otimes & H^k(X;\mathbb{Z}) & \xrightarrow{\Lambda} & H^{2k}(X;\mathbb{Z}) & \xrightarrow{\wedge \omega^{n-k}} & H^{2n}(X;\mathbb{Z}) \cong \mathbb{Z} \\
 & \cap & & & \cap & & \cap \\
\Lambda : H^k(X;\mathbb{C}) & \otimes & H^k(X;\mathbb{C}) & \xrightarrow{\Lambda} & H^{2k}(X;\mathbb{C}) & \xrightarrow{\wedge \omega^{n-k}} & H^{2n}(X;\mathbb{C}) \cong \mathbb{C} \\
 & \cup & & & \cup & & \cong \\
\Lambda : H^q(X;\Omega_X^p) & \otimes & H^p(X;\Omega_X^q) & \xrightarrow{\wedge} & H^k(X;\Omega_X^k) & \xrightarrow{\wedge \omega^{n-k}} & H^n(X,\omega_X) \cong \mathbb{C}
\end{array}$$

with two positivity conditions:

$$\Lambda(\xi,\bar{\xi}) \in \mathbb{R} \quad \text{for} \quad \xi \in H^k(X;\mathbb{R}) \qquad \xi = \bar{\xi} \text{ here}$$

$$i^{p-q+k(k-1)}\Lambda(\xi,\bar{\xi}) \geq 0 \quad \text{for} \quad \xi \in H^k_{prim}(X;\mathbb{C}) \cap H^{p,q}$$

And get "=" $\Leftrightarrow$ $\xi = 0$.

⚠ These positivities are not contradiction to each other.

If $\xi \in H^k(X;\mathbb{R}) \cap (H^k_{prim}(X;\mathbb{C}) \cap H^{p,q})$, then $\xi \in \overline{H^{p,q}} = H^{q,p}$,

then $p = q$, i.e.

$\xi \in H^{2p}_{prim}(X;\mathbb{R}) \cap H^{p,p}$, $\quad i^{p-q+k(k-1)} = (-1)^p$.

Rmk. 1. Even though $\int_X \omega^n > 0$, one don't nec have

$$\Lambda(\xi, \bar{\xi}) = \int_X \xi \wedge \bar{\xi} \wedge \omega^{n-k} \geq 0 \qquad \text{for any } \xi \in H^k(X; \mathbb{R}).$$

For example, for a cplx K3 surface X,
$H^2(X; \mathbb{Z})$ has signature $(3, 19)$, not $(0, 22)$ or $(2, 20)$

This is also a counterexample if one forgets "prim".

2. A de-Rham explanation for the following commutative diagram is here:
https://math.stackexchange.com/questions/29797/direct-proof-that-the-wedge-product-preserves-integral-cohomology-classes

$$\begin{tikzcd}
H^i(X; \mathbb{Z}) \otimes H^j(X; \mathbb{Z}) \arrow[r, "\cup"] \arrow[d] & H^{i+j}(X; \mathbb{Z}) \arrow[d] \\
H^i_{dR}(X; \mathbb{C}) \otimes H^j_{dR}(X; \mathbb{C}) \arrow[r, "\wedge"] & H^{i+j}_{dR}(X; \mathbb{C})
\end{tikzcd}$$