

$$\underline{\mathcal{F}l(n)}$$

Springer Fibers for $SL_n(\mathbb{C})$

Xiaoxiang Zhou

Universität Bonn

June 13, 2021

Recap: representation theory of finite groups

Restrict to **complex** representations, we have a nice theory:

- Any **representation** can be written ^{as a} ~~of~~ direct sum of **irreducible representation**;
- We can extract **information** of irreducible representations from the **character table**:

$$\#\{\text{irreducible representations}\} = \#\{\text{conjugation classes}\}$$

$$\sum_{\chi:\text{irr}} (\dim \chi)^2 = \#G$$

However, in general,

- NO standard way finding an **explicit construction** of all irreducible representations;
- NO **one-to-one correspondence** between irreducible representations and conjugation classes.

In this talk, we use two methods to understand representations of S_n , and find connections/analogs between them.

methods	objects
combinatorial	Young diagram, Young tableau
geometrical	Springer fiber of $SL_n(\mathbb{C})$, irreducible components

Goal of the Part I


- Explicitly **construct irreducible representations** of S_n by Young diagram;
- Compute the character table;
 - **$\dim \chi_i$** **by recursion** / Hook length formula
 - character by Frobenius formula
- Compute other representations.
 - e.g. \otimes , Sym^m , Λ^m ;
 - e.g. M_λ .

Notation

For boxes:



(Young) diagram



filling

11	78	11
6	8	

standard filling

3	5	4
1	2	

tableau


6	11	11
8	78	

standard tableau


1	3	4
2	5	

Order of Young diagram:


inclusion




dominance




Lexicographic ordering



tree of Young diagram



tree of Young diagram




Order

→ \subseteq inclusion

\trianglelefteq dominance

\preceq Lexicographic ordering



S_n & Young diagram

Proposition

We have the one-to-one correspondence

$$\left\{ \begin{array}{l} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\} \begin{array}{c} \xrightarrow{\text{partition of } n} \\ \xleftarrow{\lambda = \lambda_1^{v_1} \dots \lambda_k^{v_k}} \end{array} \left\{ \begin{array}{l} \text{Conjugation class} \\ \text{of } S_n \end{array} \right\}$$

S_n & Young diagram


Proposition

We have the one-to-one correspondence

$$\left\{ \begin{array}{l} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\} \begin{array}{c} \xleftarrow{\text{partition of } n} \\ \xrightarrow{\lambda = \lambda_1^{v_1} \dots \lambda_k^{v_k}} \end{array} \left\{ \begin{array}{l} \text{Conjugation class} \\ \text{of } S_n \end{array} \right\}$$

Example

$n = 10$ ~~10~~ ||



$$\begin{array}{c} \xleftarrow{11=5+4+1+1} \\ \xrightarrow{\lambda=5 \cdot 4 \cdot 1^2} \end{array} (12345)(6789)(10)(11)$$

S_n & Young diagram

Proposition

We have the one-to-one correspondence

$$\left\{ \begin{array}{l} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\} \begin{array}{c} \xleftarrow{\text{partition of } n} \\ \xrightarrow{\lambda = \lambda_1^{v_1} \dots \lambda_k^{v_k}} \end{array} \left\{ \begin{array}{l} \text{Conjugation class} \\ \text{of } S_n \end{array} \right\}$$

Claim

$$\left\{ \begin{array}{l} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\} \begin{array}{c} \xleftarrow{?} \\ \xrightarrow{?} \end{array} \left\{ \begin{array}{l} \text{Irreducible rep} \\ \text{of } S_n \end{array} \right\}$$

S_n & Young diagram

Claim

$$\left\{ \begin{array}{l} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\} \xleftrightarrow{\quad ? \quad} \left\{ \begin{array}{l} \text{Irreducible rep} \\ \text{of } S_n \end{array} \right\}$$

Remark

Reduced to: for each Young diagram λ ,
construct an irreducible representation S^λ , and
prove $S^\lambda = S^{\lambda'} \Rightarrow \lambda = \lambda'$.

The construction of $S^\lambda \subseteq M^\lambda$

Tabloid: equivalence class of standard filling

$$\begin{array}{|c|c|c|} \hline 3 & 5 & 4 \\ \hline 1 & 2 & \\ \hline \end{array} \begin{array}{c} \leftarrow \\ = \\ \rightarrow \end{array} \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 2 & 1 & \\ \hline \end{array} \begin{array}{c} \leftarrow \\ = \\ \rightarrow \end{array} \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 1 & 2 & \\ \hline \end{array} := \underline{\{345/12\}}$$

The construction of $S^\lambda \subseteq M^\lambda$

Tabloid: equivalence class of standard filling

$$\begin{array}{|c|c|c|} \hline 3 & 5 & 4 \\ \hline 1 & 2 & \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 2 & 1 & \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 1 & 2 & \\ \hline \end{array} := \{345/12\}$$

$$C(T) = \{Id, (13), (25), (13)(25)\}$$

$$\mathcal{T}^\lambda := \{\text{Young tabloid}\} = \{\text{standard filling } \{T\}\} / \sim$$

$$M^\lambda := \langle \{T\} \in \mathcal{T}^\lambda \rangle_{\mathbb{C}} \quad \text{ ~~$C(T) := \{\sigma \in S_n \mid \{\sigma \cdot T\} \sim \{T\}\}$~~ }$$

$$v_T := \sum_{q \in C(T)} \text{sgn}(q) \{q \cdot T\} \in M^\lambda$$

$$S^\lambda := \mathbb{C}[S_n] \cdot v_T \subseteq M^\lambda$$

invariant subspace of M^λ

The construction of $S^\lambda \subseteq M^\lambda$

$$\mathcal{T}^\lambda := \{\text{Young tabloid}\} = \{\text{standard filling } \{T\}\}$$

$$M^\lambda := \left\langle \{T\} \in \mathcal{T}^\lambda \right\rangle_{\mathbb{C}} \quad C(T) := \{\sigma \in S_n \mid \{\sigma \cdot T\} \sim \{T\}\}$$

$$v_T := \sum_{q \in C(T)} \text{sgn}(q) \{q \cdot T\} \in M^\lambda$$

$$S^\lambda := \mathbb{C}[S_n] \cdot v_T \subseteq M^\lambda \quad \text{invariant subspace of } M^\lambda$$

Example ($\lambda = 3 \cdot 2$)



$$\mathcal{T}^\lambda = \left\{ \begin{array}{l} \{123/45\}, \{124/35\}, \{125/34\}, \{134/25\}, \{135/24\}, \\ \{145/23\}, \{234/15\}, \{235/14\}, \{245/13\}, \{345/12\} \end{array} \right\}$$

$$M^\lambda = \left\langle \begin{array}{l} \{123/45\}, \{124/35\}, \{125/34\}, \{134/25\}, \{135/24\}, \\ \{145/23\}, \{234/15\}, \{235/14\}, \{245/13\}, \{345/12\} \end{array} \right\rangle_{\mathbb{C}}$$

The construction of $S^\lambda \subseteq M^\lambda$

$$\underline{v_T} := \sum_{q \in C(T)} \text{sgn}(q) \{q \cdot T\} \in M^\lambda$$

$$\boxed{\sigma v_T = v_{\sigma T}} \quad \forall \sigma \in S_n$$

$$S^\lambda := \mathbb{C}[S_n] \cdot \underline{v_T} \subseteq M^\lambda$$

invariant subspace of M^λ

Example ($\lambda = 3 \cdot 2$)

$$T = \begin{array}{|c|c|c|} \hline 3 & 5 & 4 \\ \hline 2 & 1 & \\ \hline \end{array}$$

$$C(T) = \{\text{Id}, (23), (15), (23)(15)\}$$

$$\underline{v_T} = \left\{ \begin{array}{|c|c|c|} \hline 3 & 5 & 4 \\ \hline 2 & 1 & \\ \hline \end{array} \right\} - \left\{ \begin{array}{|c|c|c|} \hline 2 & 5 & 4 \\ \hline 3 & 1 & \\ \hline \end{array} \right\} - \left\{ \begin{array}{|c|c|c|} \hline 3 & 1 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} \right\} + \left\{ \begin{array}{|c|c|c|} \hline 2 & 1 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \right\}$$

$$= \{345/12\} - \{245/13\} - \{134/25\} + \{124/35\} \in \underline{M^\lambda}$$

$$\underline{S^\lambda} = \langle \underline{v_T} \rangle_{\mathbb{C}[S_n]} = \langle \underline{v_{T'}} | T' : \text{standard tableau} \rangle_{\mathbb{C}}$$

Main theorem of S^λ

Proposition

Fix the Young diagram λ , the corresponding representation S^λ has the following properties:

- 1 the linear space S^λ has a basis $\{v_{T'} | T' : \text{standard tableau}\}$,
e.p. $\dim S^\lambda = \#\{\text{standard tableau}\}$;
- 2 the representation S^λ is irreducible;
- 3 for the Young diagram λ' , $S^{\lambda'} \cong S^\lambda \Rightarrow \lambda' = \lambda$.

Proof: basis

Proposition

- ① the linear space S^λ has a basis $\{v_{T'} | T' : \text{standard tableau}\}$,
e.p. $\dim S^\lambda = \#\{\text{standard tableau}\}$;

→ especially.

Proof

- S^λ is generated by $\{v_{T'} | T' : \text{standard filling}\}$,
eliminate the relations, we get $v_{T'} = \pm v_{T'_0}$. T'_0 : a standard tableau.

e.g. $\mathcal{V} \begin{array}{|c|c|c|} \hline 3 & 5 & 4 \\ \hline 2 & 1 & \\ \hline \end{array} \xrightarrow{\text{column}} \mathcal{V} \begin{array}{|c|c|c|} \hline 2 & 1 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \xrightarrow{\text{row}} - \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array}$

- $\{v_{T'} | T' : \text{standard tableau}\}$ are linear independent. ←

e.g. $x_1 \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} + x_2 \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} + x_3 \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} + x_4 \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & \\ \hline \end{array} - x_5 \mathcal{V} \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array} = 0 \quad x_i \in \mathbb{C}$

$\begin{array}{l} - \begin{array}{|c|c|c|} \hline 1 & 5 & 3 \\ \hline 4 & 2 & \\ \hline \end{array} \end{array}$

$\{123/45\} \rightarrow x_1 = 0 \quad \{134/25\} \rightarrow x_3 = 0$
 $\{124/35\} \rightarrow x_2 = 0 \quad \{125/34\} \rightarrow x_4 = 0$
 $\{135/24\} \rightarrow x_5 = 0$

linear ordering

We use a linear ordering of standard filling by

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} \longrightarrow 54321$$

∨

$$\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} \longrightarrow 52431$$

In the proof, we knock out the biggest one.

Example $(2, 2, 3)$

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & 6 \\ \hline \end{array} > \begin{array}{|c|c|} \hline 1 & \cancel{3} \\ \hline \cancel{3} & 4 \\ \hline 5 & 6 \\ \hline \end{array} > \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \cancel{3} & \cancel{5} \\ \hline \cancel{4} & 6 \\ \hline \end{array} > \begin{array}{|c|c|} \hline \cancel{1} & 2 \\ \hline 3 & 4 \\ \hline 5 & 6 \\ \hline \end{array} > \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & 6 \\ \hline \end{array}$$

654321 654231 645231 635241 635241

Proof: part 2&3

Proposition

- ② the representation S^λ is irreducible;
- ③ for the Young diagram λ' , $S^{\lambda'} \cong S^\lambda \Rightarrow \lambda' = \lambda$.

We have to introduce element b_T in $\mathbb{C}[S_n]$ by

$$b_T := \sum_{q \in C(T)} \text{sgn}(\sigma) \sigma$$

then

- $v_T = b_T \cdot \{T\}$;
- $\tau(b_T) = \text{sgn}(\tau)b_T$ for any $\tau \in C(T)$;
- $b_T \cdot b_T = \#C(T) \cdot b_T$;
- $b_T M^\lambda = b_T S^\lambda = \mathbb{C}v_T \neq 0$;
 $b_T M^{\lambda'} = b_T S^{\lambda'} = 0$ for $\lambda' > \lambda$.

Proof: part 2&3

Proposition

- ② *the representation S^λ is irreducible;*
- ③ *for the Young diagram λ' , $S^{\lambda'} \cong S^\lambda \Rightarrow \lambda' = \lambda$.*

$$\left[\begin{array}{l} b_T M^\lambda = b_T S^\lambda = \mathbb{C} v_T \neq 0 ; \\ b_T M^{\lambda'} = b_T S^{\lambda'} = 0 \end{array} \right. \quad \text{for } \lambda' > \lambda$$

✱To show S^λ is irreducible: only need to show indecomposability.
If $S^\lambda = V \oplus W$ as $\mathbb{C}[S_n]$ -module, then

$$\begin{aligned} \mathbb{C} v_T &= b_T S^\lambda = b_T V \oplus b_T W \\ \Rightarrow b_T V &= \mathbb{C} v_T & (\text{or } b_T W &= \mathbb{C} v_T) \\ \Rightarrow S^\lambda &= \mathbb{C}[S_n] \cdot v_T = \mathbb{C}[S_n] \cdot \mathbb{C} v_T = \mathbb{C}[S_n] \cdot b_T V \subseteq V \end{aligned}$$



Proof: part 2&3

Proposition

- ② *the representation S^λ is irreducible;*
- ③ *for the Young diagram λ' , $S^{\lambda'} \cong S^\lambda \Rightarrow \lambda' = \lambda$.*

$$\begin{aligned} b_T M^\lambda &= b_T S^\lambda = \mathbb{C}v_T \neq 0 ; \\ b_T M^{\lambda'} &= b_T S^{\lambda'} = 0 \quad \text{for } \lambda' > \lambda \end{aligned}$$


✱To show $S^{\lambda'} \cong S^\lambda \Rightarrow \lambda' = \lambda$:

If not w.l.o.g. suppose $\lambda' > \lambda$. Then


$$\underline{b_T S^{\lambda'}} = b_T S^\lambda \implies \underline{\mathbb{C}v_T} \cong 0,$$

contradiction!

Example



Example



Example: trivial representation

$$\lambda = \square\square\square = 3^1$$

$$\underline{M^\lambda} = \langle \{123\} \rangle = \underline{\mathbb{C}}$$

$$\underline{T} = \boxed{1 \mid 2 \mid 3}$$

$$\underline{C(T)} = \text{Id}$$

$$v_T = \left\{ \boxed{1 \mid 2 \mid 3} \right\}$$

$$\underline{S^\lambda} = \mathbb{C}[S_3] \cdot v_T = \mathbb{C}v_T$$

$\sigma \quad \quad \quad \tau$

Example: alternating representation

$$\lambda = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} = 1^3$$

$$M^\lambda = \langle \{1/2/3\}, \{1/3/2\}, \{2/1/3\}, \{2/3/1\}, \{3/1/2\}, \{3/2/1\} \rangle_{\mathbb{C}}$$

$$T = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$$

$$C(T) = S_3$$

$$v_T = \{1/2/3\} - \{1/3/2\} - \{2/1/3\} \\ + \{2/3/1\} + \{3/1/2\} - \{3/2/1\}$$

$$S^\lambda = \mathbb{C}[S_3] \cdot v_T = \mathbb{C}v_T$$

$$(23)v_T = \{1/3/2\} - \{1/2/3\} - \{3/1/2\} \\ + \{3/2/1\} + \{2/1/3\} - \{2/3/1\} = -v_T$$

Example: standard representation

$$\underline{\lambda} = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} = 2 \cdot 1$$

$$M^{\lambda} = \langle \{12/3\}, \{13/2\}, \{23/1\} \rangle_{\mathbb{C}}$$

$$\underline{T} = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}$$

$$\underline{C(T)} = \{\text{Id}, (13)\}$$

$$\underline{v_T} = \{12/3\} - \{23/1\} \quad 3 - 1$$

$$\underline{S^{\lambda}} = \mathbb{C}[S_3] \cdot \underline{v_T} \cong \mathbb{C}^3$$

$$\underline{(12)v_T} = \{12/3\} - \{13/2\} \quad 3 - 2$$

$$\underline{(13)v_T} = \{23/1\} - \{12/3\} = \underline{-v_T} \quad 1 - 3$$


Goal of the Part 1

- Explicitly construct irreducible representations of S_n by Young diagram;
- Compute the character table;
 - $\dim \chi_i$ by recursion / Hook length formula
 - character by Frobenius formula
- Compute other representations.
 - e.g. \otimes , Sym^m , Λ^m ;
 - e.g. M_λ .

Example: dimension of irreducible representation

$$\dim S^\lambda = \#\{\text{standard tableau of } \lambda\} = ?$$


Example



Example: dimension of irreducible representation


$$\dim S^\lambda = \#\{\text{standard tableau of } \lambda\} = ?$$

Example




$$\rightarrow \dim S^\lambda = \sum_{\substack{\lambda' \subseteq \lambda \\ |\lambda'| = n-1}} \dim S^{\lambda'}$$

Example: dimension of irreducible representation



Example: dimension of irreducible representation




1
1 1
2 1
3 3 1
4 6 4 1

Hook length formula

It helps us compute the dimension of S^λ without induction.

Step 1: count the length of hook.



$$\text{Step 2: } \dim S^\lambda = \frac{n!}{\prod(\text{hook lengths})}$$

Special case: (m, l)


Ballot problem

In an election where candidate **A** receives m votes and candidate **B** receives l votes with $m \geq l$, what is the probability that **A** will be (non-strictly) ahead of **B** throughout the count?

Proposition

Each process of the count corresponds to each standard tableau of form (m, l) .

Example



Special case: (m, m)



Corollary

$$\underline{\dim S^{(m,m)}} = C_m = \frac{1}{m+1} \binom{2m}{m}.$$

where C_m is the n -th Catalan number.

Special case: (m, m)

Corollary

$$\dim S^{(m,m)} = C_m = \frac{1}{m+1} \binom{2m}{m}.$$

where C_m is the n -th Catalan number.

Catalan number has many interpretations. For example, it counts the number of crossingless matchings of $2n$ points.

Ex. $m=3$

crossingless matchings
of 6 points



$\{(1,2), (3,4), (5,6)\}$



$\{(1,4), \dots\}$



Special case: (m, m)

Corollary

$$\dim S^{(m,m)} = C_m = \frac{1}{m+1} \binom{2m}{m}.$$

where C_m is the n -th Catalan number.

Catalan number has many interpretations. For example, it counts the number of crossingless matchings of $2n$ points.

Ex. $m=3$

crossingless matchings
of 6 points



1	3	5
2	4	6



1	3	4
2	5	6



1	2	5
3	4	6



1	2	4
3	5	6



1	2	3
4	5	6

52

Goal of the Part II

- Definition of Springer fiber;
- Some examples of Springer fiber;
- Properties: (closely connected with combinatorics)
 - irreducible component?
 - dimension? $X = \bigsqcup \mathbb{C}^k$
 - affine paving? – CW complex?
 - cohomology? – ring structure?
 - smooth?
 - explicit description?
- Weyl group action on top homology.

Goal of the Part II

- **Definition** of Springer fiber;
- Some examples of Springer fiber;
- Properties: (closely connected with combinatorics)
 - irreducible component?
 - dimension?
 - affine paving? – CW complex?
 - cohomology? – ring structure?
 - smooth?
 - explicit description?
- Weyl group action on top cohomology.

Definition

$$\begin{array}{ccc}
 \widehat{\mathfrak{g}} \subseteq \mathfrak{g} \times \mathcal{B} & \longrightarrow & \mathcal{B}_n = \mathcal{F}(n) \\
 \downarrow \mu & \rightsquigarrow & \downarrow \mu|_{\mathcal{N} \times \mathcal{B}} \\
 \mathfrak{g}_n = \mathfrak{sl}_n(\mathbb{C}) & & \mathcal{N} \quad \mathcal{X} \\
 & & \text{resolution of nilpotent cone}
 \end{array}$$

$\mu^{-1}(X)$
 \downarrow
 X

Let $X \in \mathfrak{g}$ be a nilpotent element. The Springer fiber B_X over X is defined as

$$\mathfrak{B}_X := \mu^{-1}(X)$$

$$\stackrel{\text{def}}{=} \{B \in \mathfrak{B} \mid X \in B\}$$

$$\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C}) \stackrel{\text{def}}{=} \{0 \subseteq V_1 \subseteq V_2 \subseteq \cdots \subseteq V_n = \mathbb{C}^n \mid \underbrace{XV_i \subseteq V_{i-1}}_{\uparrow} \mid \dim V_i = i\}$$

By the Jordan normal form, we have

$$\left\{ \begin{array}{c} \text{Nilpotent element} \\ \text{in } \mathfrak{gl}_n(\mathbb{C}) \end{array} \right\} / \text{conj} \longleftrightarrow \left\{ \begin{array}{c} \text{Young diagram} \\ \text{of } n \text{ boxes} \end{array} \right\}$$

$$X_\lambda = \text{diag}(\underbrace{J_{\lambda_1}, \dots, J_{\lambda_1}}_{v_1}, J_{\lambda_2}, \dots, J_{\lambda_k}) \longleftrightarrow \lambda = \lambda_1^{v_1} \dots \lambda_k^{v_k}$$

$$J_{\lambda_i} = \begin{bmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ & & & 0 \end{bmatrix}_{\lambda_i \times \lambda_i}$$

Denote $B_\lambda := B_{X_\lambda}$. $B_X \cong B_{gXg^{-1}}$ for any $g \in G$

→ Theorem (we will not give the proof.)

As S_n -representation, $S^\lambda \cong H_{\text{top}}^{\text{top}}(B_\lambda)$.


Corollary

$$\#\{\text{irreducible component of } B_\lambda\} = \dim S^\lambda$$

Goal of the Part II

- Definition of Springer fiber;
- Some **examples** of Springer fiber;
- **Properties:** (closely connected with combinatorics)
 - irreducible component?
 - dimension?
 - affine paving? – CW complex?
 - cohomology? – ring structure?
 - smooth?
 - explicit description?
- Weyl group action on top cohomology.

tree of Young diagram



Example: $\lambda = 3$



$$\{e_1, \dots, e_n\} \subseteq \mathbb{C}^3$$

$$X_\lambda = \begin{bmatrix} 0 & 1 & \\ & 0 & 1 \\ & & 0 \end{bmatrix}$$

$$B_\lambda = \left\{ 0 \subseteq \langle \overset{e_1}{?} \rangle \subseteq \langle \overset{e_1}{?}, \overset{e_2}{?} \rangle \subseteq \mathbb{C}^3 \right\} \curvearrowright X_\lambda = \{*\}$$

In general, $B_\lambda = \{*\}$ when λ has only one row.

Example: $\lambda = (1, 1, 1)$



$$X_\lambda = \begin{bmatrix} 0 & & \\ & 0 & \\ & & 0 \end{bmatrix}$$

$$\underline{B_\lambda} = \left\{ 0 \subseteq \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3 \right\} \curvearrowright X_\lambda = \underline{\mathcal{F}\ell(3)}$$

In general, $\underline{B_\lambda} = \mathcal{F}\ell(n)$ when $\lambda = 1^n$.

Example: $\lambda = (2, 1)$



$$\underline{X_\lambda} = \begin{bmatrix} 0 & 1 & \\ & 0 & \\ & & 0 \end{bmatrix}$$

$$\underline{B_\lambda} = \left\{ 0 \subseteq \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3 \right\} \curvearrowright X_\lambda = \underline{\mathbb{P}^1 \vee \mathbb{P}^1}$$

$$B_\lambda = \begin{aligned} & \{0 \subseteq \langle ae_1 + e_3 \rangle \subseteq \langle e_1, e_3 \rangle \subseteq \mathbb{C}^3\} \xrightarrow{\triangle \mathbb{C}} \text{blue circle} \\ & \cup \\ & \{0 \subseteq \langle e_1 \rangle \subseteq \langle e_1, be_2 + ce_3 \rangle \subseteq \mathbb{C}^3\} \xrightarrow{\quad} \text{red circle} \end{aligned}$$


← $\{0 \subseteq \langle e_1 \rangle \subseteq \langle e_1, e_3 \rangle \subseteq \mathbb{C}^3\}$

In general, $B_\lambda = \underbrace{\mathbb{P}^1 \vee \dots \vee \mathbb{P}^1}_{n-1}$ when $\lambda = (n-1, 1)$.



Properties of $B_\lambda = \underbrace{\mathbb{P}^1 \vee \dots \vee \mathbb{P}^1}_{n-1}$

- irreducible component: $n-1$
- $\dim B_\lambda = 1$
- affine paving: Yes
- cohomology group:
- smooth: \checkmark
- explicit description: \checkmark
- Weyl group action on $H_{\text{top}}^{\text{top}}(B_\lambda) \cong \mathbb{C}^{n-1}$:



$$H^i(B_\lambda) = \begin{cases} \mathbb{C} & i=0 \\ 0 & i=1 \\ \mathbb{C}^{n-1} & i=2 \end{cases}$$

$$H_i(B_\lambda) = \begin{cases} \mathbb{C} & i=0 \\ 0 & i=1 \\ \mathbb{C}^{n-1} & i=2 \end{cases}$$

Tool: stratification/cellular fibration/affine paving

$$\begin{array}{ccc}
 \begin{array}{l} \phi \rightarrow \\ B_\lambda \rightarrow \\ \vdots \\ B_{\lambda''} \rightarrow \end{array} & B_\lambda & \{0 \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq \mathbb{C}^n\}^{\hookrightarrow X_\lambda} \\
 & \downarrow \pi & \downarrow \\
 & \mathbb{P}^{n-1} & [V_1]
 \end{array}$$

Remark

In general, We don't have a natural CW complex structure.

We don't understand the ring structure.

Return!

For $\lambda = 1^3$, $B_\lambda \cong \mathcal{F}\ell(3)$ can be viewed as $\mathcal{F}\ell(2)$ -bundle over \mathbb{P}^2 .


$$\begin{array}{ccc} \mathcal{F}\ell(2) & \longrightarrow & \mathcal{F}\ell(3) \\ & & \downarrow \pi \\ & & \mathbb{P}^2 \quad [v] \end{array}$$

$$\pi^{-1}([v]) = \{0 \subseteq \langle v \rangle \subseteq \langle v, ? \rangle \subseteq \mathbb{C}^3\} \cong \mathcal{F}\ell(2)$$

Return!

For $\lambda = (2, 1)$, $B_\lambda \cong \mathbb{P}^1 \vee \mathbb{P}^1$:


$$\begin{array}{ccc} \mathbb{P}^1 = B_{\lambda_1} & \longrightarrow & B_{\lambda_1} \\ \{e_i\} = B_{\lambda_2} & \longrightarrow & B_{\lambda_2} \\ & & \downarrow \pi \\ & & \mathbb{P}^1 \end{array}$$



$$\begin{aligned} \pi^{-1}([e_1]) &= \{0 \subseteq \langle e_1 \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3\} \curvearrowright \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \\ &\cong \{0 \subseteq \langle ? \rangle \subseteq \mathbb{C}^2\} \curvearrowright \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = B_{\underline{0}, \underline{1}, \underline{1}} \\ \pi^{-1}([e_3]) &= \{0 \subseteq \langle e_3 \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3\} \curvearrowright \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \\ &\cong \{0 \subseteq \langle ? \rangle \subseteq \mathbb{C}^2\} \curvearrowright \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = B_{\underline{1}, \underline{1}, \underline{2}} \end{aligned}$$

Return!

For $\lambda = (2, 1)$, $B_\lambda \cong \mathbb{P}^1 \vee \mathbb{P}^1$:




$$\pi^{-1}([e_1]) \cong B_{1,1} \quad \pi^{-1}([e_3]) \cong B_2$$

$$\begin{aligned} \pi^{-1}([ae_1 + e_3]) &= \left\{ 0 \subseteq \langle ae_1 + e_3 \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3 \right\} \curvearrowright \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ &\stackrel{(f_1, f_2, f_3) = (e_1, e_2, ae_1 + e_3)}{\cong} \left\{ 0 \subseteq \langle f_3 \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3 \right\} \curvearrowright \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \cong \pi^{-1}([e_3]) \end{aligned}$$

By this way, $\pi^{-1}(\mathbb{P}^1 \setminus \{[e_1]\}) \cong \underline{B_2} \times \underline{\mathbb{C}}$ induces an affine paving.

Example: $\lambda = (2, 1, 1)$



$$X_\lambda = \begin{bmatrix} 0 & 1 & & \\ & 0 & & \\ & & 0 & \\ & & & 0 \end{bmatrix}$$

$$B_\lambda = \{0 \subseteq \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \langle ?, ?, ? \rangle \subseteq \mathbb{C}^4\} \curvearrowright X_\lambda$$

$$\mathcal{F}(1) = B_{1,1,1} \longrightarrow$$

$$\mathbb{P}^1 \vee \mathbb{P}^1 = B_{2,1} \longrightarrow B_{2,1,1}$$

$$\downarrow \uparrow$$

$$\mathbb{P}^1$$

$$\mathbb{P}^1$$

$$\mathbb{P}^2$$

$$\mathbb{P}^3$$

$$0 \subseteq \mathbb{P}^1 \subseteq \mathbb{P}^1 \subseteq \mathbb{P}^2 \subseteq \mathbb{P}^3 \subseteq \dots$$

3-dim



3-dim

1

+ 3-dim

2

= 3 components


Example: $\lambda = (2, 2)$



$$X_\lambda = \begin{bmatrix} 0 & 1 & & \\ & 0 & & \\ & & 0 & 1 \\ & & & 0 \end{bmatrix} \quad \ker X_\lambda = \langle e_1, e_3 \rangle$$

$$B_\lambda = \{0 \subseteq \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \langle ?, ?, ? \rangle \subseteq \mathbb{C}^4\} \curvearrowright X_\lambda$$

$$\mathbb{P}^1 \vee \mathbb{P}^1 = B_{2,2} \longrightarrow B_{2,2} \\ \downarrow \\ \mathbb{P}^1$$




Using the same technique, we can get


- B_λ has an affine paving \rightsquigarrow cohomology;
- Each irreducible component in B_λ has same dimension;
- It's easy to compute the dimension and the number of irreducible component.

Game: compute!


0	0	0	0
1	1	1	
2	2		



Answer: dimension



Answer: the number of irreducible component



Smooth problem

B_λ

Results


- Not all the the irreducible components of B_λ are smooth;
For example, one component of $B_{2,2,1,1}$ is not smooth.
- • All the components of B_λ are nonsingular iff

$$\lambda \in \{(\lambda_1, 1, 1, \dots), (\lambda_1, \lambda_2), (\lambda_1, \lambda_2, 1), (2, 2, 2)\}$$



.

tree of Young diagram



(n, n) case

We have an explicit description in the 2-row case when we forget the variety structure. Use this description, we can get the cohomology group structure.

Definition and Theorem

Let α be a crossingless matching, define 

$$\tilde{B}_{\alpha; m, m} := \left\{ (x_1, \dots, x_{2m}) \in (\mathbb{P}^1)^{2m} \mid x_i = x_j \text{ if } (i, j) \in \alpha \right\} \subseteq (\mathbb{P}^1)^{2m}$$

$$\tilde{B}_{m, m} := \bigcup_{\alpha} \tilde{B}_{\alpha; m, m} \subseteq (\mathbb{P}^1)^{2m}$$

then we have a homeomorphism

$$B_{m, m} \cong \tilde{B}_{m, m}$$

(n, n) case

Definition and Theorem

Let α be a crossingless matching, define

$$\tilde{B}_{\alpha; m, m} := \left\{ (x_1, \dots, x_{2m}) \in (\mathbb{P}^1)^{2m} \mid x_i = x_j \text{ if } (i, j) \in \alpha \right\} \subseteq (\mathbb{P}^1)^{2m}$$

$$\tilde{B}_{m, m} := \bigcup_{\alpha} \tilde{B}_{\alpha; m, m} \subseteq (\mathbb{P}^1)^{2m}$$

then we have a homeomorphism

$$B_{m, m} \cong \tilde{B}_{m, m}$$

Example ($m=2$)

$$\alpha = \{(1, 2), (3, 4)\} \quad \tilde{B}_{\alpha; 2, 2} = \left\{ (x_1, x_1, x_2, x_2) \in (\mathbb{P}^1)^4 \right\} \cong (\mathbb{P}^1)^2$$

$$\beta = \{(1, 4), (2, 3)\} \quad \tilde{B}_{\beta; 2, 2} = \left\{ (x_1, x_2, x_2, x_1) \in (\mathbb{P}^1)^4 \right\} \cong (\mathbb{P}^1)^2$$

$$B_{2, 2} \cong \tilde{B}_{2, 2} \cong (\mathbb{P}^1)^2 \vee (\mathbb{P}^1)^2$$

\mathbb{P}^1 -bundle over \mathbb{P}^1 \mathbb{P}^1 \mathbb{P}^1



THANKS

Thank you for listening!

Thank Rui Xiong for providing the package of Young diagram,

Thank my roommate David Cueto for pointing out typos,

Thank Prof. Eberhart for offering valuable materials and advice!

55