

Bruhat–Tits building

Xiaoxiang Zhou

Humboldt-Universität zu Berlin

January 26, 2026

Figures of Bruhat–Tits building




Figure: $\mathcal{B}_{\mathrm{SL}_3(\mathbb{Q}_p)}$, from a talk by Annette Werner

Figures of Bruhat–Tits building




Figure: $\mathcal{B}_{SL_3(\mathbb{Q}_p)}$, from buildings.gallery

Figures of Bruhat–Tits building




Figure: $\mathcal{B}_{\mathrm{SL}_2(\mathbb{Q}_2)}$

Figures of Bruhat–Tits building




Figure: $\mathcal{B}_{\mathrm{SL}_2(\mathbb{Q}_2)}$

Plan of the talk

- 1 Spherical buildings
- 2 p -adic buildings
- 3 The Gromov-Schoen theorem

Plan of the talk

- 1 Spherical buildings
- 2 p -adic buildings
- 3 The Gromov-Schoen theorem

Standard subgroups

G ^{connected}
reductive

$$\underline{GL_n(\mathbb{C}), SL_n}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/B = \{ \text{flags of } \kappa^n \}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

Standard subgroups

$$\begin{aligned} B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} & \rightsquigarrow \mathrm{GL}_n(\kappa)/B = \{ \text{flags of } \kappa^n \} \\ & = \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \\ & \\ & \mathrm{Gr}(r, n) \\ & = \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$\begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & | & * \\ \hline & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & | & * \\ \hline & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & | & * \\ \hline & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & & * \\ \vdots & & \vdots \\ * & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

T is a torus, so every rep decomposes as direct sum of 1-dim reps.

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & & * \\ \vdots & \ddots & \vdots \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

T is a torus, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & & * \\ \vdots & & \vdots \\ * & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

T is a torus, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n \quad \text{characters (1-dim reps)}$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & & * \\ \vdots & \ddots & \vdots \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

T is a torus, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n \quad \text{characters (1-dim reps)}$$

$$X_*(T) := \mathrm{Hom}(\mathbb{G}_m, T) \cong \mathbb{Z}^n$$

Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & & * \\ \vdots & \ddots & \vdots \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

T is a torus, so every rep decomposes as direct sum of 1-dim reps.

$$\begin{aligned} X^*(T) &:= \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n && \text{characters (1-dim reps)} \\ X_*(T) &:= \mathrm{Hom}(\mathbb{G}_m, T) \cong \mathbb{Z}^n && \text{cocharacters (1-parameter subgps)} \end{aligned}$$

Weyl group

Weyl group

Definition (Weyl group)

$$W := N_G(T)/T.$$

Weyl group

Definition (Weyl group)

$$W := N_G(T)/T.$$

Example

When $G = \mathrm{GL}_n(\kappa)$,

$$N_G(T) = \{ \text{monoidal matrices} \}$$

$$N_G(T)/T \cong S_n$$

Weyl group of type A

$$\begin{pmatrix} * & & & \\ & * & & \\ & & * & \\ & & & * \end{pmatrix}$$

Weyl group

WGT

G G G

$$W = N_G(T)/T \xrightarrow{\sim} W \otimes X^*(T)$$

$W \otimes X_*(T)$

Definition (Weyl group)

$$W := N_G(T)/T.$$

Example

When $G = \mathrm{GL}_n(\kappa)$,

$$N_G(T) = \{ \text{monoidal matrices} \}$$

$$N_G(T)/T \cong S_n \quad \text{Weyl group of type A}$$

Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B. \quad = \text{"BWB"}$$

So the Weyl group is the "heart" of the reductive group.

Weyl group

Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B.$$

So the Weyl group is the “heart” of the reductive group.

Weyl group

Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B.$$

So the Weyl group is the “heart” of the reductive group.




Figure: Pinned butterfly

Weyl group action on cocharacter lattices

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \underline{\mathbb{Z}\varepsilon_1} \oplus \underline{\mathbb{Z}\varepsilon_2}$, where

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where


$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where




$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$


$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$


$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where

$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \end{aligned}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$


When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where

$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{GL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\varepsilon_1 : \mathbb{G}_m \longrightarrow T \\ x \longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix}$$

$$\varepsilon_2 : \mathbb{G}_m \longrightarrow T \\ x \longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\varepsilon : \mathbb{G}_m \longrightarrow T \\ x \longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where

Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$

(12) (23)

Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where

$$\left(\begin{array}{c} \xrightarrow{\quad s_1 \quad} \\ \xrightarrow{\quad \mathrm{Id} \quad} \end{array} \right) \quad \varepsilon : \mathbb{G}_m \longrightarrow T$$

$$x \longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where

$$\varepsilon_1 : \mathbb{G}_m \longrightarrow T$$

$$x \longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix}$$

$$\varepsilon_2 : \mathbb{G}_m \longrightarrow T$$

$$x \longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$


Weyl group action on cocharacter lattices

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$

Non-standard subgroups

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

$$gTg^{-1}$$

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

Fact

*All non-standard subgroups are conjugated to standard subgroups.
Therefore,*

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

Fact

*All non-standard subgroups are conjugated to standard subgroups.
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \mid g \in G \}$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \mid g \in G \}$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \mid g \in G \}$$

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

Fact

*All non-standard subgroups are conjugated to standard subgroups.
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \mid g \in G \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \mid g \in G \}$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \mid g \in G \}$$

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

Fact

*All non-standard subgroups are conjugated to standard subgroups.
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \mid g \in G \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \mid g \in G \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \mid g \in G \}$$

Non-standard subgroups

The subgroup $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$ is not the only maximal torus.

Fact

*All non-standard subgroups are conjugated to standard subgroups.
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \mid g \in G \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \mid g \in G \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \mid g \in G \} \cong G/N_G(T)$$

G/T

Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \left\{ gBg^{-1} \mid g \in G \right\} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \left\{ gPg^{-1} \mid g \in G \right\} \cong G/P$$

$$\{ \text{maximal tori} \} = \left\{ gTg^{-1} \mid g \in G \right\} \cong G/N_G(T)$$

Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \left\{ gB_{\mathfrak{o}}g^{-1} \mid g \in G \right\} \cong G/B_{\mathfrak{o}}$$

$$\{ \text{parabolic subgroups} \} = \left\{ gP_{\mathfrak{o}}g^{-1} \mid g \in G \right\} \cong G/P_{\mathfrak{o}}$$

$$\{ \text{maximal tori} \} = \left\{ gT_{\mathfrak{o}}g^{-1} \mid g \in G \right\} \cong G/N_G(T_{\mathfrak{o}})$$

$$\{ \underset{\uparrow}{(B, T)} \mid B \supset T \} = \left\{ (g \underset{\uparrow}{B_0} g^{-1}, g \underset{\uparrow}{T_0} g^{-1}) \right\} \cong G/T_0$$


Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gB_0g^{-1} \mid g \in G \} \cong G/B_0$$

$$\{ \text{parabolic subgroups} \} = \{ gP_0g^{-1} \mid g \in G \} \cong G/P_0$$

$$\{ \text{maximal tori} \} = \{ gT_0g^{-1} \mid g \in G \} \cong G/N_G(T_0)$$

$$\{(B, T) \mid B \supset T\} = \{(gB_0g^{-1}, gT_0g^{-1})\} \cong G/T_0$$




Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \left\{ gB_0g^{-1} \mid g \in G \right\} \cong G/B_0$$

$$\{ \text{parabolic subgroups} \} = \left\{ gP_0g^{-1} \mid g \in G \right\} \cong G/P_0$$

$$\{ \text{maximal tori} \} = \left\{ gT_0g^{-1} \mid g \in G \right\} \cong G/N_G(T_0)$$

$$\{(B, T) \mid B \supset T\} = \left\{ (gB_0g^{-1}, gT_0g^{-1}) \right\} \cong G/T_0$$




Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gB_0g^{-1} \mid g \in G \} \cong G/B_0$$

$$\{ \text{parabolic subgroups} \} = \{ gP_0g^{-1} \mid g \in G \} \cong G/P_0$$


$$\{ \text{maximal tori} \} = \{ gT_0g^{-1} \mid g \in G \} \cong G/N_G(T_0)$$

$$\{ (B, T) \mid B \supset T \} = \{ (gB_0g^{-1}, gT_0g^{-1}) \} \cong G/T_0$$



Weyl group action on cocharacter lattices(revisited)

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

Weyl group action on cocharacter lattices(revisited)

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$


Weyl group action on cocharacter lattices(revisited)

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where




$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$


Weyl group action on cocharacter lattices(revisited)

When $G = \mathrm{SL}_2(\kappa)$, $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon$, where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When $G = \mathrm{SL}_3(\kappa)$, $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$, $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$, where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$

Definition (chamber, apartment and building)

Given a maximal torus T , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{B \supset T} \mathcal{C}_B,$$

Definition (chamber, apartment and building)

Given a maximal torus T , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{B \supset T} \mathcal{C}_B,$$

and the building is

$$\mathcal{B} := \left(\bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_B \mathcal{C}_B.$$

e.g. $k = \mathbb{F}_p$
 $x = \mathbb{C}$

Example of spherical building

Example of spherical building

When $G = \mathrm{SL}_2(\mathbb{F}_2)$, the building \mathcal{B} has 3 apartments and 3 chambers.




Figure: $\mathcal{B}_{\mathrm{SL}_2(\mathbb{F}_2)}$

$\mathcal{B}_{\mathrm{SL}_2(\mathbb{F}_3)}$



$(\mathcal{B} - \{o\}) / \mathbb{R}_{>0}$



Example of spherical building

When $G = \mathrm{SL}_2(\mathbb{F}_2)$, the building \mathcal{B} has 3 apartments and 3 chambers.




Figure: $\mathcal{B}_{\mathrm{SL}_2(\mathbb{F}_2)}$

When $G = \mathrm{SL}_3(\mathbb{F}_2)$, the building \mathcal{B} has 28 apartments and 21 chambers.

Remark

\mathcal{B} inherits the metric structure from $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$.

Remark

\mathcal{B} inherits the metric structure from $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$.

\mathcal{B} has also polysimplicial complex structure.

When $\kappa = \mathbb{F}_p$, \mathcal{B} is finite. i.e., having finite many chambers

Remark

\mathcal{B} inherits the metric structure from $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$.

\mathcal{B} has also polysimplicial complex structure.

When $\kappa = \mathbb{F}_p$, \mathcal{B} is finite. i.e., having finite many chambers

Proposition

- Any two chambers lie in one apartment.

$$B_1, B_2, \quad \exists T \subset B_1 \cap B_2$$

$$B_0 \cap gB_0g^{-1} = B_0 \cap b_1\omega B_0\omega^{-1}b_1^{-1} \supset b_1Tb_1^{-1}$$

$$g = b_1\omega b_2$$

Remark

\mathcal{B} inherits the metric structure from $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$.


\mathcal{B} has also polysimplicial complex structure.

When $\kappa = \mathbb{F}_p$, \mathcal{B} is finite. i.e., having finite many chambers

Proposition

- Any two chambers lie in one apartment.
- There is a unique geodesic through any two points $p_1, p_2 \in \mathcal{B}$.

$$\begin{array}{l} p_1 \in \mathcal{C}_{B_1} \\ p_2 \in \mathcal{C}_{B_2} \end{array} \quad \begin{array}{l} \subset \\ \subset \end{array} \mathcal{A}_T$$



Plan of the talk

- 1 Spherical buildings
- 2 p -adic buildings
- 3 The Gromov-Schoen theorem

p -adic notation

symbol	name	example
F	NA local field	\mathbb{Q}_p
$\mathcal{O} = \mathcal{O}_F$	ring of integers	\mathbb{Z}_p
$\mathfrak{p} = \mathfrak{p}_F$	maximal ideal	$p\mathbb{Z}_p$
$\kappa = \mathcal{O}/\mathfrak{p}$	residue field	\mathbb{F}_p
$\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$	uniformizer	p
$v : F^* \longrightarrow \mathbb{Z}$	valuation	$v\left(\frac{a}{b}p^k\right) = k$

p -adic notation

symbol	name	example
F	NA local field	\mathbb{Q}_p
$\mathcal{O} = \mathcal{O}_F$	ring of integers	\mathbb{Z}_p
$\mathfrak{p} = \mathfrak{p}_F$	maximal ideal	$p\mathbb{Z}_p$
$\kappa = \mathcal{O}/\mathfrak{p}$	residue field	\mathbb{F}_p
$\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$	uniformizer	p
$v : F^* \longrightarrow \mathbb{Z}$	valuation	$v\left(\frac{a}{b}p^k\right) = k$

standard subgroups in the p -adic world

$$\underline{GL_n(\mathcal{O})} \subset GL_n(F)$$

maximal compact subgp

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$\cup$$

$$\mathcal{B}, \mathcal{P}$$

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix}$$

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \cdots & \cdots \\ \mathfrak{p} & \mathcal{O} \end{pmatrix}$$

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \begin{array}{l} \text{Parahoric subgroup} \\ \text{=Parabolic Iwahori subgroup} \end{array}$$

standard subgroups in the p -adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \begin{array}{l} \text{Parahoric subgroup} \\ \text{=Parabolic Iwahori subgroup} \end{array}$$

Remark

They also have moduli interpretations. For example,

$$\begin{aligned} \mathrm{GL}_n(F)/I &\cong \{ \mathfrak{p}L = L_0 \subset L_1 \subset \cdots \subset L_n = L \mid L_{i+1}/L_i \cong \kappa \} \\ &= \{ \mathcal{O}\text{-lattice chains in } F^n \} \end{aligned}$$

Extended Weyl group

Extended Weyl group

To get the Iwahori decomposition

$$G(F) = \bigsqcup_{\varpi \in W_{\text{ext}}} I\varpi I, \quad = "IW_{\text{ext}}I"$$

we define the extended Weyl group as

$$W_{\text{ext}} := N_G(T(\mathcal{O}))/T(\mathcal{O}) \cong X_*(T) \rtimes W_f. \quad \leftarrow \text{finite}$$

Extended Weyl group

To get the Iwahori decomposition

$$G(F) = \bigsqcup_{\varpi \in W_{\text{ext}}} I\varpi I,$$

we define the extended Weyl group as

$$W_{\text{ext}} := N_G(T(\mathcal{O}))/T(\mathcal{O}) \cong X_*(T) \rtimes W_f.$$

Example

When $G = \text{GL}_n(F)$,

$$W_{\text{ext}} = \left\{ \text{monoidal matrices} \right\} / \begin{pmatrix} \begin{smallmatrix} * & & \\ \mathbb{Z} & & \\ & * & \end{smallmatrix} \\ \begin{smallmatrix} * \\ \mathcal{O}^* \cdots \mathcal{O}^* \end{smallmatrix} \end{pmatrix} \cong \mathbb{Z}^n \rtimes S_n.$$

Extended Weyl group action

Extended Weyl group action

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \mapsto \mu + u\lambda u^{-1}$$

Extended Weyl group action

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

Extended Weyl group action

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$


Extended Weyl group action

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




Extended Weyl group action

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$



Extended Weyl group action

When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where

Extended Weyl group action


When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where

$$s_1 = \left(\begin{array}{c|c} 1 & \\ \hline -1 & 1 \end{array} \right)_{(1\ 2)} \quad s_2 = \left(\begin{array}{c|c} 1 & \\ \hline & 1 \end{array} \right)_{(2\ 3)} \quad s_0 = \left(\begin{array}{cc} & \pi^{-1} \\ 1 & \\ -\pi & \end{array} \right)_{(3\ 1)}$$

Extended Weyl group action

When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where


$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & \\ & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ & 1 & \\ -\pi & & \end{pmatrix}$$



Extended Weyl group action

When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where


$$s_1 = \begin{pmatrix} & 1 & \\ -1 & & \\ & & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ -\pi & 1 & \\ & & \end{pmatrix}$$



Extended Weyl group action

When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 & \\ -1 & & \\ & & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ -\pi & 1 & \\ & & \end{pmatrix}$$



Extended Weyl group action




Figure: Reduced expressions labels, from Lievis

Non-standard subgroups in the p -adic world

Non-standard subgroups in the p -adic world

Similarly,

$$\{ \text{Iwahori subgroups} \} = \left\{ gI_0g^{-1} \mid g \in G \right\} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \left\{ g\tilde{P}_0g^{-1} \mid g \in G \right\} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \left\{ gT_0g^{-1} \mid g \in G \right\} \cong G/N_G(T_0(\mathcal{O}))$$

Non-standard subgroups in the p -adic world

Similarly,

$$\{ \text{Iwahori subgroups} \} = \left\{ gI_0g^{-1} \mid g \in G \right\} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \left\{ g\tilde{P}_0g^{-1} \mid g \in G \right\} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \left\{ gT_0g^{-1} \mid g \in G \right\} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \left\{ (gI_0g^{-1}, gT_0g^{-1}) \right\} \cong G/T_0(\mathcal{O})$$

Non-standard subgroups in the p -adic world


Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \mid g \in G \} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \mid g \in G \} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \mid g \in G \} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \{(gI_0g^{-1}, gT_0g^{-1})\} \cong G/T_0(\mathcal{O})$$



Non-standard subgroups in the p -adic world


Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \mid g \in G \} \cong G/I_0$$



$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \mid g \in G \} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \mid g \in G \} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \{(gI_0g^{-1}, gT_0g^{-1})\} \cong G/T_0(\mathcal{O})$$



Comparison




Extended Weyl group action(revisited)

$\downarrow T_0$ $T \otimes_0 F \cong T' \otimes_0 F$
 W_{ext} acts on $X_*(T)$ by "twisted conjugation":

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




Extended Weyl group action(revisited)

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




Extended Weyl group action(revisited)

W_{ext} acts on $X_*(T)$ by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When $G = \text{SL}_2(F)$, $W_{\text{ext}} = \langle s_0, s_1 \rangle$, where


$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




Extended Weyl group action(revisited)

When $G = \mathrm{SL}_3(F)$, $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$, where


$$s_1 = \begin{pmatrix} & 1 & \\ -1 & & \\ & & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ -\pi & 1 & \\ & & \end{pmatrix}$$




Extended Weyl group action(revisited)



Extended Weyl group action(revisited)



Extended Weyl group action(revisited)



p -adic building

p -adic building

Definition (chamber, apartment and building)

Given a maximal torus T over \mathcal{O} , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{I \supset T} \mathcal{C}_I,$$

and the p -adic building is

$$\mathcal{B} := \left(\bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_I \mathcal{C}_I.$$

p -adic building

Definition (chamber, apartment and building)

Given a maximal torus T over \mathcal{O} , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{I \supset T} \mathcal{C}_I,$$

and the p -adic building is

$$\mathcal{B} := \left(\bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_I \mathcal{C}_I.$$



Remark

Similarly, any two chambers lie in one apartment, and there is a unique geodesic through $p_1, p_2 \in \mathcal{B}$.

Plan of the talk

- 1 Spherical buildings
- 2 p -adic buildings
- 3 The Gromov-Schoen theorem

The Gromov-Schoen theorem

The Gromov-Schoen theorem

Theorem

Let F be a NA local field, (M, g) be a cpt conn Riemannian manifold with the universal covering space \widetilde{M} .

The Gromov-Schoen theorem

Theorem

Let F be a NA local field, (M, g) be a cpt conn Riemannian manifold with the universal covering space \widetilde{M} .

For any reductive homomorphism

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

The Gromov-Schoen theorem

Theorem

Let F be a NA local field, (M, g) be a cpt conn Riemannian manifold with the universal covering space \widetilde{M} .

For any reductive homomorphism

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

We call ρ reductive when $\overline{\rho(\pi_1(M))}^{\mathrm{Zar}} \subseteq \mathrm{GL}_n(F)$ is reductive.

The Gromov-Schoen theorem

Theorem

Let F be a NA local field, (M, g) be a cpt conn Riemannian manifold with the universal covering space \widetilde{M} .

For any reductive homomorphism

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

there exists a ^① $\pi_1(M)$ -equivariant ^② Lipschitz continuous ^④ regular ^③ harmonic map

$$h_\rho : \mathcal{U} \rightarrow \mathbb{R}^n$$

$$h_\rho : \widetilde{M} \longrightarrow \mathcal{B}_{\mathrm{GL}_n(F)}$$

$$\exists C \\ d(h_\rho(x), h_\rho(y)) \leq C d(x, y)$$

We call ρ reductive when $\overline{\rho(\pi_1(M))}^{\mathrm{Zar}} \subseteq \mathrm{GL}_n(F)$ is reductive.

$$\nabla h_\rho = 0$$

regularity

regularity

Definition

h_ρ is regular at $x \in \widetilde{M}$ if



a neighbourhood of x has image inside an apartment \mathcal{A}_T of \mathcal{B} .

regularity

Definition

h_ρ is regular at $x \in \widetilde{M}$ if

a neighbourhood of x has image inside an apartment \mathcal{A}_T of \mathcal{B} .

h_ρ is regular if

$$\mathrm{codim}_{\widetilde{M}} \left\{ x \in \widetilde{M} \mid h_\rho \text{ is not regular at } x \right\} \geq 2.$$

regularity

Definition

h_ρ is regular at $x \in \widetilde{M}$ if

a neighbourhood of x has image inside an apartment \mathcal{A}_T of \mathcal{B} .

h_ρ is regular if

$$\operatorname{codim}_{\widetilde{M}, \mathbb{R}} \left\{ x \in \widetilde{M} \mid h_\rho \text{ is not regular at } x \right\} \geq 2.$$

Example

The map

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

is regular.

regularity

Example

The map

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

is regular.


regularity

Example

The map

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

is regular.



$$\mathbb{R}^2 \longrightarrow \{y^2 = x^2\}$$


regularity

Example

The map

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

is regular.



$$\mathbb{R}^2 \longrightarrow \{y^2 = x^2\}$$


regularity

Example


The map

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

is regular.



f



\mathbb{R}^2

\longrightarrow

$\{y^2 = x^2\}$

$\mathcal{B}_{SL_2(\mathbb{F}_3)}$

Thanks for listening!

You can get this slide at:

[https://github.com/ramified/personal_tex_collection/raw/main/
Bruhat-Tits_building/Bruhat-Tits_building.pdf](https://github.com/ramified/personal_tex_collection/raw/main/Bruhat-Tits_building/Bruhat-Tits_building.pdf)