

# Bruhat–Tits building

Xiaoxiang Zhou

Humboldt-Universität zu Berlin

January 24, 2026

# Figures of Bruhat–Tits building




Figure:  $\mathcal{B}_{SL_3(\mathbb{Q}_p)}$ , from Annette Werner's talk

# Figures of Bruhat–Tits building




Figure:  $\mathcal{B}_{\mathrm{SL}_3(\mathbb{Q}_p)}$ , from buildings.gallery

# Figures of Bruhat–Tits building




Figure:  $\mathcal{B}_{SL_2(\mathbb{Q}_2)}$

# Figures of Bruhat–Tits building




Figure:  $\mathcal{B}_{\mathrm{SL}_2(\mathbb{Q}_2)}$

# Process

- 1 Spherical building
- 2 p-adic building
- 3 Gromov-Schoen theorem

# Process

- 1 Spherical building
- 2 p-adic building
- 3 Gromov-Schoen theorem

# Standard subgroups



# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/B = \{ \text{flags of } \kappa^n \}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

# Standard subgroups

$$\begin{aligned}
 B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} & \rightsquigarrow \mathrm{GL}_n(\kappa)/B = \{ \text{flags of } \kappa^n \} \\
 & = \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \\
 & \\
 & \mathrm{Gr}(r, n) \\
 & = \{ V \subset \kappa^n \mid \dim V = r \}
 \end{aligned}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$\begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$



# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

$T$  is comm, so every rep decomposes as direct sum of 1-dim reps.

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

$T$  is comm, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & * \\ \vdots & \vdots \\ * & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

$T$  is comm, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n \quad \text{characters (1-dim reps)}$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & \vdots & * \\ \vdots & \ddots & \vdots \\ \vdots & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

$T$  is comm, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n \quad \text{characters (1-dim reps)}$$

$$X_*(T) := \mathrm{Hom}(\mathbb{G}_m, T) \cong \mathbb{Z}^n$$

# Standard subgroups

$$B = \begin{pmatrix} * & * & * \\ & * & * \\ & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/B &= \{ \text{flags of } \kappa^n \} \\ &= \{ V_1 \subset \cdots \subset V_n = \kappa^n \mid \dim V_i = i \} \end{aligned}$$

$$P = \begin{pmatrix} * & \vdots & * \\ \vdots & \ddots & \vdots \\ \vdots & & * \end{pmatrix} \rightsquigarrow \begin{aligned} \mathrm{GL}_n(\kappa)/P &= \mathrm{Gr}(r, n) \\ &= \{ V \subset \kappa^n \mid \dim V = r \} \end{aligned}$$

$$T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix} \rightsquigarrow \mathrm{GL}_n(\kappa)/T = \{ \kappa^n = W_1 \oplus \cdots \oplus W_n \mid \dim W_i = 1 \}$$

$T$  is comm, so every rep decomposes as direct sum of 1-dim reps.

$$X^*(T) := \mathrm{Hom}(T, \mathbb{G}_m) \cong \mathbb{Z}^n \quad \text{characters (1-dim reps)}$$

$$X_*(T) := \mathrm{Hom}(\mathbb{G}_m, T) \cong \mathbb{Z}^n \quad \text{cocharacters (1-parameter subgps)}$$

# Weyl group

# Weyl group

Definition (Weyl group)

$$W := N_G(T)/T.$$

# Weyl group

## Definition (Weyl group)

$$W := N_G(T)/T.$$

## Example

When  $G = \mathrm{GL}_n(\kappa)$ ,

$$N_G(T) = \{ \text{monoidal matrixes} \}$$

$$N_G(T)/T \cong S_n \quad \text{Weyl group of type } A$$



# Weyl group

## Definition (Weyl group)

$$W := N_G(T)/T.$$

## Example

When  $G = \mathrm{GL}_n(\kappa)$ ,

$$N_G(T) = \{ \text{monoidal matrixes} \}$$

$$N_G(T)/T \cong S_n \quad \text{Weyl group of type A}$$

## Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B.$$

So the Weyl group is the “heart” of the reductive group.

# Weyl group

## Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B.$$

So the Weyl group is the “heart” of the reductive group.

# Weyl group

## Remark

We have Bruhat decomposition proved by Gauss elimination

$$G = \bigsqcup_{\omega \in W} B\omega B.$$

So the Weyl group is the “heart” of the reductive group.




Figure: Pinned butterfly

# Weyl group action on cocharacter lattices

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where


$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$


$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$



# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$


$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$


$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where

$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \end{aligned}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$


When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where

$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{GL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & b \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & \\ & x \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_2 = \{\mathrm{Id}, s_1\}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & b^{-2} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where

# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & b^{-2} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & b^{-2} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where

$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$




# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & b^{-2} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \\ \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \\ W = S_3 &= \langle s_1, s_2 \rangle \end{aligned}$$


# Weyl group action on cocharacter lattices

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$

# Non-standard subgroups

# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

## Fact

*All non-standard subgroups are conjugated to standard subgroups.  
Therefore,*

# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

## Fact

*All non-standard subgroups are conjugated to standard subgroups.  
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \}$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \}$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \}$$

# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

## Fact

*All non-standard subgroups are conjugated to standard subgroups.  
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \}$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \}$$

# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

## Fact

*All non-standard subgroups are conjugated to standard subgroups.  
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \}$$



# Non-standard subgroups

The subgroup  $T = \begin{pmatrix} * & & \\ & \ddots & \\ & & * \end{pmatrix}$  is not the only maximal torus.

## Fact

*All non-standard subgroups are conjugated to standard subgroups.  
Therefore,*

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \} \cong G/N_G(T)$$

# Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \} \cong G/N_G(T)$$

# Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \} \cong G/N_G(T)$$

$$\{(B, T) \mid B \supset T\} = \{(gB_0g^{-1}, gT_0g^{-1})\} \cong G/T_0$$


# Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \} \cong G/N_G(T)$$

$$\{(B, T) \mid B \supset T\} = \{(gB_0g^{-1}, gT_0g^{-1})\} \cong G/T_0$$




# Non-standard subgroups

$$\{ \text{Borel subgroups} \} = \{ gBg^{-1} \} \cong G/B$$

$$\{ \text{parabolic subgroups} \} = \{ gPg^{-1} \} \cong G/P$$

$$\{ \text{maximal tori} \} = \{ gTg^{-1} \} \cong G/N_G(T)$$

$$\{(B, T) \mid B \supset T\} = \{(gB_0g^{-1}, gT_0g^{-1})\} \cong G/T_0$$




# Weyl group action on cocharacter lattices(revisit)

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$


# Weyl group action on cocharacter lattices(revisit)

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$


# Weyl group action on cocharacter lattices(revisit)

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$




# Weyl group action on cocharacter lattices(revisit)

When  $G = \mathrm{SL}_2(\kappa)$ ,  $T = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon$ , where



$$\begin{aligned} \varepsilon : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \\ W = S_2 &= \{\mathrm{Id}, s_1\} \end{aligned}$$

When  $G = \mathrm{SL}_3(\kappa)$ ,  $T = \begin{pmatrix} a & & \\ & b & \\ & & a^{-1}b^{-1} \end{pmatrix}$ ,  $X_*(T) = \mathbb{Z}\varepsilon_1 \oplus \mathbb{Z}\varepsilon_2$ , where



$$\begin{aligned} \varepsilon_1 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} x & & \\ & x^{-1} & \\ & & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \varepsilon_2 : \mathbb{G}_m &\longrightarrow T \\ x &\longmapsto \begin{pmatrix} 1 & & \\ & x & \\ & & x^{-1} \end{pmatrix} \end{aligned}$$

$$W = S_3 = \langle s_1, s_2 \rangle$$

## Definition (chamber, apartment and building)

Given a maximal torus  $T$ , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{B \supset T} \mathcal{C}_B,$$

## Definition (chamber, apartment and building)

Given a maximal torus  $T$ , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{B \supset T} \mathcal{C}_B,$$

and the building is

$$\mathcal{B} := \left( \bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_B \mathcal{C}_B.$$

# Example of spherical building

## Example of spherical building

When  $G = \mathrm{SL}_2(\mathbb{F}_2)$ , the building  $\mathcal{B}$  has 3 apartments and 3 chambers.




Figure:  $\mathcal{B}_{\mathrm{SL}_2(\mathbb{F}_2)}$

## Example of spherical building

When  $G = \mathrm{SL}_2(\mathbb{F}_2)$ , the building  $\mathcal{B}$  has 3 apartments and 3 chambers.




Figure:  $\mathcal{B}_{\mathrm{SL}_2(\mathbb{F}_2)}$

When  $G = \mathrm{SL}_3(\mathbb{F}_2)$ , the building  $\mathcal{B}$  has 28 apartments and 21 chambers.

## Remark

$\mathcal{B}$  inherits the metric structure from  $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$ .

## Remark

$\mathcal{B}$  inherits the metric structure from  $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$ .

$\mathcal{B}$  has also polysimplicial complex structure.

When  $\kappa = \mathbb{F}_p$ ,  $\mathcal{B}$  is finite.



## Remark

$\mathcal{B}$  inherits the metric structure from  $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$ .

$\mathcal{B}$  has also polysimplicial complex structure.

When  $\kappa = \mathbb{F}_p$ ,  $\mathcal{B}$  is finite.

## Proposition

- *Two chambers lie in one apartment.*

## Remark

$\mathcal{B}$  inherits the metric structure from  $\mathcal{A}_T = X_*(T)_{\mathbb{R}}$ .

$\mathcal{B}$  has also polysimplicial complex structure.

When  $\kappa = \mathbb{F}_p$ ,  $\mathcal{B}$  is finite.

## Proposition

- *Two chambers lie in one apartment.*
- *There is a unique geodesic passing any two points  $p_1, p_2 \in \mathcal{B}$ .*

# Process

- 1 Spherical building
- 2 p-adic building
- 3 Gromov-Schoen theorem

## p-adic notation

symbol	name	example
$F$	NA local field	
$\mathcal{O} = \mathcal{O}_F$	integral ring	
$\mathfrak{p} = \mathfrak{p}_F$	maximal ideal	
$\kappa = \mathcal{O}/\mathfrak{p}$	residue field	
$\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$	uniformizer	
$v : F^* \longrightarrow \mathbb{Z}$	valuation	

## p-adic notation

symbol	name	example
$F$	NA local field	$\mathbb{Q}_p$
$\mathcal{O} = \mathcal{O}_F$	integral ring	$\mathbb{Z}_p$
$\mathfrak{p} = \mathfrak{p}_F$	maximal ideal	$p\mathbb{Z}_p$
$\kappa = \mathcal{O}/\mathfrak{p}$	residue field	$\mathbb{F}_p$
$\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$	uniformizer	$p$
$v : F^* \longrightarrow \mathbb{Z}$	valuation	$v\left(\frac{a}{b}p^k\right) = k$

# standard subgroups in p-adic world

# standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

## standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix}$$



## standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

# standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \hline \mathfrak{p} & \mathcal{O} \end{pmatrix}$$

# standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \hline \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Parahoric subgroup}$$

# standard subgroups in p-adic world

$$\pi : \mathrm{GL}_n(\mathcal{O}) \longrightarrow \mathrm{GL}_n(\kappa)$$

$$I = \pi^{-1}(B) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p} & \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Iwahori subgroup}$$

$$\tilde{P} = \pi^{-1}(P) = \begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \hline \mathfrak{p} & \mathcal{O} \end{pmatrix} \quad \text{Parahoric subgroup}$$

## Remark

They also have moduli interpretations. For example,

$$\begin{aligned} \mathrm{GL}_n(F)/I &\cong \{L = L_0 \subset L_1 \subset \cdots \subset L_n = \mathfrak{p}L \mid L_{i+1}/L_i \cong \kappa\} \\ &= \{\mathcal{O}\text{-lattice chains in } F^n\} \end{aligned}$$

# Extended Weyl group

# Extended Weyl group

To get the Iwahori decompositionn

$$G(F) = \bigsqcup_{\varpi \in W_{\text{ext}}} I\varpi I,$$

we define the extended Weyl group as

$$W_{\text{ext}} := N_G(T(\mathcal{O}))/T(\mathcal{O}) \cong X_*(T) \rtimes W_f.$$

# Extended Weyl group

To get the Iwahori decomposition

$$G(F) = \bigsqcup_{\varpi \in W_{\text{ext}}} I\varpi I,$$

we define the extended Weyl group as

$$W_{\text{ext}} := N_G(T(\mathcal{O}))/T(\mathcal{O}) \cong X_*(T) \rtimes W_f.$$

## Example

When  $G = \text{GL}_n(F)$ ,

$$W_{\text{ext}} = \{ \text{monoidal matrixes} \} / \begin{pmatrix} \mathcal{O}^* & & \\ & \ddots & \\ & & \mathcal{O}^* \end{pmatrix} \cong \mathbb{Z}^n \rtimes S_n.$$

# Extended Weyl group action



# Extended Weyl group action

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

# Extended Weyl group action

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

# Extended Weyl group action

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$


# Extended Weyl group action

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




# Extended Weyl group action

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$



# Extended Weyl group action

When  $G = \mathrm{SL}_3(F)$ ,  $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$ , where

# Extended Weyl group action


When  $G = \mathrm{SL}_3(F)$ ,  $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 & \\ -1 & & \\ & & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ & 1 & \\ -\pi & & \end{pmatrix}$$

# Extended Weyl group action

When  $G = \mathrm{SL}_3(F)$ ,  $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 & \\ -1 & & \\ & & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ & 1 & \\ -\pi & & \end{pmatrix}$$








# Extended Weyl group action

When  $G = \mathrm{SL}_3(F)$ ,  $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \\ & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ & 1 & \\ -\pi & & \end{pmatrix}$$



# Extended Weyl group action




Figure: Reduced expressions labels, from Lievis

# Non-standard subgroups in p-adic world

# Non-standard subgroups in p-adic world

Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \} \cong G/N_G(T_0(\mathcal{O}))$$

# Non-standard subgroups in p-adic world

Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \{(gI_0g^{-1}, gT_0g^{-1})\} \cong G/T_0(\mathcal{O})$$

# Non-standard subgroups in p-adic world


Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \} \cong G/\tilde{P}_0$$

$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \{(gI_0g^{-1}, gT_0g^{-1})\} \cong G/T_0(\mathcal{O})$$



# Non-standard subgroups in p-adic world


Similarly,

$$\{ \text{Iwahori subgroups} \} = \{ gI_0g^{-1} \} \cong G/I_0$$

$$\{ \text{paraholic subgroups} \} = \{ g\tilde{P}_0g^{-1} \} \cong G/\tilde{P}_0$$



$$\{ \text{maximal tori over } \mathcal{O} \} = \{ gT_0g^{-1} \} \cong G/N_G(T_0(\mathcal{O}))$$

$$\{(I, T) \mid I \supset T\} = \{(gI_0g^{-1}, gT_0g^{-1})\} \cong G/T_0(\mathcal{O})$$





# Comparison




# Extended Weyl group action(revisit)

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




# Extended Weyl group action(revisit)

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where

$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




# Extended Weyl group action(revisit)

$W_{\text{ext}}$  acts on  $X_*(T)$  by “twisted conjugation”:

$$W_{\text{ext}} \times X_*(T) \longrightarrow X_*(T) \quad (\mu \rtimes u, \lambda) \longmapsto \mu + u\lambda u^{-1}$$

When  $G = \text{SL}_2(F)$ ,  $W_{\text{ext}} = \langle s_0, s_1 \rangle$ , where


$$s_1 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \quad s_0 = \begin{pmatrix} & \pi^{-1} \\ -\pi & \end{pmatrix}$$




# Extended Weyl group action(revisit)

When  $G = \mathrm{SL}_3(F)$ ,  $W_{\mathrm{ext}} = \langle s_0, s_1, s_2 \rangle$ , where


$$s_1 = \begin{pmatrix} & 1 \\ -1 & \\ & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & & \\ & 1 & \\ & -1 & 1 \end{pmatrix} \quad s_0 = \begin{pmatrix} & & \pi^{-1} \\ -\pi & 1 & \\ & & \end{pmatrix}$$




# Extended Weyl group action(revisit)



## Extended Weyl group action(revisit)



## Extended Weyl group action(revisit)





# p-adic building

# p-adic building

## Definition (chamber, apartment and building)

Given a maximal torus  $T$  over  $\mathcal{O}$ , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{I \supset T} \mathcal{C}_I,$$

and the p-adic building is

$$\mathcal{B} := \left( \bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_I \mathcal{C}_I.$$

# p-adic building

## Definition (chamber, apartment and building)

Given a maximal torus  $T$  over  $\mathcal{O}$ , the apartment is

$$\mathcal{A}_T := X_*(T)_{\mathbb{R}} = \bigcup_{I \supset T} \mathcal{C}_I,$$

and the p-adic building is

$$\mathcal{B} := \left( \bigsqcup_T \mathcal{A}_T \right) / \sim = \bigcup_I \mathcal{C}_I.$$

## Remark

Similarly, two chambers lie in one apartment,  
and there is a unique geodesic passing  $p_1, p_2 \in \mathcal{B}$ .

# Process

- 1 Spherical building
- 2 p-adic building
- 3 Gromov-Schoen theorem

# Gromov-Schoen theorem

# Gromov-Schoen theorem

## Theorem

*Let  $F$  be a NA local field,  $(M, g)$  be a cpt conn Riemannian manifold with the universal covering space  $\widetilde{M}$ .*

# Gromov-Schoen theorem

## Theorem

*Let  $F$  be a NA local field,  $(M, g)$  be a cpt conn Riemannian manifold with the universal covering space  $\widetilde{M}$ .*

*For any reductive map*

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

# Gromov-Schoen theorem

## Theorem

*Let  $F$  be a NA local field,  $(M, g)$  be a cpt conn Riemannian manifold with the universal covering space  $\widetilde{M}$ .*

*For any reductive map*

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

*there exists a  $\pi_1(M)$ -equivariant Lipschitz continuous regular harmonic map*

$$h_\rho : \widetilde{M} \longrightarrow \mathcal{B}_{\mathrm{GL}_n(F)}$$



# Gromov-Schoen theorem

## Theorem

*Let  $F$  be a NA local field,  $(M, g)$  be a cpt conn Riemannian manifold with the universal covering space  $\widetilde{M}$ .*

*For any reductive map*

$$\rho : \pi_1(M) \longrightarrow \mathrm{GL}_n(F),$$

*there exists a  $\pi_1(M)$ -equivariant Lipschitz continuous regular harmonic map*

$$h_\rho : \widetilde{M} \longrightarrow \mathcal{B}_{\mathrm{GL}_n(F)}$$

We call  $\rho$  reductive when  $\overline{\rho(\pi_1(M))}^{\mathrm{Zar}} \subseteq \mathrm{GL}_n(F)$  is reductive.

# regularity

## regularity

## Definition

$h_\rho$  is regular at  $x \in \widetilde{M}$  if

a neighbourhood of  $x$  has image inside an apartment  $\mathcal{A}_T$  of  $\mathcal{B}$ .

## regularity

## Definition

$h_\rho$  is regular at  $x \in \widetilde{M}$  if

a neighbourhood of  $x$  has image inside an apartment  $\mathcal{A}_T$  of  $\mathcal{B}$ .

$h_\rho$  is regular if

$$\operatorname{codim}_{\widetilde{M}} \left\{ x \in \widetilde{M} \mid h_\rho \text{ is not regular at } x \right\} \geq 2.$$

# regularity

## Definition

$h_\rho$  is regular at  $x \in \widetilde{M}$  if

a neighbourhood of  $x$  has image inside an apartment  $\mathcal{A}_T$  of  $\mathcal{B}$ .

$h_\rho$  is regular if

$$\operatorname{codim}_{\widetilde{M}} \left\{ x \in \widetilde{M} \mid h_\rho \text{ is not regular at } x \right\} \geq 2.$$

## Example

*The map*

$$f : \mathbb{R}^2 \longrightarrow \{y^2 = x^2\} \quad (a, b) \longmapsto (a|b|, b|a|)$$

*is regular.*

# Thanks for listening!

You can get this slide at:

[https://github.com/ramified/personal\\_tex\\_collection/raw/main/Bruhat-Tits\\_building/Bruhat-Tits\\_building.pdf](https://github.com/ramified/personal_tex_collection/raw/main/Bruhat-Tits_building/Bruhat-Tits_building.pdf)