

# SUBVARIETIES IN COMPLEX ABELIAN VARIETIES

XIAOXIANG ZHOU

## CONTENTS

1. Introduction	1
Acknowledgement	3
2. Tangent Gauss map and conormal Gauss map	3
3. Monodromy group	8
4. Families of subvarieties	15
5. Dimension and homology class	18
References	26

## 1. INTRODUCTION

Let  $A$  be a complex abelian variety of dimension  $n$ . To any closed subvariety  $Z \subset A$  of dimension  $r$ , one can associate a reductive group  $G_Z$  through the convolution of perverse sheaves. This correspondence allows to reformulate problems in algebraic geometry in representation-theoretic terms, thereby offering new geometric insights. A crucial role in this context is played by the characteristic cycle, a fundamental invariant attached to a perverse sheaf, [8, §5–§8]. Recent work demonstrates that one can approach these characteristic cycles directly as conic Lagrangian cycles, without invoking the full machinery of perverse sheaves.

In [11], Krämer establishes a correspondence between Weyl group orbits and conic Lagrangian cycles in the cotangent bundle which arise as the conormal varieties of certain subvarieties. Despite this correspondence, an explicit geometric description of these cycles themselves remains elusive. In this article, we propose a purely geometric construction of these cycles and analyze three key properties of the associated subvarieties: irreducibility, dimension, and homology class.

To present our results, we begin by recalling several fundamental notions, which will be discussed in detail in Section 2. On a complex abelian variety  $A$ , the *translations* and *reflections* are special automorphisms of  $A$  of the form  $x \mapsto x + p$  and  $x \mapsto -x + p$ , respectively. For  $\sigma \in \text{Aut}(A)$  and a closed subvariety  $Z \subset A$ , we say that  $Z$  is *invariant* under  $\sigma$  if  $\sigma(Z) = Z$ . Given a subvariety  $Z \subset A$ , the *tangent Gauss map*  $\phi_Z$  assigns to  $p \in Z^{\text{sm}}$  the embedded tangent space to  $Z$  at  $p$  (cf. Definition 2.1), while the *conormal Gauss map*  $\gamma_Z : \mathbb{P}\Lambda_Z \subset A \times \mathbb{P}T_0A \rightarrow \mathbb{P}T_0A$  (cf. Definition 2.3) arises from the projectivized conormal variety of  $Z$  and is typically generically finite. The *monodromy group*  $\text{Gal}(\gamma_Z)$  encodes the permutation of the sheets of the covering  $\gamma_Z$  around its branch locus, while  $W_Z$  denotes the *Weyl group* of the reductive group  $G_Z$ .

Concerning irreducibility, it follows from the constructions in [11, 2.c] that the irreducible components of the appearing cycles correspond precisely to the orbits of a certain monodromy group. The discrepancy between the monodromy group and a closely related Weyl group provides a measure of the failure of irreducibility of these subvarieties. A natural question is how big this discrepancy could be. For this, we formulate a conjecture that refines [10, Conjecture 8]:

**Conjecture 1.1.** *Let  $A$  be an abelian variety of dimension  $n$ , and let  $C \subset A$  be a non-degenerate curve that is not invariant under any non-trivial translation. Then, the monodromy group  $\text{Gal}(\gamma_C)$  is big (see Definition 3.1).*

For  $n > 2$ , this conjecture admits a purely geometric reformulation:

**Conjecture 1.2.** *Let  $A$  be an abelian variety of dimension  $n > 2$ , and let  $C \subset A$  be a non-degenerate curve that is not invariant under any non-trivial translation in  $A$ . Then:*

- *If  $C$  is invariant under some reflection, the tangent Gauss map  $\phi_C : C \rightarrow \mathbb{P}^{n-1}$  is a double cover onto its image;*
- *If  $C$  is not invariant under any reflection, the tangent Gauss map  $\phi_C : C \rightarrow \mathbb{P}^{n-1}$  is generically injective.*

In general Conjecture 1.1 is still open. We analyze the Prym case thoroughly, which will lead to the following result:

**Theorem A** *For any double cover  $h : C \rightarrow C'$  of smooth, projective, non-hyperelliptic curves, one may view  $C$  as a subvariety of the Prym variety  $\text{Prym}(C/C')$  via the Abel–Prym map. Then the associated monodromy group is big, except in the case where  $C'$  is bielliptic and  $h$  arises as the pullback of an étale double cover of elliptic curves. In this exceptional situation, the curve  $C \subset \text{Prym}(C/C')$  is stable under a non-trivial translation.*

Theorem A follows from a combination of Proposition 3.9 and Corollary 3.11.

For the dimension and homology class of a subvariety  $Z$ , both invariants can be recovered from the homology class of  $\mathbb{P}\Lambda_Z \subset \mathbb{P}T^*A$  — that is, from the total Chern–Mather class of  $Z$ . Standard computational techniques are available when the monodromy group  $\text{Gal}(\gamma_Z)$  is a full or signed symmetric group. The notions of the Chern–Mather class and the Pontryagin product will be recalled in Definition 5.1 and Definition 5.6, respectively.

**Theorem B** *Consider a non-degenerate subvariety  $Z \subset A$  and a tuple  $(m) = (m_1, \dots, m_d) \in \mathbb{Z}^d$ . Let  $Z^{(m)} \subset A$  be as in Definition 4.7. Let  $d = \deg \gamma_Z$ ,  $\tilde{d} = d/2$ , and let  $c_i := c_{M,i}(\Lambda_Z)$  be the Chern–Mather class of  $Z$ . Let  $*$  be the Pontryagin product. We write  $\lambda \vdash l$  to indicate that  $\lambda = [\lambda_1, \dots, \lambda_{k'}]$  is a partition of  $l$ .*

- (1) When  $\text{Gal}(\gamma_Z) = S_d$ , the Chern–Mather classes of  $Z^{(m)}$  can be written as

$$c_{M,l}(\Lambda_{Z^{(m)}}) = \frac{1}{c_Z^{(m)}} \sum_{\lambda \vdash l} \mu_d^\lambda \left( \bigstar_{i=1}^{k'} c_{\lambda_i} \right)$$

for certain explicit coefficients  $c_Z^{(m)} \in \mathbb{N}_{>0}$  and  $\mu_d^\lambda \in \mathbb{Z}$ .

- (2) When  $Z = -Z$  and  $\text{Gal}(\gamma_Z) = W(C_{d/2})$ , the Chern–Mather classes of  $Z^{(m)}$  can be written as

$$c_{M,l}(\Lambda_{Z^{(m)}}) = \frac{1}{c_Z^{(m)}} \sum_{\lambda \vdash l} \tilde{\mu}_{\tilde{d}}^\lambda \left( \bigstar_{i=1}^{k'} c_{\lambda_i} \right)$$

for certain explicit coefficients  $c_Z^{(m)} \in \mathbb{N}_{>0}$  and  $\tilde{\mu}_{\tilde{d}}^\lambda \in \mathbb{Z}$ .

The proof of theorem B will occupy Section 5. In fact, we will show (1) in (5.1), and (2) in (5.2). The combinatorial coefficients  $\mu_d^\lambda$  and  $\tilde{\mu}_{\tilde{d}}^\lambda$  appearing in the formulas are defined in Definition 5.12, while the case of Jacobian varieties is discussed in detail in Example 5.14.

The structure of the article is as follows.

In Section 2, we fix notation and describe the Gauss map from three different perspectives. We also define the monodromy group and highlight its connection to the degree of the tangent Gauss map.

Section 3 is devoted to the explicit computation of the monodromy group in selected cases, with a particular focus on the Prym curve situation. Subsection 3.3 presents an example in which

the monodromy group is small; however, this does not constitute a counterexample to Conjecture 1.1, as the curve remains invariant under a non-trivial translation, as discussed in Lemma 3.14. In Subsection 3.1, we collect several sufficient conditions for the monodromy group to be large, which we then employ in Subsection 3.2 to show that, for all remaining Prym curve cases with  $g(C') > 9$ , the monodromy group is necessarily large.

In Section 4, we define the subvarieties  $Z^{(m)}$ , which serve as the central objects of this study. In Subsection 4.2, we briefly discuss how these varieties naturally arise as the characteristic cycles of certain perverse sheaves.

In Section 5, we focus on the computation of the dimension and the homology class of  $Z^{(m)}$ . After recalling the definition of the Chern-Mather class in Subsection 5.1, the task reduces to computing  $[\mathbb{P}\Lambda_{Z^{(m)}}]$ . Subsections 5.3 and 5.4 treat the type  $A$  case, where  $\text{Gal}(\gamma_Z) = S_d$ , and the type  $C$  case, where  $\text{Gal}(\gamma_Z) = W(C_{d/2})$ , respectively. The derivation of the formulas involves certain coefficients that are purely combinatorial in nature. These combinatorial contributions are treated separately in Subsection 5.5.

#### ACKNOWLEDGEMENT

I would like to express my sincere gratitude to my supervisor, Thomas Krämer, without whose support this work would not have been possible. He proposed the problem, guided me throughout the project, and generously spent many hours discussing drafts and explaining important ideas. I am also thankful to Haohao Liu, Chenji Fu, Zhixing Chen, Vincent Ariksoy, Aaron Beumer, Roben de Preter, Andreas Kretschmer, Andrés Rojas, Søren Gammelgaard, Paweł Borówka, and Gavril Farkas for their insightful discussions and valuable comments. Funding is provided by the DFG (German Research Foundation) under Germany's Excellence Strategy – The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project ID: 390685689).

## 2. TANGENT GAUSS MAP AND CONORMAL GAUSS MAP

We work over the base field  $\kappa = \mathbb{C}$ , and by a variety we mean a integral separated scheme of finite type over  $\mathbb{C}$ . Let  $A/\mathbb{C}$  be an abelian variety of dimension  $n$ , and let  $Z \subseteq A$  be an irreducible closed subvariety of dimension  $r$ . We denote by  $\iota_Z : Z \hookrightarrow A$  the inclusion morphism.

### 2.1. Gauss map and monodromy group.

**Definition 2.1.** *For a subvariety  $Z \subset A$ , the tangent Gauss map of  $Z$  is the map*

$$\phi_Z : Z^{\text{sm}} \longrightarrow \text{Gr}(r, T_0 A) \quad p \longmapsto T_p Z \subseteq T_p A \cong T_0 A$$

*sending a smooth point to the tangent space at that point (seen as a subspace of  $T_0 A$ ).*

*Remark 2.2.* Any map  $X \longrightarrow \text{Gr}(r, V)$  is induced by a rank  $r$  vector bundle  $\mathcal{E}$  on  $X$  together with an epimorphism  $V^* \otimes \mathcal{O}_X \twoheadrightarrow \mathcal{E}$ . In our case, the map  $\phi_Z$  is induced by the tangent bundle  $\mathcal{T}_{Z^{\text{sm}}}$  and the sections

$$T_0^* A \otimes_{\mathbb{C}} \mathcal{O}_{Z^{\text{sm}}} \rightarrow H^0(Z^{\text{sm}}, \mathcal{T}_{Z^{\text{sm}}}^*) \otimes_{\mathbb{C}} \mathcal{O}_{Z^{\text{sm}}} \twoheadrightarrow \mathcal{T}_{Z^{\text{sm}}}^*.$$

The definition of the conormal Gauss map requires a brief recollection of the conormal variety. On the smooth locus, the normal and conormal bundles are the vector bundles  $\mathcal{N}_{Z^{\text{sm}}/A}$  resp.  $\mathcal{N}_{Z^{\text{sm}}/A}^*$  defined by the short exact sequences

$$0 \longrightarrow \mathcal{T}_{Z^{\text{sm}}} \longrightarrow \mathcal{T}_A|_{Z^{\text{sm}}} \longrightarrow \mathcal{N}_{Z^{\text{sm}}/A} \longrightarrow 0$$

$$0 \longrightarrow \mathcal{N}_{Z^{\text{sm}}/A}^* \longrightarrow \Omega_A|_{Z^{\text{sm}}} \longrightarrow \Omega_{Z^{\text{sm}}} \longrightarrow 0$$

We write  $\Lambda_{Z^{\text{sm}}}$  for the total space of  $\mathcal{N}_{Z^{\text{sm}}/A}^*$ . The conormal variety  $\Lambda_Z$  is the closure of  $\Lambda_{Z^{\text{sm}}}$  viewed as a subvariety in  $T^*A$ :

$$\Lambda_Z := \overline{\Lambda_{Z^{\text{sm}}}} \subset T^*A \cong A \times T_0^*A$$

It is a conical Lagrangian subvariety of  $T^*A$ . The (affine) conormal Gauss map is defined as

$$\gamma_Z^{\text{aff}} : \Lambda_Z \subset A \times T_0^*A \longrightarrow T_0^*A$$

For intersection-theoretic purposes it is more natural to work with projectivized conormal spaces. We therefore pass from the affine conormal Gauss map to its projectivized version:

**Definition 2.3.** *The projectivized conormal variety of a subvariety  $Z \subset A$  is defined by*

$$\mathbb{P}\Lambda_Z := \overline{\mathbb{P}\Lambda_{Z^{\text{sm}}}} \subset \mathbb{P}T^*A \cong A \times \mathbb{P}T_0^*A,$$

and the (projectivized) conormal Gauss map is defined by

$$\gamma_Z : \mathbb{P}\Lambda_Z \subset A \times \mathbb{P}T_0^*A \longrightarrow \mathbb{P}T_0^*A.$$

By [8, Theorem 2.8 (1)],  $\gamma_Z$  is generically finite when  $Z$  is (an integral variety) of general type. A lot of geometry of  $Z$  is encoded in the map  $\gamma_Z$ . For instance, if  $Z \subset A$  is smooth, then

$$\deg \gamma_Z = (-1)^r \chi(Z),$$

where  $\chi(Z) = \sum_i (-1)^i b_i(Z)$  is the topological Euler characteristic of  $Z$ , see [12, Theorem 1.1.1].

Further insight can be gained by analyzing the fibers of  $\gamma_Z$ . For instance, if  $Z$  is preserved by a translation  $t_v : A \longrightarrow A$ , then each fiber  $\gamma_Z^{-1}(\xi) \subset A$  is also invariant under  $t_v$ . Likewise, if  $Z = -Z$ , then the fiber satisfies  $\gamma_Z^{-1}(\xi) = -\gamma_Z^{-1}(\xi)$ . Finding further constraints is more challenging.

An important invariant arising from the fiber  $\gamma_Z^{-1}(\xi)$  is the monodromy group  $\text{Gal}(\gamma_Z)$ ; for completeness, we recall its definition below.

**Definition 2.4.** *Let  $f : Y \rightarrow X$  be a generically finite morphism between algebraic varieties. Then there exists a non-empty open subset  $U \subseteq X$  such that the restriction  $f^{-1}(U) \rightarrow U$  is a finite étale cover. Moving along a closed loop in  $U$  induces a permutation of the points in the fiber  $f^{-1}(\xi_0)$ , which defines the map<sup>1</sup>*

$$\rho_f : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(f^{-1}(\xi_0)) \cong S_{\deg f}.$$

The monodromy group is then defined as the image of  $\rho_f$ , i.e.,

$$\text{Gal}(f) := \text{Im } \rho_f.$$

When  $U$  is smooth, the isomorphism class of  $\text{Gal}(f)$  doesn't depend on the choice of  $U$ .

**2.2. Interpolation via hyperplanes.** In this subsection, we reinterpret  $\gamma_Z$  using a functorial and more transparent framework. This permits to define the conormal Gauss map and its monodromy group for any morphism  $\phi_Z : Z \rightarrow \text{Gr}(r, n)$  with  $\dim Z = r$ , and clarifies the relation between the monodromy group and  $\deg \phi_Z$ .

To simplify notation, we abbreviate  $T_0^*A$  by  $W$ . Since each non-zero conormal vector  $\xi \in T_0^*A$  determines a hyperplane  $H_\xi \in \text{Gr}(n-1, T_0A)$ , we have the isomorphisms

$$\begin{aligned} \mathbb{P}T_0^*A &\cong \text{Gr}(n-1, T_0A), \\ \mathbb{P}\Lambda_{Z^{\text{sm}}} &= \{ (p, \xi) \in Z^{\text{sm}} \times \mathbb{P}T_0^*A \mid \xi|_{T_p Z} \equiv 0 \} \\ &\cong \{ (p, H) \in Z^{\text{sm}} \times \text{Gr}(n-1, W) \mid \phi_Z(p) \subseteq H \} \\ &\cong (\phi_Z, \text{Id})^{-1} I_{r, n-1}, \end{aligned}$$

where

$$I_{r, n-1} := \{ (V, H) \in \text{Gr}(r, W) \times \text{Gr}(n-1, W) \mid V \subseteq H \}$$

is the incidence variety relating  $\text{Gr}(r, W)$  and  $\text{Gr}(n-1, W)$ . In these terms,

$$\begin{aligned} \gamma_Z^{-1}(H) \cap Z^{\text{sm}} &= \{ p \in Z^{\text{sm}} \mid \phi_Z(p) \subseteq H \} \\ &\cong \phi_Z^{-1}(\text{Gr}(r, H)) \end{aligned}$$

is the collection of points whose tangent spaces lie entirely within  $H$ .

<sup>1</sup>In the last isomorphism we implicitly give a labelling from  $\{1, 2, \dots, \deg f\}$  to the fiber  $f^{-1}(\xi)$ .

Geometrically, the monodromy can be described as follows: Given a general hyperplane  $H \in \mathbb{P}T_0^*A$ , its preimage under  $\gamma_Z$  consists of  $d$  points  $p_1, \dots, p_d$ . Moving  $H$  continuously along a closed loop we obtain a permutation of these points, and the monodromy group  $\text{Gal}(\gamma_Z)$  consists of all permutations obtained this way. With this formulation, it suffices to consider the Gauss map  $\phi_Z$  alone; the inclusion  $\iota_Z : Z \rightarrow A$  is no longer required for computing the monodromy group.

**Definition 2.5.** *Let  $Z$  be an  $r$ -dimensional variety and  $\phi : Z \rightarrow \text{Gr}(r, n)$  a morphism. Suppose that  $\phi_*[Z] \cup [\text{Gr}(r, H)] = d \neq 0$  in  $H^{2r(n-r)}(\text{Gr}(r, n); \mathbb{Q})$ . The monodromy group  $\text{Mon}(\phi)$  is defined as  $\text{Gal}(f_\phi)$ , where*

$$f_\phi : (\phi, \text{Id})^{-1}I_{r, n-1} \subset Z \times \text{Gr}(n-1, n) \xrightarrow{\pi_2} \text{Gr}(n-1, n)$$

is a generically finite morphism of degree  $d$ .

When  $\phi$  is not generically injective, the monodromy group is subject to additional constraints, as captured by the next lemma.

**Lemma 2.6.** *Let  $\phi : Z \rightarrow \text{Gr}(r, n)$  be generically  $k$ -to-1 onto its image. With a suitable ordering of the fibers, the associated monodromy group  $\text{Mon}(\phi)$  is contained in a wreath product*

$$S_k \wr S_{d/k} := \left( S_k^{\oplus d/k} \right) \rtimes S_{d/k}.$$

*Proof.* For a general hyperplane  $H$ , consider the Cartesian diagram below:

$$\begin{array}{ccccc} \phi : & Z & \xrightarrow{k:1} & \text{Im } Z & \hookrightarrow & \text{Gr}(r, n) \\ & \cup & & \cup & & \cup \\ & \{p_1, \dots, p_d\} & \longrightarrow & \{q_1, \dots, q_{d/k}\} & \longrightarrow & \text{Gr}(r, H) \end{array}$$

The fiber  $\phi^{-1}(\text{Gr}(r, H_0))$  splits into  $d/k$  groups of points, with the monodromy group acting by permutations within each group and among the groups.  $\square$

**2.3. Interpolation via Albanese morphisms.** Examples of subvarieties of abelian varieties can be obtained in two ways: one may fix an abelian variety and consider cycles within it, or begin with a variety  $Z$  and construct a map to an abelian variety, such as the Albanese map. In the latter perspective, the setting changes slightly:  $X$  is a smooth projective variety of dimension  $r$ ,  $\iota_X : X \rightarrow A$  is a morphism to an abelian variety  $A$  of dimension  $n$ , and  $Z := \iota_X(X)$  denotes its image in  $A$ .

In this subsection, we begin with the Albanese morphism  $X \rightarrow \text{Alb}(X)$ , from which we derive some basic properties of the tangent Gauss map. Any other morphism to an abelian variety factors over the Albanese morphism, and then analogous methods apply, see Proposition 2.10.

Let  $X$  be a smooth complex projective variety of dimension  $r$ , and set  $n := \dim_{\mathbb{C}} H^0(X, \Omega_X) = h^{1,0}$ . Recall that the Albanese variety of  $X$  is defined as

$$\text{Alb}(X) := H^0(X, \Omega_X)^* / H_1(X, \mathbb{Z})_{\text{free}},$$

and the Albanese map is given by (for some fixed base point  $p_0 \in X$ )

$$\alpha : X \rightarrow \text{Alb}(X) \quad p \mapsto \left[ \omega \mapsto \int_{\gamma: p_0 \sim p} \omega \right].$$

One classical question is the dimension of  $\alpha(X)$ . Before proceeding, we fix the following notation: For any closed subscheme  $S \subset X$ , let  $\mathcal{I}_S \subset \mathcal{O}_X$  be its ideal sheaf. Given a point  $p \in X$  with inclusion  $i_p : p \hookrightarrow X$ , we denote by  $i_{p,*}\mathcal{O}_p$  the skyscraper sheaf at  $p$ . We set  $\Omega_X(-p) := \Omega_X \otimes \mathcal{I}_p$ , and denote by  $\Omega_X|_p := \Omega_X \otimes i_{p,*}\mathcal{O}_p = i_{p,*}i_p^*\Omega_X$  the fiber of  $\Omega_X$  at  $p$ . With this notation, the cotangent map of  $\alpha$  at  $p \in X$  is

$$T_p^*\alpha : T_{\alpha(p)}^*\text{Alb}(X) = H^0(X, \Omega_X) \rightarrow T_p^*X = H^0(X, \Omega_X|_p).$$

Consider the short exact sequence of coherent sheaves on  $X$

$$0 \longrightarrow \Omega_X(-p) \longrightarrow \Omega_X \longrightarrow \Omega_X|_p \longrightarrow 0$$

which induces a long exact sequence

$$\begin{array}{ccccccc} & & \mathrm{H}^1(X, \Omega_X(-p)) & \longrightarrow & \mathrm{H}^1(X, \Omega_X) & \longrightarrow & 0 \\ & \nearrow & & & & & \\ 0 & \longrightarrow & \mathrm{H}^0(X, \Omega_X(-p)) & \longrightarrow & \mathrm{H}^0(X, \Omega_X) & \xrightarrow{T_p^* \alpha} & \mathrm{H}^0(X, \Omega_X|_p) \end{array}$$

The proposition below follows from standard arguments in homological algebra:

**Proposition 2.7.** *For a general point  $p \in X$ ,*

$$\begin{aligned} \dim_{\mathbb{C}} \alpha(X) &= \operatorname{rank} T_p^* \alpha \\ &= n - h^0(X, \Omega_X(-p)) \\ &= h^1(X, \Omega_X(-p)) - h^1(X, \Omega_X). \end{aligned}$$

*In particular,*

$$\begin{aligned} \alpha \text{ is surjective} &\iff h^0(X, \Omega_X(-p)) = 0 \\ \alpha \text{ is generically finite onto its image} &\iff h^0(X, \Omega_X(-p)) = n - r \\ \alpha \text{ is constant} &\iff h^0(X, \Omega_X(-p)) = n \\ &\iff h^1(X, \Omega_X(-p)) = h^1(X, \Omega_X) \\ &\iff n = 0. \end{aligned}$$

*Proof.* We know that

$$\begin{aligned} \alpha \text{ is surjective} &\iff \dim_{\mathbb{C}} \alpha(X) = n \\ \alpha \text{ is generically finite onto its image} &\iff \dim_{\mathbb{C}} \alpha(X) = r \\ \alpha \text{ is constant} &\iff \dim_{\mathbb{C}} \alpha(X) = 0 \\ &\iff \operatorname{Alb}(X) = 0. \end{aligned}$$

□

We will concentrate on the case where  $\alpha$  is generically injective.<sup>2</sup> Under this assumption, we set  $Z = \iota_X(X)$  and  $A = \operatorname{Alb}(X)$ . The corresponding Gauss map is then a rational map:

$$\begin{array}{ccc} \phi_Z : Z & \dashrightarrow & \operatorname{Gr}(r, T_0 A) \cong \operatorname{Gr}(n-r, \mathrm{H}^0(X, \Omega_X)) \\ p & \longmapsto & \mathrm{H}^0(X, \Omega_X(-p)) \end{array}$$

and we have the isomorphisms

$$\begin{aligned} \mathbb{P}T_0^* A &\cong \mathbb{P}\mathrm{H}^0(X, \Omega_X), \\ \mathbb{P}\Lambda_Z &= \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^* A \mid \omega(p) = 0 \} \\ &\cong \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^* A \mid \omega \in \mathrm{H}^0(X, \Omega_X(-p)) \} \\ &\cong (\phi_Z, \operatorname{Id})^{-1} I_{n-r,1}, \end{aligned}$$

where

$$I_{n-r,1} := \{ (V, [\omega]) \in \operatorname{Gr}(n-r, n) \times \operatorname{Gr}(1, n) \mid \omega \in V \}$$

is the incidence variety relating  $\operatorname{Gr}(n-r, n)$  and  $\operatorname{Gr}(1, n)$ . In that case,

$$\gamma_Z^{-1}([\omega]) = \{ p \in X \mid \omega(p) = 0 \}$$

<sup>2</sup>The general method remains valid in the broader setting, but the Gauss map then takes values in a different space. In fact, there is always a semicontinuous map

$$\phi_X : X \longrightarrow \bigsqcup_{i=0}^r \operatorname{Gr}(n-i, \mathrm{H}^0(X, \Omega_X)) \quad p \longmapsto \mathrm{H}^0(X, \Omega_X(-p)).$$

is the zero set of the section  $\omega \in H^0(X, \Omega_X)$ . The number  $(-1)^r \deg \gamma_Z$  is the index (of the vector field) in the Poincaré–Hopf index formula (see [15, p35]), and the monodromy group  $\text{Gal}(\gamma_Z)$  serves as a more refined invariant.

The next proposition shows when the Gauss map  $\phi_Z$  is not generically injective. We denote by  $X^{[m]}$  the Hilbert scheme of  $m$ -points on  $X$ .

**Proposition 2.8.** *When  $\alpha$  is generically finite onto its image,*

*$\phi_Z$  is not generically injective*

$\iff$  *For general  $p \in X$ , there exists  $q \neq p$  such that  $h^0(X, \Omega_X(-p-q)) = n-r$*

$\stackrel{(1)}{\iff}$  *For general  $p \in X$ , there exists  $S \in X^{[2]}$  such that  $p \in S$  and  $h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r$*

$\stackrel{(2)}{\iff}$  *For all  $p \in X$ , there exists  $S \in X^{[2]}$  such that  $p \in S$  and  $h^0(X, \Omega_X \otimes \mathcal{I}_S) \geq n-r$ .*

*Proof.* When  $n = r$ , the map  $\varphi_Z$  has a point as its target, so the equivalence is trivial. We shall thus restrict to the case  $n > r$  in the subsequent discussion.

(1): The implication “ $\Rightarrow$ ” is immediate. For the converse “ $\Leftarrow$ ”, it suffices to show that the image of

$$\left\{ S \in X^{[2]} \mid h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r \right\}$$

under the map  $\pi_X : X^{[2]} \rightarrow X^{(2)}$  does not include the diagonal. Indeed, we can choose a section  $s \in H^0(X, \Omega_X)$  that cuts out finitely many reduced points  $p_1, \dots, p_d$  in  $X$ . (Well this maybe not so true, since  $\phi_Z$  is not always regular, such a section  $s$  may vanish along some indeterminacy locus. However, for a general  $s$ , there will always be at least one isolated zero  $p_1$ , which is sufficient for our purposes.) Then for any  $S \in \pi_X^{-1}(2p_1)$ , we have

$$H^0(X, \Omega_X \otimes \mathcal{I}_S) \subsetneq H^0(X, \Omega_X(-p_1)).$$

(2): The implication “ $\Rightarrow$ ” follows from the geometry of the projection  $\text{pr}_1 : X \times X^{[2]} \rightarrow X$ . By Lemma 2.9 and the closedness of the tautological correspondence in  $X \times X^{[2]}$ , the subset

$$I^{n-r} := \left\{ (p, S) \in X \times X^{[2]} \mid p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) \geq n-r \right\}$$

is closed in  $X \times X^{[2]}$ . Consequently, its image

$$\text{pr}_1(I^{n-r}) = \left\{ p \in X \mid \text{there exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r \right\}$$

is closed in  $X$ . Since the hypothesis ensures that  $\text{pr}_1(I^{n-r})$  contains a nonempty open subset of  $X$ , we obtain  $\text{pr}_1(I^{n-r}) = X$ .

For “ $\Leftarrow$ ”: as  $\alpha$  is generically injective, one has  $h^0(X, \Omega_X(-p)) = n-r$  for a general point  $p \in X$ . Therefore, for any subscheme  $S$  containing such a general point,  $h^0(X, \Omega_X \otimes \mathcal{I}_S) \leq n-r$ .  $\square$

**Lemma 2.9.** *Let  $m \in \mathbb{Z}_{>0}$ . The function*

$$h^\Omega : X^{[m]} \longrightarrow \mathbb{Z}_{\geq 0} \quad S \longmapsto h^0(X, \Omega_X \otimes \mathcal{I}_S)$$

*is Zariski upper semicontinuous.*

*Proof.* Consider the coherent sheaf  $\mathcal{F} \in \text{Coh}(X^{[m]} \times X)$  characterized by the property

$$\mathcal{F}|_{\{S\} \times X} \cong \Omega_X \otimes \mathcal{I}_S,$$

The proposition then follows directly from the semicontinuity theorem [16, 28.1.1].  $\square$

The stratification of  $X^{[m]}$  by  $h^\Omega$  offers a natural generalization of Brill–Noether theory beyond the setting of curves.

At the end of this subsection, let us turn to the setting of a general abelian variety  $A$  and a smooth subvariety  $\iota_Z : Z \hookrightarrow A$ , where  $n = \dim_{\mathbb{C}} A$  and  $r = \dim_{\mathbb{C}} Z$ . Observe that  $\iota_Z$  factors through the Albanese variety of  $Z$ :

$$\iota_Z : Z \xrightarrow{\alpha_Z} \text{Alb}(Z) \xrightarrow{\pi} A$$

We shall also assume that  $Z$  generates  $A$ ; it then follows that the map  $\pi$  is surjective. The cotangent map of  $\iota_Z$  at a point  $p \in Z$  factors through  $H^0(Z, \Omega_Z)$ :

$$T_p^* \iota_Z : T_p^* A \hookrightarrow H^0(Z, \Omega_Z) \longrightarrow T_p^* Z$$

For convenience, abbreviate  $V := T_0^* A \cong T_p^* A$ , and view  $V$  as a subspace of  $H^0(Z, \Omega_Z)$ .

**Proposition 2.10.** *Assume that  $Z$  is embedded in  $A$  and generates  $A$ , and let  $V := T_0^* A$ . Then*

$$\dim_{\mathbb{C}} H^0(Z, \Omega_Z(-p)) \cap V = n - r \quad \text{for all } p \in Z$$

*It follows that the Gauss map is a regular morphism*

$$\begin{array}{ccc} \phi_Z : Z & \longrightarrow & \text{Gr}(r, T_0 A) \cong \text{Gr}(n - r, V) \\ p & \longmapsto & H^0(Z, \Omega_Z(-p)) \cap V \end{array}$$

Furthermore,

$\phi_Z$  is not generically injective

$\iff$  For general  $p \in Z$ , there exists  $q \neq p$  such that  $h^0(Z, \Omega_Z(-p - q)) \cap V = n - r$

$\iff$  For all  $p \in Z$ , there exists  $S \in Z^{[2]}$  such that  $p \in S$  and  $h^0(Z, \Omega_Z \otimes \mathcal{I}_S) \cap V = n - r$ .

### 3. MONODROMY GROUP

#### 3.1. Criteria for big monodromy group.

**Definition 3.1** (big monodromy group). *We refer to the big monodromy group as any group of the following types:*

notation	name	alias
$W(A_{m+1}) = S_m$	full symmetric group	
$W(C_m) = S_2^{\oplus m} \rtimes S_m$	signed symmetric group	hyperoctahedral group
$W(D_m) = (S_2^{\oplus m})_0 \rtimes S_m$	even-signed symmetric group	demihyperoctahedral group

TABLE 1. big monodromy group

*In practice, the term “big monodromy group” refers to the full symmetric group  $S_n$  when the subset  $Z \subset A$  is not symmetric, and to the (even-)signed symmetric group when  $Z \subset A$  is symmetric.*

**Proposition 3.2** (See [2, p111] for a detailed proof). *Suppose that  $\iota_C : C \subseteq \mathbb{P}^{n-1}$  is an irreducible nondegenerate<sup>3</sup> curve of degree  $d$ , then  $\text{Mon}(\iota_C) \cong S_d$ .*

*Sketch of proof.* Because  $S_d$  is generated by its transpositions, we are reduced to verifying that:

- $\text{Mon}(\iota_C)$  acts doubly transitively on the fiber;
- $\text{Mon}(\iota_C)$  contains a transposition.

□

In fact, a degree 2 : 1 map does not give rise to any exceptional monodromy groups beyond those listed in Table 1.

<sup>3</sup>A curve  $C \subseteq \mathbb{P}^{n-1}$  is said to be nondegenerate if it is not contained in any hyperplane  $H \subseteq \mathbb{P}^{n-1}$ .

**Proposition 3.3.** *Let  $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$  be an irreducible nondegenerate curve of degree  $d/2$ , and let  $h : C \rightarrow C'$  be a degree 2 ramified covering. Then*

$$\text{Mon}(\iota_{C'} \circ h) \cong W(C_{d/2}) \text{ or } W(D_{d/2}).$$

*Sketch of proof.* By Lemma 2.6 we know that  $\text{Mon}(\iota_{C'} \circ h) \subseteq W(C_{d/2})$ . By Lemma 3.4, we are reduced to verifying that:

- The quotient map  $\text{Mon}(\iota_{C'} \circ h) \rightarrow \text{Mon}(\iota_{C'}) \cong S_{d/2}$  is surjective;
- (signed doubly transitive)  $\text{Mon}(\iota_{C'} \circ h)$  acts transitively on pairs  $(x, y)$  with  $x \neq \pm y$ .

□

**Lemma 3.4.** *Let  $G$  be a subgroup of  $W(C_m)$ , acting naturally on the set  $\pm 1, \dots, \pm m$ . If the projection  $G \rightarrow S_m$  is surjective then*

$$G \cong W(C_m) \text{ or } W(D_m) \text{ or } S_2 \times S_m \text{ or } S_m.$$

*Sketch of proof.* Let  $H$  denote the kernel of the natural quotient map  $G \rightarrow S_m$ . Then  $H \subseteq (S_2)^{\oplus m}$  is stable under the action of  $S_m$ . There are only four possible forms that  $H$  can take:<sup>4</sup>

- $H = 0$ . Then  $G \cong S_m$ .
- $H = \langle (-1, \dots, -1) \rangle \cong S_2$ . Then  $G \cong S_2 \times S_m$ .
- $H = (S_2^{\oplus m})_0$ . Then  $G$  is a index 2 subgroup of  $W(C_m)$ , so  $G \cong W(D_m)$ .<sup>5</sup>
- $H = S_2^{\oplus m}$ . Then  $G = W(C_m)$ .

□

By incorporating further information about the covering, we are able to determine the monodromy group.

**Proposition 3.5.** *Let  $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$  be an irreducible nondegenerate curve of degree  $d/2$ , and let  $h : C \rightarrow C'$  be a degree 2 ramified covering, with ramification occurring at at least one smooth point of  $C'$ . Then  $\text{Mon}(\iota_{C'} \circ h) \cong W(C_{d/2})$  is the hyperoctahedral group/signed symmetric group.*

*Sketch of proof.* By Lemma 2.6 we know that  $\text{Mon}(\iota_{C'} \circ h) \subseteq W(C_{d/2})$ . By Lemma 3.6, we are reduced to verifying that:

- The quotient map  $\text{Mon}(\iota_{C'} \circ h) \rightarrow \text{Mon}(\iota_{C'}) \cong S_{d/2}$  is surjective;
- $\text{Mon}(\iota_{C'} \circ h)$  contains a transposition of a pair of points in the fiber of  $h$ .

□

**Lemma 3.6.** *Let  $G$  be a subgroup of  $W(C_m)$ , acting naturally on the set  $\pm 1, \dots, \pm m$ . If the projection  $G \rightarrow S_m$  is surjective and the transposition  $\sigma_0$  of  $\pm 1$  lies in  $G$ , then  $G = W(C_m)$ .*

*Sketch of proof.* Let  $\varepsilon_i$  denote the transposition of  $\pm i$ . For any  $\sigma \in S_m$ , choose a lift  $\tilde{\sigma} \in G$ , then

$$\varepsilon_{\sigma(1)} = \tilde{\sigma} \circ \sigma_0 \circ \tilde{\sigma}^{-1} \in G.$$

Thus,  $S_2^{\oplus m} \subset G$ , and since  $G$  maps onto  $S_m$ , we obtain  $G = W(C_m)$ . □

*Proof that Conjecture 1.1 is equivalent to Conjecture 1.2 when  $n > 2$ .* If  $\phi_C$  is generically injective, Proposition 3.2 yields  $\text{Gal}(\gamma_C) \cong S_d$ . When  $C = -C$  and  $\phi_C$  is a double cover, Proposition 3.3 implies  $\text{Gal}(\gamma_C) \cong W(C_{d/2})$  or  $W(D_{d/2})$ , both of which are big monodromy groups. In all other cases, Lemma 2.6 shows that  $\text{Gal}(\gamma_C) \subseteq S_k \wr S_{d/k}$ , which is not big whenever  $k > 2$  and  $k \neq d$ . Note that  $k = d$  occurs only when  $n = 2$ . □

<sup>4</sup>Here is a brief argument showing that  $H$  must be one of  $0$ ,  $S_2$ ,  $(S_2^{\oplus m})_0$ , or  $S_2^{\oplus m}$ . If  $H$  contains an element  $h = (a_1, \dots, a_m)$  with  $a_i \neq a_j$  for some  $i \neq j$ , then

$$(ij)h + h = (1, \dots, \underset{i\text{-th}}{\uparrow} -1, \dots, \underset{j\text{-th}}{\uparrow} -1, \dots, 1) \in H,$$

implying  $(S_2^{\oplus m})_0 \subseteq H$ . Hence,  $H$  must be either  $(S_2^{\oplus m})_0$  or  $S_2^{\oplus m}$ . Otherwise, if all  $h \in H$  have identical coordinates, then  $H$  is either  $0$  or  $S_2$ .

<sup>5</sup>Check [stackexchange discussions](#)

**3.2. Curves with big monodromy group.** The availability of these criteria permits the systematic construction of numerous cases where the associated monodromy group is big.

**Example 3.7.** Let  $C$  be a smooth curve of genus  $g$  embedded in its Jacobian  $A := \text{Jac}(C)$  via the Abel–Jacobi map  $\text{AJ}_C : C \hookrightarrow A$ . In this case, the tangent Gauss map is the canonical map  $|\omega_C|$ .

When  $C$  is non-hyperelliptic, the corresponding Gauss map

$$|\omega_C| : C \longrightarrow \mathbb{P}^{g-1}$$

makes  $C$  as an irreducible nondegenerate curve of degree  $2g - 2$ , by Proposition 3.2 we get

$$\text{Gal}(\gamma_C) \cong S_{2g-2}.$$

When  $C$  is hyperelliptic, the corresponding Gauss map is  $2 : 1$  onto a rational normal curve  $R \subset \mathbb{P}^{g-1}$ :

$$|\omega_C| : C \xrightarrow{2:1} R \hookrightarrow \mathbb{P}^{g-1}$$

By Proposition 3.5 we get

$$\text{Gal}(\gamma_C) \cong S_2^{\oplus g-1} \rtimes S_{g-1}.$$

The Prym case is more intricate, since the associated monodromy group may fail to be large.

**Setting 3.8.** Let  $C'/\mathbb{C}$  be a smooth projective curve,  $\eta \in \text{Pic}(C')$  a line bundle, and  $B$  a reduced effective (possibly zero) divisor on  $C'$ . Suppose  $k > 1$  and that there is an isomorphism  $\eta^{\otimes k} \cong \mathcal{O}_{C'}(B)$ . According to [3, §17], these data determine a cyclic  $k$ -fold covering  $h : C \rightarrow C'$  of smooth projective curves, branched exactly along  $B$ . Denote by  $\sigma$  the generator of the Galois group  $\text{Gal}(C/C') \cong \mathbb{Z}/k\mathbb{Z}$ .

Recall that the Prym variety  $A := \text{Prym}(C/C')$  is defined as the connected component of the identity in

$$\ker [\text{Nm} : \text{Jac}(C) \longrightarrow \text{Jac}(C')]$$

Its dimension is denoted by  $n$ .

The Abel–Prym map is defined by

$$\text{AP}_{C/C'} : C \longrightarrow A \quad p \longmapsto \mathcal{O}_C(p - \sigma(p)).$$

The following example, which essentially restates [5, Corollary 2.2], illustrates the case where the cover is an étale double cover.

**Proposition 3.9.** In Setting 3.8, assume further that  $k = 2$ ,  $B = \emptyset$ , and that  $C'$  is non-hyperelliptic of genus  $g(C') \geq 4$ .

- If  $C'$  is bielliptic and  $\eta = \text{pr}^* \eta_0$ , where  $\text{pr} : C' \rightarrow E'$  is a bielliptic map and  $\eta_0 \in \text{Pic}^0(E')[2]$ , then the tangent Gauss map  $\phi_C : C \rightarrow \mathbb{P}^{n-1}$  has degree 4, and the curve  $C \subset A$  is invariant under a 2-torsion translation in  $A$ .
- Otherwise, the tangent Gauss map  $\phi_C : C \rightarrow \mathbb{P}^{n-1}$  is a double cover onto its image, and  $\text{Gal}(\gamma_C)$  is big.

*Proof.* When  $k = 2$ , the corresponding Gauss map  $\gamma_C$  factors through  $h$ :

$$\begin{aligned} \gamma_C : C &\xrightarrow{h} C' \xrightarrow{|\omega_{C'} \otimes \eta|} \mathbb{P}^{n-1} = \mathbb{P}(\text{H}^0(\omega_{C'} \otimes \eta)^*) \\ &\cong \text{Gr}(n-1, \text{H}^0(\omega_{C'} \otimes \eta)) \\ \tilde{p} &\longmapsto p \longmapsto \text{H}^0(\omega_{C'} \otimes \eta(-p)) \end{aligned}$$

where  $|\omega_{C'} \otimes \eta|$  denotes the Prym–canonical map. According to [5, Corollary 2.2], one has  $\deg |\omega_{C'} \otimes \eta| \in \{1, 2\}$ , and the case  $\deg |\omega_{C'} \otimes \eta| = 2$  occurs precisely when  $C'$  is bielliptic and  $\eta = \text{pr}^* \eta_0$ . In

this situation, let  $h_E: E \rightarrow E'$  denote the unramified double covering corresponding to  $\eta_0$ , and let  $\iota \in \text{Gal}(E/E')$  be the associated involution. The following diagram is Cartesian:

$$\begin{array}{ccc} & \eta & \\ & \swarrow & \searrow \\ C' & & E \\ \text{pr} \swarrow & & \nwarrow \eta_0 \\ & E' & \end{array} \quad \begin{array}{ccc} & C & \\ h \swarrow & & \searrow p \\ C' & & E \\ \text{pr} \swarrow & & \nwarrow h_E \\ & E' & \end{array}$$

For any  $p_0 \in E$ , set

$$\mathcal{L}_0 := \mathcal{O}_E(p_0 - \iota(p_0)) \in \text{Pic}^0(E)[2],$$

which does not depend on the choice of  $p_0$ . The curve  $C \subset A$  is then invariant under translation by  $p^*\mathcal{L}_0 \in A$ .  $\square$

We now examine all possible cases of double covers. Motivated by [14, Lemma 2.1], we formulate the following result:

**Lemma 3.10.** *Let  $C'$  be a smooth projective curve of genus  $g(C') \geq 2$ , and let  $\eta \in \text{Pic}(C')$  be a line bundle of degree  $s \geq 0$ . Then the linear system  $|\omega_{C'} \otimes \eta|$  is not generically injective if and only if one of the following holds:*

- (a)  $s = 2$  and for all  $x \in C'$ , there exists  $y \in C'$  such that  $\eta \cong \mathcal{O}_{C'}(x + y)$ ;
- (b)  $s = 1$  and for all  $x \in C'$ , there exist  $y, u \in C'$  such that  $\eta \cong \mathcal{O}_{C'}(x + y - u)$ ;
- (c)  $s = 0$  and for all  $x \in C'$ , there exist  $y, u, v \in C'$  such that  $\eta \cong \mathcal{O}_{C'}(x + y - u - v)$ .

Furthermore, in cases (a) and (b), the curve  $C'$  must be hyperelliptic.

*Proof.* This is a consequence of Riemann–Roch. In case (a), since  $\eta \in g_2^1$ , the curve  $C'$  is hyperelliptic. In case (b), one obtains a family of  $g_3^1$ 's of the form  $\mathcal{O}_{C'}(x + y + u)$ , which implies that  $\mathcal{O}_{C'}(x + y) \in g_2^1$ .  $\square$

**Corollary 3.11.** *In Setting 3.8, assume further that  $k = 2$ ,  $B \neq \emptyset$ , and that  $C'$  is non-hyperelliptic. Then the Prym-canonical map  $|\omega_{C'} \otimes \eta|$  is generically injective, and therefore  $\text{Gal}(\gamma_C)$  is big.*

In the process of persuing proving Conjecture 1.2, we propose the following lema which will roll out a lot of trivial cases.

**3.3. Curves in Prym variety with small monodromy group(Delete later).** This subsection is devoted to presenting an example that has small monodromy group.

The classical theory treats the case where the Prym variety  $A$  is principally polarized—this occurs exactly when  $B = \emptyset$  or  $B = \{p_0, q_0\}$ , that is, when  $h$  is either unramified or ramified at two points.<sup>6</sup> For clarity, we restrict our attention to these cases. Table 2 lists the relevant numerical data.<sup>7</sup>

	$g(C)$	$g(C')$	$\deg \eta$
$B = \emptyset$	$2n + 1$	$n + 1$	0
$B = \{p_0, q_0\}$	$2n$	$n$	1

TABLE 2. numerical data of Prym pair

The Abel–Prym map  $\text{AP}_{C/C'}$  does not always behave as nicely as the Abel–Jacobi map; it may fail to be an embedding. When  $C$  is hyperelliptic, the map  $\text{AP}_{C/C'}$  fails to be generically injective, and therefore falls outside the scope of our discussion. When  $C$  is non-hyperelliptic, we

<sup>6</sup>See [6, Theorem 3.2.6] for the proof.

<sup>7</sup>As usual,  $n = \dim_{\mathbb{C}} A$  is the dimension of the abelian variety.

may regard it as a subvariety of  $A$  for our purposes, although it may be not strictly embedded.<sup>8</sup> However, it remains unclear whether  $C$  is stable under any translation in  $A$ .

**Example 3.12.** Let  $C'$  be a non-hyperelliptic bielliptic curve, and let  $\text{pr} : C' \rightarrow E$  denote a  $2 : 1$  covering onto an elliptic curve. For any nontrivial 2-torsion line bundle  $\eta_0 \in \text{Pic}^0(E)[2]$ , the pullback  $\eta := \text{pr}^* \eta_0$  satisfies  $\eta^{\otimes 2} \cong \mathcal{O}_{C'}$ , and the map  $|\omega_{C'} \otimes \eta|$  factors through  $\text{pr}$ .

*Proof of Example 3.12.* For any  $x \in C'$ , write  $x_0 := \text{pr}(x)$ , then

$$\deg \eta_0^\vee(x_0) = 1 \implies \text{there exists } y_0 \in E \text{ such that } \eta_0^\vee(x_0) = \mathcal{O}_E(y_0).$$

Write  $\text{pr}^{-1}(x_0) = \{x, x'\}$ ,  $\text{pr}^{-1}(y_0) = \{y, y'\}$ , we get

$$\begin{aligned} \eta_0^\vee(x_0) &= \mathcal{O}_E(y_0) \\ \implies \eta^\vee(x + x') &= \mathcal{O}_{C'}(y + y') && \text{Via pulling back along pr} \\ \implies h^0(\eta^\vee(x + x')) &= 1 \\ \iff h^0(\omega_{C'} \otimes \eta(-x - x')) &= n - 1 && \text{Via Riemann-Roch} \\ \iff H^0(\omega_{C'} \otimes \eta(-x)) &\cong H^0(\omega_{C'} \otimes \eta(-x')) \\ \iff |\omega_{C'} \otimes \eta|(x) &= |\omega_{C'} \otimes \eta|(x') \end{aligned}$$

□

The next proposition tells us, if we put some Galois condition for the covering, then the resulting curve  $C \subset A$  is not invariant under any non-trivial translation of  $A$ .

**Proposition 3.13.** In Example 3.12, let  $C$  be the curve corresponding to  $\eta \in \text{Pic}^0(C')[2]$ . If  $\text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ , then  $C$  admits three intermediate quotients over  $E$ . Denote them by  $C'$ ,  $C_1$ , and  $C_2$ , where  $C_1$  and  $C_2$  are the two intermediate curves different from  $C'$ .

$$\begin{array}{ccc} \begin{array}{c} C \\ \downarrow h \\ C' \\ \downarrow \\ E \\ \downarrow \\ \mathbb{P}^1 \end{array} & & \begin{array}{ccccc} & C & & & \\ & \swarrow h_1 & \downarrow h & \searrow h_2 & \\ C_1 & & C' & & C_2 \\ & \searrow & \downarrow & \swarrow & \\ & E & & & \\ & \downarrow & & & \\ & \mathbb{P}^1 & & & \end{array} \\ \text{Gal}(C/E) \cong \mathbb{Z}/4\mathbb{Z} & & \text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \end{array}$$

If either  $\text{Gal}(C/E) \cong \mathbb{Z}/4\mathbb{Z}$ , or  $\text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$  with the coverings  $h_i : C \rightarrow C_i$  ramified, then the curve  $C \subset A$  is not fixed under any non-trivial translation on  $A$ .

*Proof.* We prove by contradiction. Assume there exists a non-trivial translation  $\sigma : A \rightarrow A$  preserving  $C$ . Then the restriction  $\sigma|_C : C \rightarrow C$  is an automorphism, and the induced map

$$h_\sigma : C \longrightarrow C/\sigma$$

is an unramified double covering. Since  $\sigma$  is a translation, the Gauss map  $\gamma_C$  necessarily factors through  $h_\sigma$ :


$$\begin{array}{ccccc} \gamma_C : C & \xrightarrow{2:1} & C' & \xrightarrow{2:1} & E \subset \mathbb{P}^{n-1} \\ & \searrow h_\sigma & & \nearrow \exists! & \\ & & C/\sigma & & \end{array}$$

<sup>8</sup>For a detailed description of the map  $\text{AP}_{C/C'}$ , see [4, Proposition 12.5.2, Corollary 12.5.6]. Although  $\text{AP}_{C/C'}$  may collapse  $p_0$  and  $q_0$ , this does not affect the relevant computation of  $\text{Gal}(\gamma_Z)$ .

Without loss of generality, we may assume that  $\deg(h_\sigma) = 2$ . By assumption,  $C/\sigma$  coincides with  $C'$ , and  $\sigma|_C$  is the involution  $\iota$  associated with the cover  $h : C \rightarrow C'$ , which extends to an involution  $\tilde{\iota}$  of  $A$ . According to [4, Proposition 12.4.2],  $\tilde{\iota}$  is a reflection. Hence  $\sigma \circ \tilde{\iota}^{-1}$  is another reflection fixing  $C$ , which is impossible.  $\square$

Unfortunately, the condition in Proposition 3.13 is never met, and consequently the curve  $C \subset A$  is preserved by a 2-torsion translation of  $A$ , as explained below.

**Lemma 3.14.** *In Example 3.12, let  $(\mathcal{L}_3, \mathcal{L}_3^{\otimes 2} \cong \mathcal{O}_E(R))$  denote the line bundle associated with the double cover  $C' \rightarrow E$ , where  $R$  is its branch divisor. Set  $\mathcal{L}_1 := \eta_0$ ,  $\mathcal{L}_2 := \eta_0 \otimes \mathcal{L}_3$ , and let  $u_i : C_i \rightarrow E$  be the ramified covering determined by  $\mathcal{L}_i$ .*



- (1) We have  $C_3 = C'$ ,  $g(C_1) = 1$  and  $g(C_2) = g(C_3) = n + 1$ .
- (2) The curve  $C$  arises as the fiber product  $C = C_3 \times_E C_1$ . The covering  $C \rightarrow E$  is Galois with

$$\mathrm{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \hat{=} \{\mathrm{Id}, \sigma_1, \sigma_2, \sigma_3\},$$

where each  $\sigma_i$  is the involution associated with the projection  $h_i : C \rightarrow C_i$  (so that  $\iota = \sigma_3$ ). By the Riemann–Hurwitz formula, the maps  $u_1$ ,  $h_2$ , and  $h_3$  are unramified.

- (3) With  $\sigma = \{\sigma_2, \sigma_3\} \in \mathrm{Gal}(C_1/E)$  the involution of  $u_1$ , one can define

$$\mathcal{L}_0 := \mathcal{O}_{C_1}(p_0 - \sigma(p_0)) \in \mathrm{Pic}^0(C_1)[2],$$

independent of the choice of  $p_0 \in C_1$ . The curve  $C \subset A$  is invariant under the translation corresponding to  $h_1^* \mathcal{L}_0$ . Concretely, for every  $p \in C$ , one has

$$\mathcal{O}_C(p - \sigma_3(p)) \cong \mathcal{O}_C(\sigma_2(p) - \sigma_3\sigma_2(p)) \otimes h_1^* \mathcal{L}_0.$$

*Proof.*

- (1) Since

$$\mathcal{L}_1^{\otimes 2} \cong \mathcal{O}_E, \quad \mathcal{L}_2^{\otimes 2} \cong \mathcal{O}_E(R), \quad \mathcal{L}_3^{\otimes 2} \cong \mathcal{O}_E(R),$$

the Riemann–Hurwitz formula yields the genus of  $C_i$ .

- (2) Notice that the curve  $C$  corresponds to the line bundle  $\eta := \mathrm{pr}^* \eta_0$ .
- (3) Since  $u_1$  is unramified, the involution  $\sigma \in \mathrm{Aut}(C_1/E)$  acts by translation and thus preserves the Jacobian:

$$\mathcal{O}_{C_1}(p_0 - q_0) \cong \mathcal{O}_{C_1}(\sigma(p_0 - q_0)), \quad \text{for any } p_0, q_0 \in C_1.$$

For  $p \in C$ , write  $p_0 = h_1(p)$ , one can compute that

$$\begin{aligned}
 h_1^* \mathcal{L}_0 &\cong h_1^* \mathcal{O}_{C_1}(p_0 - \sigma(p_0)) \\
 &\cong \mathcal{O}_C(p + \sigma_1(p) - \sigma_2(p) - \sigma_3(p)) \\
 &\cong \mathcal{O}_C(p - \sigma_3(p)) \otimes \mathcal{O}_C(-\sigma_2(p) + \sigma_3\sigma_2(p))
 \end{aligned}$$

$\square$

**Lemma 3.15.** *When  $\mathrm{gon}(C') > 4$ ,  $|\omega_{C'} \otimes \eta|$  is injective. As a result, the monodromy group is big.*

*Proof.* Suppose that  $|\omega_{C'} \otimes \eta|$  is not injective, we need to find a line bundle of degree 4 and rank  $\geq 1$ . In fact, for  $p \neq q$ ,

$$\begin{aligned} |\omega_{C'} \otimes \eta|(p) &= |\omega_{C'} \otimes \eta|(q) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-p - q)) &= 1 && \text{By [16, 19.2.8]} \\ \iff h^0(\omega_{C'} \otimes \eta(-p - q)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^{\vee}(p + q)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

When  $B = \emptyset$ , write  $\eta^{\vee}(p + q) = \mathcal{O}_{C'}(p' + q')$ , then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^{\vee})^{\otimes 2}(2p + 2q) = \mathcal{O}_{C'}(2p' + 2q') \in g_4^1;$$

When  $B = \{p_0, q_0\}$ , write  $\eta^{\vee}(p + q) = \mathcal{O}_{C'}(p')$ , then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^{\vee})^{\otimes 2}(2p + 2q + p_0 + q_0) = \mathcal{O}_{C'}(2p' + p_0 + q_0) \in g_4^1.$$

□

*Remark 3.16.* Based on the strategy used in the proof of Lemma 2.10, we can in fact obtain stronger results. For  $p \in C'$ ,

$$\begin{aligned} |\omega_{C'} \otimes \eta| \text{ is ramified at } p & \quad (\text{i.e., the tangent map is 0 at } p) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-2p)) &= 1 && \text{By [16, 19.2.9]} \\ \iff h^0(\omega_{C'} \otimes \eta(-2p)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^{\vee}(2p)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

For the remainder of this discussion, we focus exclusively on the case  $B = \emptyset$ . Combining both,

$$\begin{aligned} |\omega_{C'} \otimes \eta| \text{ is not generically injective} \\ \iff \text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } h^0(\eta^{\vee}(p + q)) &= 1 \\ \iff \text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } \eta^{\vee} + p \in C' + C' - q \\ \iff \eta^{\vee} + C' \subseteq C' + C' - C'. \end{aligned}$$

Gonality is not the only invariant forcing the monodromy group to be large. Indeed, the Castelnuovo–Severi inequality implies that Example 3.12 is the sole instance of a non-trivial small monodromy group when  $g(C') > 9$ .

**Fact 3.17** (Castelnuovo–Severi inequality, [9, p26, Corollary]). *Let  $C$  be a smooth projective curve equipped with two ramified coverings  $f_i : C \rightarrow C_i$  of degrees  $d_i$  ( $i = 1, 2$ ). Suppose that there is no morphism  $h : C \rightarrow \tilde{C}$  with  $\deg(h) > 1$  such that both  $f_1$  and  $f_2$  factor through  $h$ . Then*

$$g(C) \leq d_1 \cdot g(C_1) + d_2 \cdot g(C_2) + (d_1 - 1)(d_2 - 1).$$

**Proposition 3.18.** *If, in Setting 3.8, we additionally require that  $C'$  be non-hyperelliptic and non-bielliptic, and  $h : C \rightarrow C'$  is étale, then any such curve with  $g(C') > 9$  must have big monodromy group.*

*Proof.* Assume that  $C'$  and  $\eta$  satisfies

$$\eta^{\vee} + C' \subseteq C' + C' - C'.$$


**Step 1.** we can find two distinct  $g_4^1$  of  $C'$ .

For any  $p \in C'$ , there exist  $q, p', q' \in C'$  such that

$$\eta^{\vee}(p + q) = \mathcal{O}_{C'}(p' + q').$$

This implies

$$\mathcal{O}_{C'}(2p + 2q) = \mathcal{O}_{C'}(2p' + 2q'),$$



<sup>9</sup>For  $g(C') > 5$ , Castelnuovo–Severi implies that  $C'$  admits exactly one ramified double cover onto an elliptic curve.

#### 4.1. Clean Lagrangian cycles.

**Proposition 4.1.** *All irreducible conic Lagrangian cycles in  $T^*A$  are of the form  $\Lambda_Z$  for some irreducible subvariety  $Z \subset A$ . This yields a one-to-one correspondence between irreducible conic Lagrangian cycles in  $T^*A$  and irreducible subvarieties of  $A$ :*

$$\{ \text{irreducible conic Lagrangian cycles in } T^*A \} \longleftrightarrow \{ \text{irreducible subvarieties in } A \}$$

*Sketch of proof.* For any irreducible conic Lagrangian cycle  $\Lambda \subset T^*A$ , let  $Z$  denote the image of  $\Lambda$  under the natural projection  $T^*A \rightarrow A$ . Our goal is to show that  $\Lambda = \Lambda_Z$ .

- By definition,  $\Lambda \subset T^*A|_Z$ .
- Since  $\Lambda$  is conic, we have  $s(Z) \subset \Lambda$ , where  $s : A \rightarrow T^*A$  denotes the zero section.
- Since  $\Lambda$  is Lagrangian and  $s(Z) \subset \Lambda$ , we have  $\Lambda \subset \Lambda_Z$ .
- Since  $\Lambda_Z$  is irreducible with  $\dim_{\mathbb{C}} \Lambda = \dim_{\mathbb{C}} \Lambda_Z = n$ , we have  $\Lambda = \Lambda_Z$ .

□

Why do we shift attention from  $Z$  to  $\Lambda_Z$  as the main object of study? One reason is the uniformity of  $\Lambda_Z$ : it always has dimension  $n$ , and in most cases, the natural map  $\Lambda_Z \rightarrow T_0^*A$  is generically finite, with fibers lying inside  $A$ .

**Definition 4.2** (Clean cycle). *An irreducible Lagrangian cycle  $\Lambda \subset T^*A$  is called clean if the composed projection*

$$\Lambda \longrightarrow T^*A \twoheadrightarrow T_0^*A$$

*is generically finite.*

Another important reason is that the space of weighted clean conic Lagrangian cycles naturally acquires a convolution structure, arising from the group law on  $A$ , which plays a central role in the analysis.

**Proposition 4.3.** *The group of weighted clean conic Lagrangian cycles*

$$\begin{aligned} \mathcal{L}(A) &:= \{ \text{weighted clean conic Lagrangian cycles in } T^*A \} \\ &= \left\{ \sum_{\substack{Z_i \subset A \\ \text{irr clean}}} n_i \Lambda_{Z_i} \mid n_i \in \mathbb{Z} \right\} \end{aligned}$$

*has a natural convolution structure as follows:*

$$\begin{aligned} \Lambda_{Z_1} \circ \Lambda_{Z_2} &= \text{the clean part of } (a, \text{Id}_{T_0^*A})_* (\Lambda_{Z_1} \times_{T_0^*A} \Lambda_{Z_2}) \\ &= \overline{(a, \text{Id}_U)_* (\Lambda_{Z_1}|_U \times_U \Lambda_{Z_2}|_U)} \end{aligned}$$

*where*

$$U := \left\{ \xi \in T_0^*A \mid \deg \phi_{Z_i} = \# \phi_{Z_i}^{-1}(\xi) \text{ for } i = 1, 2 \right\}$$

*and  $a : A \times A \rightarrow A$  is the addition map in  $A$ . The general convolution is defined by  $\mathbb{Z}$ -linear extension.*

*Sketch of proof.* To establish the claim, it suffices to show that  $\Lambda_{Z_1} \circ \Lambda_{Z_2}$  defines a weighted conic Lagrangian cycle. The conic property follows directly from the definition, while the Lagrangian condition can be verified at a general point  $(p_1 + p_2, \xi) \in \Lambda_{Z_1} \circ \Lambda_{Z_2}$ . □

We now consider the projective versions of all objects involved, so that we may make use of properness. To simplify notation, we abbreviate  $\mathbb{P}T_0^*A$  by  $\mathbb{P}^\vee$ .

**Lemma 4.4.**

(1) Suppose that  $\mathbb{P}\Lambda_{Z_1}, \mathbb{P}\Lambda_{Z_2} \subset \mathbb{P}T^*A$  admit monodromy representations

$$\rho_{\gamma_{Z_i}} : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_i}^{-1}(\xi_0)),$$

then  $\mathbb{P}\Lambda_{Z_1} \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_{Z_2} \subset A \times A \times \mathbb{P}^\vee$  admits monodromy representation given by

$$(\rho_{\gamma_{Z_1}}, \rho_{\gamma_{Z_2}}) : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_1}^{-1}(\xi_0) \times \gamma_{Z_2}^{-1}(\xi_0)),$$

(2) When  $Z_1 = Z_2 = Z$ , we obtain an one-to-one correspondence:

$$\left\{ \begin{array}{c} \text{irr components of } \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \end{array} \right\}$$

*Sketch of proof.* Statement (1) holds by definition. The proof of (2) reduces to the following purely topological statement:

**Claim 4.5.** *Let  $\pi : E \rightarrow B$  be a (unramified) covering space over a manifold  $B$  with deck transformation group  $G$ , then*

$$\{ \text{connected components of } E \times_B E \} \longleftrightarrow \{ G\text{-orbits of } \pi^{-1}(b_0) \times \pi^{-1}(b_0) \}.$$

The claim follows directly from the correspondence between covering spaces over  $B$  and  $\pi_1(B)$ -sets; see [7, Theorem 1.38].  $\square$

Generalizing the argument of Lemma 4.4, we arrive at the following lemma.

**Lemma 4.6.** *For  $d = \deg \gamma_Z$ ,  $\xi_0 \in T_0^*A$  a general point, write*

$$\begin{aligned} \mathbb{P}\Lambda_Z^{\times d} &:= \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \cdots \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \subset A^d \times \mathbb{P}^\vee \\ \gamma_Z^{-1}(\xi_0)^d &:= \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \times \cdots \times \gamma_Z^{-1}(\xi_0) \subset A^d \end{aligned}$$

(1) we obtain an one-to-one correspondence:

$$\left\{ \begin{array}{c} \text{irr components of } \mathbb{P}\Lambda_Z^{\times d} \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0)^d \end{array} \right\}$$

(2) Write

$$\begin{aligned} \Delta_d &:= \{ (p_1, \dots, p_d) \in A^d \mid p_i = p_j \text{ for some } i \neq j \} \subset A^d \\ \mathbb{P}\Lambda_Z^{[d]} &:= \overline{(\mathbb{P}\Lambda_Z^{\times d} \setminus (\Delta_d \times \mathbb{P}^\vee))} \big|_U \subset A^d \times \mathbb{P}^\vee \end{aligned}$$

Fix a general point  $\xi_0 \in \mathbb{P}^\vee$  and a well-order for  $\gamma_Z^{-1}(\xi_0)$ , one can identify  $S_d \cong \gamma_Z^{-1}(\xi_0)^d \setminus \Delta_d$ , and

$$\left\{ \text{irr components of } \mathbb{P}\Lambda_Z^{[d]} \right\} \longleftrightarrow \left\{ \text{Gal}(\gamma_Z)\text{-orbits of } S_d \right\}$$

In reference,  $\Delta_d$  is usually called the big diagonal.

From this point on, we fix an irreducible component of  $\mathbb{P}\Lambda_Z^{[d]}$ , denoted by  $\mathbb{P}\Lambda_Z^{\text{univ}}$ . As we will see in Definition 4.7, this variety generates all subvarieties within the families under consideration.

**Definition 4.7** (The subvariety  $Z^{(m)}$ ). *For any tuple  $(m) = (m_1, \dots, m_d) \in \mathbb{Z}^d$ , we define the weighted sum map*

$$a^{(m)} : A^d \rightarrow A \quad (p_1, \dots, p_d) \mapsto \sum_{i=1}^d m_i p_i.$$

We also define

$$\mathbb{P}\Lambda_Z^{(m)} := \left( a^{(m)}, \text{Id}_{\mathbb{P}^\vee} \right)_* \mathbb{P}\Lambda_Z^{\text{univ}}.$$

as the (projectivized) weighted Lagrangian cycle in  $\mathbb{P}T^*A$ . The projective cycle  $\mathbb{P}\Lambda_Z^{(m)}$  is irreducible but may appear with multiplicities. We can therefore write

$$\mathbb{P}\Lambda_Z^{(m)} = c_Z^{(m)} \mathbb{P}\Lambda_{Z^{(m)}}$$

where  $c_Z^{(m)} \in \mathbb{Z}_{>0}$  and  $Z^{(m)} \subset A$  are uniquely determined. This gives rise to a family of subvarieties parametrized by  $\mathbb{Z}^d$ .

The next lemma gathers some basic properties of  $Z^{(m)}$ . Observe that  $S_d = \text{Aut}(\gamma_Z^{-1}(\xi_0))$  acts naturally on  $\mathbb{Z}^d$  via

$$g(m) = (m_{g(1)}, \dots, m_{g(d)}) \in \mathbb{Z}^d.$$

**Lemma 4.8.**

- (1) For all  $g \in \text{Gal}(\gamma_Z)$ , we have  $Z^{g(m)} = Z^{(m)}$ ,  $c_Z^{g(m)} = c_Z^{(m)}$ ;
- (2) For  $(m) = (1, 0, \dots, 0) \in \mathbb{Z}^d$ ,  $Z^{(m)} = Z$ ;
- (3) For all  $(m), (m') \in \mathbb{Z}^d$ , we have  $\mathbb{P}\Lambda_Z^{(m)} \circ \mathbb{P}\Lambda_Z^{(m')} \supseteq \mathbb{P}\Lambda_Z^{(m+m')}$ ;
- (4) The group  $\langle \mathbb{P}\Lambda_{Z^{(m)}} \rangle_{\text{Abel}}$  is closed under the convolution product.

**4.2. Realized as characteristic cycles.** In fact, the Lagrangian cycles  $\mathbb{P}\Lambda_{Z^{(m)}}$  coincide with the irreducible components of the clean cycles described in [11, 2.c] and [13, p5, Theorem 1.7], leading to the following relations:

$$\begin{array}{ccc} \text{Perv}(A)/N(A) & \supset & \langle \delta_Z \rangle \cong \text{Rep}(G_u) \\ & & \downarrow \text{cc} \quad \downarrow \\ \mathcal{L}(A) & \supset & \langle \text{cc}(\delta_Z) \rangle \stackrel{10}{\cong} \text{Rep}(T_u \rtimes \text{Gal}(\gamma_Z)) \end{array}$$

Here,  $\delta_Z$  denotes the perverse intersection complex associated with the subvariety  $Z$  (in particular,  $\delta_Z = \iota_{Z,*} \mathbb{Q}_Z[-\dim Z]$  when  $Z$  is smooth), and  $\mathcal{L}(A)$  stands for the  $\lambda$ -ring of clean conic Lagrangian cycles on  $T^*A$  [12, p5].

*Remark 4.9.* Suppose that  $Z \subset A$  is smooth of general type. Then the characteristic cycle  $\text{cc}(\delta_Z)$  is irreducible and equals  $\mathbb{P}\Lambda_Z$ . In this situation, let  $\lambda_1, \dots, \lambda_d \subset X^*(T_u)$  denote the weights corresponding to the points  $p_1, \dots, p_d \in \gamma_Z^{-1}(\xi_0)$ . For each tuple  $(m) \in \mathbb{Z}^d$ , define  $\lambda^{(m)} = \sum m_i \lambda_i \in X^*(T_u)$ , and consider the “highest weight representation”

$$V_{\lambda^{(m)}} = \bigoplus_{\lambda \in \text{Gal}(\gamma_Z) \cdot \lambda^{(m)}} \mathbb{C}_\lambda \in \text{Rep}(T_u \rtimes \text{Gal}(\gamma_Z))$$

where  $\mathbb{C}_\lambda$  is the one-dimensional representation of  $T_u$  with weight  $\lambda$ . Under this correspondence, one has an explicit identification

$$\langle \mathbb{P}\Lambda_Z \rangle \cong \text{Rep}(T_u \rtimes \text{Gal}(\gamma_Z)) \quad \mathbb{P}\Lambda_{Z^{(m)}} \longleftrightarrow V_{\lambda^{(m)}}.$$

Moreover, the Weyl group  $W_Z$  acts naturally on  $X^*(T)$ . For the orbit  $W_Z \cdot \chi_0 \subset X^*(T)$ , the associated Lagrangian cycle is given by

$$\sum_{\sigma \in W_Z / \text{Gal}(\gamma_Z)} \mathbb{P}\Lambda_{Z^{\sigma(\chi_0)}}.$$

## 5. DIMENSION AND HOMOLOGY CLASS

In this section, all cohomology groups are taken with  $\mathbb{Q}$ -coefficients for convenience in applying the Künneth formula.

**5.1. Reminder on the (homological) Chern–Mather class.** We begin by recalling the definition of the Chern–Mather class. Suppose  $\dim A = n$ , and denote by

$$p : \mathbb{P}T^*A \cong A \otimes \mathbb{P}^\vee \longrightarrow \mathbb{P}^\vee$$

the natural projection.

**Definition 5.1.** For a conic Lagrangian cycle  $\Lambda$  on  $T^*A$  and  $i \geq 0$ , the Chern–Mather class is defined as

$$c_{M,i}(\Lambda) = p_*([\mathbb{P}\Lambda] \cdot [A \times H_i]) \in H_{2i}(A) \cong H^{2(n-i)}(A),$$

where  $H_i \subseteq \mathbb{P}^\vee$  denotes a general linear subspace of dimension  $i$ . For brevity, we may later write  $c_{M,i}(\Lambda)$  as  $c_i(\Lambda)$ , and  $c_{M,i}(\Lambda_Z)$  simply as  $c_i$ .<sup>11</sup>

<sup>10</sup>I believe that this isomorphism should be already known, so I should probably cite it somewhere (rather than making it up all by myself).

<sup>11</sup>Note that although  $c_i$  arises as the pushforward of the classical Chern class in the smooth case, the indexing order is reversed. In particular,  $c_0 = d$ , rather than 1.

By the Künneth formula, we may write

$$[\mathbb{P}\Lambda] = \sum_{i=0}^{n-1} a_i \otimes H^i,$$

where  $H \in H^2(\mathbb{P}^\vee)$  denotes the hyperplane class and  $a_i \in H^{2(n-i)}(A)$ . A direct computation gives

$$\begin{aligned} c_i(\Lambda) &= p_*([\mathbb{P}\Lambda] \cdot [A \times H_i]) \\ &= p_*\left(\sum_{j=0}^{n-1} (a_j \otimes H^j) \cup (1 \otimes H^{n-1-i})\right) \\ &= p_*\left(\sum_{j=0}^{n-1} a_j \otimes H^{n-1-i+j}\right) \\ &= a_i. \end{aligned}$$

Consequently,

$$[\mathbb{P}\Lambda] = \sum_{i=0}^{n-1} c_i(\Lambda) \otimes H^i \in H^{2n}(A \times \mathbb{P}^\vee),$$

showing that the Chern–Mather classes  $c_i(\Lambda)$  are precisely the coefficients of the class  $[\mathbb{P}\Lambda]$  in the Künneth decomposition. <sup>12</sup>

*Remark 5.2.* For a subvariety  $Z \subset A$ , both its dimension  $\dim Z$  and its cohomology class  $[Z] \in H^{2(n-\dim Z)}(A)$  can be determined from  $[\mathbb{P}\Lambda] \in H^{2n}(A \times \mathbb{P}^\vee)$ , as shown in [12, Lemma 3.1.2(2)]. Indeed,

$$\begin{aligned} \dim Z &= \max \{i \in \mathbb{Z} \mid c_i \neq 0\}, \\ [Z] &= c_{\dim Z}. \end{aligned}$$

**5.2. The homology class of  $\mathbb{P}\Lambda_Z^{\times d}$ .** By Remark 5.2, it suffices to consider the homology class of  $\mathbb{P}\Lambda_{Z(m)}$ . If we are not concerned with the scalar factor  $c_Z^{(m)}$ , we may equivalently compute  $[\mathbb{P}\Lambda_Z^{(m)}]$ , which by definition is the pushforward of  $[\mathbb{P}\Lambda_Z^{\text{univ}}]$ . Consequently, the problem reduces to determining  $[\mathbb{P}\Lambda_Z^{\text{univ}}] \in H^{2dn}(A^d \times \mathbb{P}^\vee)$ .

Typically, for an initial subvariety  $Z \subset A$ , the Chern–Mather classes are known. Our ultimate goal is to express  $[\mathbb{P}\Lambda_Z^{\text{univ}}]$  in terms of these classes; as a preparatory step, we first examine  $[\mathbb{P}\Lambda_Z^{\times d}]$ .

By definition,

$$\mathbb{P}\Lambda_Z^{\times d} = \underbrace{\mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \cdots \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z}_{d \text{ factors}} = \bigcap_{i=1}^d \pi_{i, \mathbb{P}^\vee}^{-1}(\mathbb{P}\Lambda_Z),$$

---

<sup>12</sup>If the Chern–Mather classes are considered in the Chow ring, this argument does not apply, since the Künneth decomposition is unavailable at the level of Chow groups.

where  $\pi_{i, \mathbb{P}^\vee} : A^d \times \mathbb{P}^\vee \rightarrow A \times \mathbb{P}^\vee$  denotes the projection onto the  $i$ -th factor. The transversality of these intersections is immediate, so

$$\begin{aligned} [\mathbb{P}\Lambda_Z^{\times d}] &= \cup_{i=1}^d \pi_{i, \mathbb{P}^\vee}^* [\mathbb{P}\Lambda_Z] \\ &= \cup_{i=1}^d \pi_{i, \mathbb{P}^\vee}^* \left( \sum_{j=0}^{n-1} c_j \otimes H^j \right) \\ &= \cup_{i=1}^d \sum_{j=0}^{n-1} \left( 1 \otimes \cdots \otimes \underset{\substack{\uparrow \\ i\text{-th}}}{c_j} \otimes \cdots \otimes H^j \right) \\ &= \sum_{j=0}^{n-1} \left( \sum_{\sum_{k=1}^d j_k = j} c_{j_1} \otimes \cdots \otimes c_{j_d} \right) \otimes H^j. \end{aligned}$$

**5.3. The homology class of  $\mathbb{P}\Lambda_Z^{[d]}$ .** This subsection explains how to eliminate the contribution of the big diagonal and how to compute  $\mathbb{P}\Lambda_Z^{[d]}$  from  $\mathbb{P}\Lambda_Z^{\times d}$ . For this purpose, we introduce some combinatorial preliminaries.

**Definition 5.3.** For  $n \in \mathbb{N}_{>0}$ , let  $[n] := \{1, \dots, n\}$ , and denote by  $\mathcal{P}(n)$  the lattice of partitions of  $[n]$ , ordered by refinement:  $\alpha' \leq \alpha$  if and only if any two elements  $i, j$  belonging to the same block of  $\alpha'$  also belong to the same block of  $\alpha$ . For a partition  $\alpha = \{A_1, \dots, A_k\} \in \mathcal{P}(d)$ , we associate a surjective map

$$f_\alpha : [d] \longrightarrow [k] \quad a \longmapsto j \quad \text{if } a \in A_j$$

which is well-defined up to the natural  $S_k$ -action; this indeterminacy will not affect our discussion. Each map  $f_\alpha$  naturally gives rise to a partial diagonal embedding

$$\Delta_\alpha : A^k \times \mathbb{P}^\vee \longrightarrow A^d \times \mathbb{P}^\vee \quad ((p_i), \xi) \longmapsto ((p_{f_\alpha(i)}), \xi).$$

This construction determines a subvariety of  $\mathbb{P}\Lambda_Z^{\times d}$ , defined by

$$\begin{aligned} \mathbb{P}\Lambda_Z^{\geq \alpha} &:= \Delta_\alpha (\mathbb{P}\Lambda_Z^{\times k}) \\ &= \{((p_i), \xi) \in \mathbb{P}\Lambda_Z^{\times d} \mid p_i = p_j \text{ if } i \sim_\alpha j\}. \end{aligned}$$

We can in addition define the locus corresponding precisely to  $\alpha$ :

$$\mathbb{P}\Lambda_Z^\alpha := \overline{\{((p_i), \xi) \in \mathbb{P}\Lambda_Z^{\times d} \mid p_i = p_j \text{ iff } i \sim_\alpha j\}}.$$

*Remark 5.4.* Denote by

$$\hat{0} := \{\{1\}, \dots, \{d\}\} \in \mathcal{P}(d)$$

the finest partition. Then  $\mathbb{P}\Lambda_Z^{\times d} = \mathbb{P}\Lambda_Z^{\geq \hat{0}}$ , and  $\mathbb{P}\Lambda_Z^{[d]} = \mathbb{P}\Lambda_Z^{\hat{0}}$ .

By definition,

$$[\mathbb{P}\Lambda_Z^{\geq \alpha}] = \sum_{\alpha' \geq \alpha} [\mathbb{P}\Lambda_Z^{\alpha'}].$$

Applying Möbius inversion on the partition lattice yields

$$[\mathbb{P}\Lambda_Z^\alpha] = \sum_{\alpha' \geq \alpha} \mu(\alpha, \alpha') [\mathbb{P}\Lambda_Z^{\geq \alpha'}],$$

where  $\mu(\alpha, \alpha')$  denotes the Möbius function as defined in [1, p141].

**Fact 5.5** (See [1, IV.3]). Let  $\alpha' \geq \alpha$  be two partitions of  $[d]$ , where  $\alpha' = \{A_1, \dots, A_k\}$ . For each  $i$ , denote by  $r_i$  the number of blocks of  $\alpha$  that are contained in  $A_i$ . Then

$$\mu(\alpha, \alpha') = (-1)^{|\alpha| - k} \prod_{i=1}^k (r_i - 1)!$$

In particular,

$$\mu(\hat{0}, \alpha') = (-1)^{d-k} \prod_{i=1}^k (|A_i| - 1)!$$

With Fact 5.5, one can compute  $[\mathbb{P}\Lambda_Z^{[d]}] = [\mathbb{P}\Lambda_Z^{\hat{0}}]$  by the Möbius inverse formula:

$$\begin{aligned} [\mathbb{P}\Lambda_Z^{[d]}] &= \sum_{\alpha'} \mu(\hat{0}, \alpha') [\mathbb{P}\Lambda_Z^{\geq \alpha'}] \\ &= \sum_{k=1}^d \sum_{\alpha' = \{A_1, \dots, A_k\}} (-1)^{d-k} \prod_{i=1}^k (|A_i| - 1)! \cdot \Delta_{\alpha, *} [\mathbb{P}\Lambda_Z^{\times k}] \end{aligned}$$

The computation of pushforwards along diagonal embeddings can be rather cumbersome. However, upon composing with  $a^{(m)}$ , it suffices to compute those of certain weighted sum maps. These pushforwards admit a more transparent description via the Pontryagin product, whose definition we now recall.

**Definition 5.6** (Pontryagin product). *Let  $A$  be an abelian variety, and denote by  $a : A \times A \rightarrow A$  the addition map. The Pontryagin product on  $A$  is defined by*

$$H^{2n-i}(A) \times H^{2n-j}(A) \subseteq H^{4n-i-j}(A \times A) \xrightarrow{a_*} H^{2n-i-j}(A) \quad a \otimes b \mapsto a * b.$$

*Remark 5.7.* The Pontryagin product is unital and associative, but in general only anti-commutative [4, 1.5.(7) b)]. In the context of our work, we are concerned with Chern–Mather classes, which have even degrees; consequently, the anti-commutativity of the Pontryagin product does not pose any complications.

In particular, for any sum map  $a : A^k \rightarrow A$ , one can unambiguously write

$$a_*(a_1 \otimes \cdots \otimes a_k) = a_1 * \cdots * a_k \triangleq \bigstar_{i=1}^k a_i$$

where no additional parentheses are required.

**Lemma 5.8.** *For any tuple  $(m) = (m_1, \dots, m_k) \in \mathbb{Z}^k$ ,  $a_i \in H^{2(n-l_i)}(A)$ ,*

$$a_*^{(m)}(a_1 \otimes \cdots \otimes a_k) = \left( \prod_{i=1}^k m_i^{2l_i} \right) \cdot \bigstar_{i=1}^k a_i.$$

*Proof.* Notice that  $a^{(m)}$  can be written as compositions of basic functions:

$$a^{(m)} : A^k \xrightarrow{(m_1, \dots, m_k)} A^k \xrightarrow{a} A$$

□

**Definition 5.9.** *For any tuple  $(m) = (m_1, \dots, m_d) \in \mathbb{Z}^d$  and any partition  $\alpha = \{A_1, \dots, A_k\} \in \mathcal{P}(d)$ , we define*

$$\alpha(m) := \left( \sum_{i \in A_1} m_i, \dots, \sum_{i \in A_k} m_i \right) \in \mathbb{Z}^k.$$

Moreover, for  $\mathbf{l} = (l_1, \dots, l_k) \in \mathbb{N}_{\geq 0}^k$  with  $\sum l_i = l$ , we set

$$\alpha(m)^{2\mathbf{l}} := \left( \sum_{i \in A_1} m_i \right)^{2l_1} \cdots \left( \sum_{i \in A_k} m_i \right)^{2l_k}$$

which defines a homogeneous polynomial of degree  $2l$  in  $\mathbb{Z}[m_1, \dots, m_k]$ .

We can now determine  $[\mathbb{P}\Lambda_Z^{(m)}]$  in the case where the monodromy group is  $S_d$ :

$$\begin{aligned}
[\mathbb{P}\Lambda_Z^{(m)}] &= \left(a^{(m)}, \text{Id}_{\mathbb{P}^\vee}\right)_* [\mathbb{P}\Lambda_Z^{\text{univ}}] \\
&= \left(a^{(m)}, \text{Id}_{\mathbb{P}^\vee}\right)_* [\mathbb{P}\Lambda_Z^{[d]}] \\
&= \sum_{\alpha} \mu(\hat{0}, \alpha) \left(a^{(m)}, \text{Id}_{\mathbb{P}^\vee}\right)_* \Delta_{\alpha,*} [\mathbb{P}\Lambda_Z^{\times k}] \\
&= \sum_{\alpha} \mu(\hat{0}, \alpha) \left(a^{\alpha(m)}, \text{Id}_{\mathbb{P}^\vee}\right)_* [\mathbb{P}\Lambda_Z^{\times k}] \\
&= \sum_{\alpha} \mu(\hat{0}, \alpha) \left(a^{\alpha(m)}, \text{Id}_{\mathbb{P}^\vee}\right)_* \left( \sum_{l=0}^{n-1} \left( \sum_{\sum l_i=l} c_{l_1} \otimes \cdots \otimes c_{l_k} \right) \otimes H^l \right) \\
&= \sum_{\alpha} \mu(\hat{0}, \alpha) \sum_{l=0}^{n-1} \sum_{\sum l_i=l} \alpha(m)^{2l} \left( \bigstar_{i=1}^k c_{l_i} \otimes H^l \right) \\
&= \sum_{l=0}^{n-1} \left( \sum_{\alpha} \sum_{\sum l_i=l} \left( \mu(\hat{0}, \alpha) \alpha(m)^{2l} \bigstar_{i=1}^k c_{l_i} \right) \right) \otimes H^l
\end{aligned}$$

Therefore,

$$c_l(\Lambda_{Z^{(m)}}) = \frac{1}{c_Z^{(m)}} \sum_{\alpha} \sum_{\sum l_i=l} \left( \mu(\hat{0}, \alpha) \alpha(m)^{2l} \bigstar_{i=1}^k c_{l_i} \right) \quad (5.1)$$

The following corollary collects some quantitative implications of (5.1).

**Corollary 5.10.** *Assume that  $\text{Gal}(\gamma_Z) = S_d$ .*

- (1) The Chern–Mather classes

$$c_l(\Lambda_{Z^{(m)}}) \in H^{2(n-l)}(A)[m_1, \dots, m_d]^{S_d}$$

are polynomials of degree  $2l$  with exponent at most  $2r$ , when expressed in the variables  $m_1, \dots, m_d$ .

- (2) If  $Z \subset A$  is a smooth curve, there exists a homogeneous symmetric polynomial  $f_{Z,l} \in \mathbb{Q}[m_1, \dots, m_d]^{S_d}$  of degree  $2l$  and exponent at most 2, such that

$$c_l(\Lambda_{Z^{(m)}}) = f_{Z,l}(m_1, \dots, m_d) \left( \bigstar_{i=1}^k c_{l_i} \right).$$

In this case,

$$Z^{(m)} \subset A \text{ is a divisor} \iff f_{Z,n-1}(m_1, \dots, m_d) \neq 0.$$

**5.4. The homology class in type C case.** In many instances, the subvariety  $Z \subset A$  is invariant under a reflection, and we may, without loss of generality, assume that the reflection is taken with respect to the origin, so that  $Z = -Z$ . In this situation, the degree  $d = \deg \gamma_Z$  is necessarily even, and the locus  $[\mathbb{P}\Lambda_Z^{[d]}]$  admits a decomposition into a finite union of subvarieties. To describe these components, let

$$\mathcal{P}_2(d) := \{ \alpha \in \mathcal{P}(d) \mid |A| = 2 \text{ for all } A \in \alpha \}$$

denote the set of all perfect matchings of  $[d]$ . Each  $\alpha \in \mathcal{P}_2(d)$  induces an involution  $\tau_\alpha$  of  $[d]$ , and we define

$$\begin{aligned}
\mathbb{P}\Lambda_Z^{[\widetilde{\alpha}]} &:= \left\{ ((p_i), \xi) \in \mathbb{P}\Lambda_Z^{[d]} \mid p_{\tau_\alpha(i)} = -p_i \text{ for all } i \right\} \\
\Delta_{\widetilde{\alpha}} : A^{d/2} \times \mathbb{P}^\vee &\longrightarrow A^d \times \mathbb{P}^\vee \quad ((p_i), \xi) \longmapsto ((\pm p_{f_\alpha(i)}), \xi)
\end{aligned}$$

where the sign is chosen so that  $p_{\tau_\alpha(i)} = -p_i$ . Furthermore, there exists a distinguished subvariety in  $\mathbb{P}\Lambda_Z^{\times d/2}$ :

$$\mathbb{P}\Lambda_Z^{\widetilde{[d/2]}} := \overline{\left\{ ((p_i), \xi) \in \mathbb{P}\Lambda_Z^{\times d/2} \mid p_i \neq \pm p_j \text{ iff } i \neq j \right\}}.$$

With these conventions, we can now express the following decomposition:

$$\begin{aligned} \mathbb{P}\Lambda_Z^{[d]} &= \bigcup_{\alpha \in \mathcal{P}_2(d)} \mathbb{P}\Lambda_Z^{[\alpha]} \\ &= \bigcup_{\alpha \in \mathcal{P}_2(d)} \Delta_{\tilde{\alpha}} \left( \mathbb{P}\Lambda_Z^{\widetilde{[d/2]}} \right) \end{aligned}$$

Evidently, determining  $[\mathbb{P}\Lambda_Z^{[\alpha]}]$  amounts to determining  $[\mathbb{P}\Lambda_Z^{\widetilde{[d/2]}}]$ , and the latter computation reduces again to combinatorics.

For  $\beta \in \mathcal{P}(d/2)$ , define

$$\begin{aligned} \mathbb{P}\Lambda_Z^{\widetilde{\geq \beta}} &:= \left\{ ((p_i), \xi) \in \mathbb{P}\Lambda_Z^{\times d/2} \mid p_i = \pm p_j \text{ if } i \sim_\beta j \right\} \\ \mathbb{P}\Lambda_Z^{\tilde{\beta}} &:= \overline{\left\{ ((p_i), \xi) \in \mathbb{P}\Lambda_Z^{\times d/2} \mid p_i = \pm p_j \text{ iff } i \sim_\beta j \right\}} \end{aligned}$$

Then one has the relations

$$\begin{aligned} [\mathbb{P}\Lambda_Z^{\widetilde{\geq \beta}}] &= \sum_{\beta' \geq \beta} [\mathbb{P}\Lambda_Z^{\tilde{\beta}'}], \\ [\mathbb{P}\Lambda_Z^{\tilde{\beta}}] &= \sum_{\beta' \geq \beta} \mu(\beta, \beta') [\mathbb{P}\Lambda_Z^{\widetilde{\geq \beta'}}], \end{aligned}$$

where  $\mu(\beta, \beta')$  denotes the Möbius function of the poset  $\mathcal{P}(d/2)$ .

Furthermore, the classes  $[\mathbb{P}\Lambda_Z^{\widetilde{\geq \beta}}]$  admit a decomposition as sums of pushforwards of  $[\mathbb{P}\Lambda_Z^{\times k}]$  under appropriately defined signed diagonal maps. Explicitly, for every pair of maps  $f_\beta : [d/2] \rightarrow [k]$  and  $\eta : [d/2] \rightarrow \{\pm 1\}$ , one introduces the signed diagonal embedding

$$\Delta_\beta^\eta : A^k \times \mathbb{P}^\vee \longrightarrow A^{d/2} \times \mathbb{P}^\vee \quad ((p_i), \xi) \longmapsto ((\eta(i)p_{f_\beta(i)}), \xi),$$

which yields the expression

$$[\mathbb{P}\Lambda_Z^{\widetilde{\geq \beta}}] = \frac{1}{2^k} \sum_{\eta : [d/2] \rightarrow \{\pm 1\}} \Delta_{\beta,*}^\eta [\mathbb{P}\Lambda_Z^{\times k}].$$

Assuming that the monodromy group is the signed symmetric group, we may identify  $\mathbb{P}\Lambda_Z^{\text{univ}}$  with one of the components  $\mathbb{P}\Lambda_Z^{[\alpha]}$ . Without loss of generality, we choose

$$\alpha = \{ \{i, i + d/2\} \mid i = 1, \dots, d/2 \}$$

and fix the sign convention for  $\Delta_{\tilde{\alpha}}$ :

$$\Delta_{\tilde{\alpha}} : A^{d/2} \times \mathbb{P}^\vee \xrightarrow{(\Delta, \text{Id})} A^{d/2} \times A^{d/2} \times \mathbb{P}^\vee \xrightarrow{(\text{Id}, -\text{Id}, \text{Id})} A^d \times \mathbb{P}^\vee$$

Under this convention, the tuples  $(m_1, \dots, m_d)$  and  $(m_1 - m_{d/2+1}, \dots, m_{d/2} - m_d, 0, \dots, 0)$  define the same subvariety. Hence, without loss of generality, we may assume that

$$(m) := (m_1, \dots, m_{d/2}, 0, \dots, 0) \in \mathbb{Z}^d.$$

We now introduce some notations for later computations:

$$\begin{aligned} (\tilde{m}) &:= (m_1, \dots, m_{d/2}) \in \mathbb{Z}^{d/2} \\ \beta^\eta(\tilde{m}) &:= \left( \sum_{i \in A_1} \eta(i)m_i, \dots, \sum_{i \in A_k} \eta(i)m_i \right) \in \mathbb{Z}^k \\ \beta^\eta(\tilde{m})^{2\mathbf{l}} &:= \left( \sum_{i \in A_1} \eta(i)m_i \right)^{2l_1} \cdots \left( \sum_{i \in A_k} \eta(i)m_i \right)^{2l_k} \\ \beta(\tilde{m}^2)^{\mathbf{l}} &:= \left( \sum_{i \in A_1} m_i^2 \right)^{l_1} \cdots \left( \sum_{i \in A_k} m_i^2 \right)^{l_k} \end{aligned}$$

where  $\beta = \{A_1, \dots, A_k\} \in \mathcal{P}(d/2)$  is a partition of length  $k$ ,  $\eta : [d/2] \rightarrow \{\pm 1\}$  is a sign function, and  $\mathbf{l} = (l_1, \dots, l_k) \in \mathbb{N}_{\geq 0}^k$  with  $\sum l_i = l$ . Notice that

$$\sum_{\eta} \beta^\eta(\tilde{m})^{2\mathbf{l}} = 2^{d/2} \beta(\tilde{m}^2)^{\mathbf{l}}.$$

We can now determine  $[\mathbb{P}\Lambda_Z^{(m)}]$  in the case where  $Z = -Z$  and  $\text{Gal}(\gamma_Z) \cong W(C_{d/2})$ :

$$\begin{aligned} [\mathbb{P}\Lambda_Z^{(m)}] &= \left( a^{(m)}, \text{Id}_{\mathbb{P}^V} \right)_* [\mathbb{P}\Lambda_Z^{[\alpha]}] \\ &= \left( a^{(\tilde{m})}, \text{Id}_{\mathbb{P}^V} \right)_* [\mathbb{P}\Lambda_Z^{[d/2]}] \\ &= \left( a^{(\tilde{m})}, \text{Id}_{\mathbb{P}^V} \right)_* [\mathbb{P}\Lambda_Z^{\tilde{0}}] \\ &= \sum_{\beta} \mu(\hat{0}, \beta) \left( a^{(\tilde{m})}, \text{Id}_{\mathbb{P}^V} \right)_* [\mathbb{P}\Lambda_Z^{\geq \beta}] \\ &= \sum_{\beta, \eta} \frac{1}{2^k} \mu(\hat{0}, \beta) \left( a^{\beta^\eta(\tilde{m})}, \text{Id}_{\mathbb{P}^V} \right)_* [\mathbb{P}\Lambda_Z^{\times k}] \\ &= \sum_{\beta, \eta} \frac{1}{2^k} \mu(\hat{0}, \beta) \left( a^{\beta^\eta(\tilde{m})}, \text{Id}_{\mathbb{P}^V} \right)_* \left( \sum_{l=0}^{n-1} \left( \sum_{\sum l_i=l} c_{l_1} \otimes \cdots \otimes c_{l_k} \right) \otimes H^l \right) \\ &= \sum_{\beta, \eta} \frac{1}{2^k} \mu(\hat{0}, \beta) \sum_{l=0}^{n-1} \sum_{\sum l_i=l} \beta^\eta(\tilde{m})^{2\mathbf{l}} \left( \bigstar_{i=1}^k c_{l_i} \otimes H^l \right) \\ &= \sum_{l=0}^{n-1} \left( \sum_{\beta, \eta} \sum_{\sum l_i=l} \left( \frac{1}{2^k} \mu(\hat{0}, \beta) \beta^\eta(\tilde{m})^{2\mathbf{l}} \bigstar_{i=1}^k c_{l_i} \right) \right) \otimes H^l \\ &= \sum_{l=0}^{n-1} \left( \sum_{\beta} \sum_{\sum l_i=l} \left( 2^{d/2-k} \mu(\hat{0}, \beta) \beta(\tilde{m}^2)^{\mathbf{l}} \bigstar_{i=1}^k c_{l_i} \right) \right) \otimes H^l \end{aligned}$$

Therefore,

$$c_l(\Lambda_{Z^{(m)}}) = \frac{1}{c_Z^{(m)}} \sum_{\beta} \sum_{\sum l_i=l} \left( 2^{d/2-k} \mu(\hat{0}, \beta) \beta(\tilde{m}^2)^{\mathbf{l}} \bigstar_{i=1}^k c_{l_i} \right) \quad (5.2)$$

The following corollary collects some quantitative implications of (5.2).

**Corollary 5.11.** *Suppose that  $Z = -Z$  and  $\text{Gal}(\gamma_Z) \cong W(C_{d/2})$ , and assume that  $(m) := (m_1, \dots, m_{d/2}, 0, \dots, 0) \in \mathbb{Z}^d$ .*

(1) The Chern–Mather classes

$$c_l(\Lambda_{Z^{(m)}}) \in H^{2(n-l)}(A)[m_1^2, \dots, m_{d/2}^2]^{S_{d/2}}$$

are polynomials of degree  $l$  with exponent at most  $2r$ , when expressed in the variables  $m_1^2, \dots, m_{d/2}^2$ .

- (2) If  $Z \subset A$  is a smooth curve, there exists a homogeneous symmetric polynomial  $\tilde{f}_{Z,l} \in \mathbb{Q}[m_1^2, \dots, m_{d/2}^2]^{S_{d/2}}$  of degree  $l$  and exponent at most 2, such that

$$c_l(\Lambda_{Z^{(m)}}) = \tilde{f}_{Z,l}(m_1^2, \dots, m_{d/2}^2) \left( \begin{smallmatrix} k \\ * \\ c_1 \end{smallmatrix} \right).$$

In this case,

$$Z^{(m)} \subset A \text{ is a divisor} \iff \tilde{f}_{Z,n-1}(m_1^2, \dots, m_{d/2}^2) \neq 0.$$

**5.5. Möbius-transforms block-sum polynomial.** The formulas (5.1) and (5.2) admit further simplifications in certain special situations, such as the case of curves. In what follows, we extract the corresponding polynomial coefficient, which may be of independent combinatorial interest.

**Definition 5.12.** Fix  $d$  and  $\tilde{d} = d/2$ , and let  $l \in \mathbb{N}_{\geq 0}$ . For a partition  $\lambda = [\lambda_1, \dots, \lambda_{k'}]$  of  $l$  with  $\lambda_1 \geq \dots \geq \lambda_{k'} > 0$ , the Möbius-transforms block-sum polynomial of type  $A$  (resp. type  $C$ ) is defined by

$$\begin{aligned} \mu_d^\lambda &:= \sum_{\alpha \in \mathcal{P}(d)} \sum_{\substack{\mathbf{l} \vdash \lambda \\ \mathbf{l}: \text{length } k}} \mu(\hat{0}, \alpha) \alpha(m)^{2\mathbf{l}} d^{k-k'} \in \mathbb{Z}[m_1, \dots, m_d]^{S_d} \\ \tilde{\mu}_{\tilde{d}}^\lambda &:= \sum_{\beta \in \mathcal{P}(\tilde{d})} \sum_{\substack{\mathbf{l} \vdash \lambda \\ \mathbf{l}: \text{length } k}} 2^{\tilde{d}-k} \mu(\hat{0}, \beta) \beta(\tilde{m}^2)^{\mathbf{l}} (2\tilde{d})^{k-k'} \in \mathbb{Z}[m_1^2, \dots, m_{\tilde{d}}^2]^{S_{\tilde{d}}} \end{aligned}$$

In this definition:

- $\alpha = \{A_1, \dots, A_k\} \in \mathcal{P}(d)$  is a set partition of  $[d]$  with  $k = \#\alpha$  blocks (analogously for  $\beta$ );
- $\mathbf{l} = (l_1, \dots, l_k) \in \mathbb{Z}^k$ , and  $\mathbf{l} \vdash \lambda$  indicates that  $\mathbf{l}$  is of type  $\lambda$ , meaning that its nonzero components, counted with multiplicities, form the partition  $\lambda$ ;
- $\mu(\hat{0}, \alpha) = (-1)^{d-k} \prod_{i=1}^k (|A_i| - 1)!$  is the Möbius function;
- $\alpha(m)^{2\mathbf{l}} = (\sum_{i \in A_1} m_i)^{2l_1} \cdots (\sum_{i \in A_k} m_i)^{2l_k}$  and  $\beta(\tilde{m}^2)^{\mathbf{l}} = (\sum_{i \in A_1} m_i^2)^{l_1} \cdots (\sum_{i \in A_k} m_i^2)^{l_k}$ .

Now, the formulas (5.1) and (5.2) can be written as

$$\begin{aligned} c_l(\Lambda_{Z^{(m)}}) &= \frac{1}{c_Z^{(m)}} \sum_{\lambda \vdash l} \mu_d^\lambda \left( \begin{smallmatrix} k' \\ * \\ c_{\lambda_i} \end{smallmatrix} \right) \quad \text{type } A \text{ case} \\ c_l(\Lambda_{Z^{(m)}}) &= \frac{1}{c_Z^{(m)}} \sum_{\lambda \vdash l} \tilde{\mu}_{\tilde{d}}^\lambda \left( \begin{smallmatrix} k' \\ * \\ c_{\lambda_i} \end{smallmatrix} \right) \quad \text{type } C \text{ case} \end{aligned}$$

The following identities have been verified in low-degree cases using SageMath, though a purely combinatorial explanation, as well as a general formulation for the remaining types, is still lacking.

**Fact 5.13.** Let  $1^{k'} := [1, \dots, 1] \vdash k'$ . When  $k' < \tilde{d}$ , we have

$$\begin{aligned} \mu_d^{1^{k'}} &= \frac{1}{2^{k'} k'!} \sum_{\sigma \in S_d} \prod_{i=1}^{k'} (m_{\sigma(2i-1)} - m_{\sigma(2i)})^2 \\ \tilde{\mu}_{\tilde{d}}^{1^{k'}} &= 2^{\tilde{d}} \frac{1}{2^{k'} k'!} \sum_{\sigma \in S_d} \prod_{i=1}^{k'} m_{\sigma(i)}^2 \\ &= 2^{\tilde{d}-k'} (\tilde{d} - k')! \sum_{1 \leq i_1 < \dots < i_{k'} \leq \tilde{d}} m_{i_1}^2 \cdots m_{i_{k'}}^2 \end{aligned}$$

**Example 5.14** (Continuation of Example 3.7). *Let  $C$  be a smooth curve of genus  $g$  embedded in its Jacobian  $A := \text{Jac}(C)$  via the Abel–Jacobi map  $\text{AJ}_C : C \hookrightarrow A$ .*

*When  $C$  is non-hyperelliptic, we get*

$$\dim Z^{(m)} = \min_{k \in \mathbb{Z}} \{g - 1, \# \{i \in [2g - 2] \mid m_i \neq k\}\}.$$

*When  $C$  is hyperelliptic, with  $(m) = (m_1, \dots, m_{g-1}, 0, \dots, 0)$ , we get*

$$\dim Z^{(m)} = \# \{i \in [g - 1] \mid m_i \neq 0\}.$$

#### REFERENCES

- [1] Martin Aigner. *Combinatorial theory*, volume 234 of *Grundlehren Math. Wiss.* Springer, Cham, 1979.
- [2] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Volume I*, volume 267 of *Grundlehren Math. Wiss.* Springer, Cham, 1985.
- [3] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergeb. Math. Grenzgeb., 3. Folge*. Berlin: Springer, 2nd enlarged ed. edition, 2004.
- [4] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 302 of *Grundlehren Math. Wiss.* Berlin: Springer, 2nd augmented ed. edition, 2004.
- [5] Ciro Ciliberto, Thomas Dedieu, Concettina Galati, and Andreas Leopold Knutsen. On the locus of Prym curves where the Prym-canonical map is not an embedding. *Ark. Mat.*, 58(1):71–85, 2020.
- [6] Giulio Codogni and Thomas Krämer. Semicontinuity of Gauss maps and the Schottky problem. *Math. Ann.*, 382(1-2):607–630, 2022.
- [7] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge University Press, 2002.
- [8] Ariyan Javanpeykar, Thomas Krämer, Christian Lehn, and Marco Maculan. The monodromy of families of subvarieties on abelian varieties. *Duke Math. J.*, 174(6):1045–1149, 2025.
- [9] Ernst Kani. On Castelnuovo’s equivalence defect. *J. Reine Angew. Math.*, 352:24–70, 1984.
- [10] Thomas Krämer. Cubic threefolds, Fano surfaces and the monodromy of the Gauss map. *Manuscr. Math.*, 149(3-4):303–314, 2016.
- [11] Thomas Krämer. Summands of theta divisors on Jacobians. *Compos. Math.*, 156(7):1457–1475, 2020.
- [12] Thomas Krämer. Characteristic cycles and the microlocal geometry of the Gauss map. II. *J. Reine Angew. Math.*, 774:53–92, 2021.
- [13] Thomas Krämer. Characteristic cycles and the microlocal geometry of the Gauss map. I. *Ann. Sci. Éc. Norm. Supér. (4)*, 55(6):1475–1527, 2022.
- [14] Herbert Lange and Ángela Ortega. Prym varieties of cyclic coverings. *Geom. Dedicata*, 150:391–403, 2011.
- [15] John W. Milnor. *Topology from the differentiable viewpoint*. Charlottesville: The University Press of Virginia. IX, 64 p. (1965)., 1965.
- [16] Ravi Vakil. *The rising sea: Foundations of algebraic geometry. preprint*, 2017.

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, BERLIN, 12489, GERMANY,  
*Email address:* email:xiaoxiang.zhou@hu-berlin.de