

SUBVARIETIES IN COMPLEX ABELIAN VARIETIES

XIAOXIANG ZHOU

CONTENTS

1. Introduction	1
Acknowledgement	2
2. Tangent Gauss map and Conormal Gauss map	2
3. Families of subvarieties	7
4. When is monodromy group small	9
5. Tannakian formalism	17
6. delete later	18
References	18

1. INTRODUCTION

From any subvariety of an abelian variety one obtains a reductive group via the convolution of perverse sheaves; in this way, traditional questions in algebraic geometry can be reinterpreted in representation-theoretic terms, providing new geometric perspectives.

In [10], Prof. Krämer shows that Weyl group orbits correspond to conic Lagrangian cycles in the cotangent bundle, which are realized as the conormal bundles of certain subvarieties. A careful analysis of these subvarieties is frequently central to resolving the problems at hand, cf. [7, §5-§8]. Here we investigate three principal properties of these subvarieties: irreducibility, dimension, and homology class.

Concerning irreducibility, we observe that the irreducible components correspond precisely to the orbits of the monodromy group. Hence, the discrepancy between the monodromy group and the Weyl group provides a measure of the failure of irreducibility of these subvarieties. Recently, we constructed an example where the monodromy group is significantly smaller than expected, thereby disproving [9, Conjecture 8]:

Proposition 1.1. *There exists an étale double cover $h : C \rightarrow C'$ of smooth projective curves such that the Abel–Prym map*

$$\mathrm{AP}_{C/C'} : C \longrightarrow A \quad p \longmapsto \mathcal{O}_C(p - \iota(p))$$

is an embedding, and

$$W_C \cong S_2^{\oplus d/2} \rtimes S_{d/2}$$

$$M_C \subseteq S_4^{\oplus d/4} \rtimes S_{d/4}$$

where $d = 2g(C') - 2$, and W_C, M_C denote the Weyl and monodromy groups, respectively.

This turns out to be the only nontrivial example in the Prym setting when $g(C') > 9$.

For the dimension and homology class of a subvariety Z , both invariants can be recovered from the homology class of $\mathbb{P}\Lambda_Z \subset \mathbb{P}T^*A$ — that is, from the Chern–Mather class of Z . Standard computational techniques are available when the group is a full or signed symmetric group. This applies in particular to the Jacobian case, which will be analyzed in detail in Section ???.

ACKNOWLEDGEMENT

I would like to express my sincere gratitude to my supervisor, Thomas Krämer, without whose support this work would not have been possible. He proposed the problem, guided me throughout the project, and generously spent many hours discussing drafts and explaining important ideas. I am also thankful to Haohao Liu, Vincent Ariksoy, Aaron Beumer, Roben de Preter, Andreas Kretschmer, Andrés Rojas, Søren Gammelgaard, Paweł Borówka, and Gavril Farkas for their insightful discussions and valuable comments. Funding is provided by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project ID: 390685689).

2. TANGENT GAUSS MAP AND CONORMAL GAUSS MAP

For simplicity, we work over the base field $\kappa = \mathbb{C}$, and by a variety we mean a integral separated scheme of finite type over \mathbb{C} . Let A/\mathbb{C} be an abelian variety of dimension n , and let $Z \subseteq A$ be an irreducible closed subvariety of dimension r . We denote by $\iota_Z : Z \hookrightarrow A$ the inclusion morphism.¹

2.1. Gauss map and monodromy group.

Definition 2.1. For a subvariety $Z \subset A$, the tangent Gauss map of Z is defined as

$$\phi_Z : Z^{\text{sm}} \longrightarrow \text{Gr}(r, T_0 A) \quad p \longmapsto T_p Z \subseteq T_p A \cong T_0 A$$

which describes the tangent space information at each point.

Remark 2.2. Any map $X \longrightarrow \text{Gr}(r, V)$ is induced by a rank r vector bundle \mathcal{E} on X together with an epimorphism $V^* \otimes \mathcal{O}_X \twoheadrightarrow \mathcal{E}$. In our case, the map ϕ_Z is induced by the tangent bundle $\mathcal{T}_{Z^{\text{sm}}}$ and the sections

$$T_0^* A \otimes_{\mathbb{C}} \mathcal{O}_{Z^{\text{sm}}} \subseteq H^0(Z^{\text{sm}}, \mathcal{T}_{Z^{\text{sm}}}^\vee) \otimes_{\mathbb{C}} \mathcal{O}_{Z^{\text{sm}}} \twoheadrightarrow \mathcal{T}_{Z^{\text{sm}}}^\vee.$$

The definition of the conormal Gauss map requires a brief recollection of the conormal variety. On the smooth locus, the normal and conormal bundles behave well as vector bundles:²

$$\mathcal{N}_{Z^{\text{sm}}/A} := \mathcal{T}_A|_{Z^{\text{sm}}} / \mathcal{T}_{Z^{\text{sm}}} \quad \mathcal{N}_{Z^{\text{sm}}/A}^* = \ker \left(\Omega_A|_{Z^{\text{sm}}} \rightarrow \Omega_{Z^{\text{sm}}} \right).$$

We write $\Lambda_{Z^{\text{sm}}}$ as the total space of $\mathcal{N}_{Z^{\text{sm}}/A}^*$. The conormal variety Λ_Z is the closure of $\Lambda_{Z^{\text{sm}}}$ viewed as a subvariety in T^*A :

$$\Lambda_Z := \overline{\Lambda_{Z^{\text{sm}}}} \subset T^*A \cong A \times T_0^*A$$

this is a conical Lagrangian cycle in T^*A . The (affine) conormal Gauss map is defined as

$$\gamma_Z^{\text{aff}} : \Lambda_Z \subset A \times T_0^*A \longrightarrow T_0^*A$$

For intersection-theoretic purposes it is more natural to work with projectivized conormal spaces. We therefore pass from the affine conormal Gauss map to its projectivized version:

Definition 2.3. the projectivized conormal variety

$$\mathbb{P}\Lambda_Z := \overline{\mathbb{P}\Lambda_{Z^{\text{sm}}}} \subset \mathbb{P}T^*A \cong A \times \mathbb{P}T_0^*A$$

is a Legendrian cycle in the contact variety $A \times \mathbb{P}T_0^*A$. $\mathbb{P}\Lambda_{Z^{\text{sm}}}$ is a \mathbb{P}^{r-1} -bundle over Z^{sm} , and the map

$$\gamma_Z : \mathbb{P}\Lambda_Z \subset A \times \mathbb{P}T_0^*A \longrightarrow \mathbb{P}T_0^*A$$

¹I'm not sure whether we should consider the more general cases in the future—such as working over a field of characteristic p , letting A be a semiabelian variety or a complex torus, or allowing ι to be a covering onto its image. For now, I will omit these possibilities from this document.

²This is more symmetric when writing them as short exact sequences:

$$\begin{aligned} 0 &\longrightarrow \mathcal{T}_{Z^{\text{sm}}} \longrightarrow \mathcal{T}_A|_{Z^{\text{sm}}} \longrightarrow \mathcal{N}_{Z^{\text{sm}}/A} \longrightarrow 0 \\ 0 &\longrightarrow \mathcal{N}_{Z^{\text{sm}}/A}^* \longrightarrow \Omega_A|_{Z^{\text{sm}}} \longrightarrow \Omega_{Z^{\text{sm}}} \longrightarrow 0 \end{aligned}$$

is called the (projectivized) conormal Gauss map.

By [7, Theorem 2.8 (1)], γ_Z is generically finite when Z is (an integral variety) of general type. A lot of geometry of Z is encoded in the map γ_Z . For instance, if $Z \subset A$ is smooth, then

$$\deg \gamma_Z = (-1)^r \chi(Z),$$

where $\chi(Z) = \sum_i (-1)^i b_i(Z)$ is the topological Euler characteristic of Z .

Further insight can be gained by analyzing the fibers of γ_Z . For instance, if Z is preserved by a translation $t_v : A \rightarrow A$, then each fiber $\gamma_Z^{-1}(\xi)$ is also invariant under t_v . Likewise, if $Z = -Z$, then the fiber satisfies $\gamma_Z^{-1}(\xi) = -\gamma_Z^{-1}(\xi)$. Finding further constraints is more challenging.³

An important invariant arising from the fiber $\gamma_Z^{-1}(\xi)$ is the monodromy group $\text{Gal}(\gamma_Z)$; for completeness, we recall its definition below.

Definition 2.4. *Let $f : Y \rightarrow X$ be a generically finite morphism between algebraic varieties. Then there exists a non-empty open subset $U \subseteq X$ such that the restriction $f^{-1}(U) \rightarrow U$ is a finite étale cover. Moving along a closed loop in U induces a permutation of the points in the fiber $f^{-1}(\xi_0)$, which defines the map⁴*

$$\rho_f : \pi_1(U, \xi_0) \rightarrow \text{Aut}(f^{-1}(\xi_0)) \cong S_{\deg f}.$$

The monodromy group is then defined as the image of ρ_f , i.e.,

$$\text{Gal}(f) := \text{Im } \rho_f.$$

It is clear that $\text{Gal}(f)$ doesn't depend on the choice of U .

2.2. Interpolation via hyperplanes. In this subsection, we reinterpret γ_Z using a functorial and more transparent framework. This permits the definition of the conormal Gauss map and its monodromy group for any morphism $\phi : Z \rightarrow \text{Gr}(r, n)$ with $\dim Z = r$, and clarifies the relation between the monodromy group and $\deg \phi$.

Since each non-zero conormal vector $\xi \in T_0^* A$ determines a hyperplane $H_\xi \in \text{Gr}(n-1, T_0 A)$, we have the isomorphisms (To simplify notation, we abbreviate $T_0^* A$ by W .)

$$\begin{aligned} \mathbb{P}T_0^* A &\cong \text{Gr}(n-1, T_0 A), \\ \mathbb{P}\Lambda_{Z^{\text{sm}}} &= \{ (p, \xi) \in Z^{\text{sm}} \times \mathbb{P}T_0^* A \mid \xi|_{T_p Z} \equiv 0 \} \\ &\cong \{ (p, H) \in Z^{\text{sm}} \times \text{Gr}(n-1, W) \mid \phi_Z(p) \subseteq H \} \\ &\cong (\phi_Z, \text{Id})^{-1} I_{r, n-1}, \end{aligned}$$

where

$$I_{r, n-1} := \{ (V, H) \in \text{Gr}(r, W) \times \text{Gr}(n-1, W) \mid V \subseteq H \}$$

is the incidence variety relating $\text{Gr}(r, W)$ and $\text{Gr}(n-1, W)$. In these terms,

$$\begin{aligned} \gamma_Z^{-1}(H) \cap Z^{\text{sm}} &= \{ p \in Z^{\text{sm}} \mid \phi_Z(p) \subseteq H \} \\ &\cong \phi_Z^{-1}(\text{Gr}(r, H)) \end{aligned}$$

is the collection of points whose tangent spaces lie entirely within H .

Geometrically, the monodromy can be described as follows: given a general hyperplane $H \in \mathbb{P}T_0^* A$, its preimage consists of d points p_1, \dots, p_d . Moving H continuously along a closed loop causes these points to permute, and the monodromy group $\text{Gal}(\gamma_Z)$ consists of all permutations obtained this way. With this formulation, it suffices to consider the Gauss map ϕ_Z alone; the inclusion $\iota_Z : Z \rightarrow A$ is no longer required for computing the monodromy group.

³You can imagine the fiber $\gamma_Z^{-1}(\xi)$ as a cluster of stars projected onto a celestial dome. As ξ varies, these points shift, tracing out paths much like stars drifting across the night sky. The constraints that govern them are subtle, like the imagined lines that shape constellations. And in the long arc of variation, monodromy emerges—like the slow turning that replaces Kochab with Polaris among the stars.

⁴In the last isomorphism we implicitly give an order to the fiber $f^{-1}(\xi)$.

Definition 2.5. Let Z be an r -dimensional variety and $\phi : Z \rightarrow \mathrm{Gr}(r, n)$ a morphism. The monodromy group $\mathrm{Mon}(\phi)$ is defined as $\mathrm{Gal}(f_\phi)$, where

$$f_\phi : (\phi, \mathrm{Id})^{-1} I_{r, n-1} \subset Z \times \mathrm{Gr}(n-1, n) \xrightarrow{\pi_2} \mathrm{Gr}(n-1, n).$$

When ϕ is not generically injective, the monodromy group is subject to additional constraints, as captured by the next lemma.

Lemma 2.6. Let $\phi : Z \rightarrow \mathrm{Gr}(r, n)$ be generically k -to-1 onto its image. With a suitable ordering of the fibers, the associated monodromy group $\mathrm{Mon}(\phi)$ is contained in a wreath product

$$S_k \wr S_{d/k} := \left(S_k^{\oplus d/k} \right) \rtimes S_{d/k}.$$

Proof. Consider the diagram below:

$$\begin{array}{ccccc} \phi : & Z & \xrightarrow{k:1} & \mathrm{Im} Z & \hookrightarrow & \mathrm{Gr}(r, n) \\ & \cup & & \cup & & \cup \\ & \{p_1, \dots, p_d\} & \longrightarrow & \{q_1, \dots, q_{d/k}\} & \longrightarrow & \mathrm{Gr}(r, H) \end{array}$$

The fiber $\phi^{-1}(\mathrm{Gr}(r, H_0))$ splits into d/k groups of points, with the monodromy group acting by permutations within each group and among the groups. \square

2.3. Interpolation via higher Brill–Noether theory. Examples of subvarieties of abelian varieties can be obtained in two ways: one may fix an abelian variety and consider cycles within it, or begin with a variety Z and construct a map to an abelian variety, such as the Albanese map. When adopting the latter perspective, we write X for the same variety Z to stress the focus on the variety itself.

In this subsection, we begin with the case $X \rightarrow \mathrm{Alb}(X)$, from which we derive some basic properties of the Albanese map and the tangent Gauss map. The remaining cases can then be treated in an analogous manner.

Let X be a smooth complex projective variety of dimension r , and set $n := \dim_{\mathbb{C}} H^0(X, \Omega_X) = h^{1,0}$. Recall that the Albanese variety of X is defined as

$$\mathrm{Alb}(X) := H^0(X, \Omega_X)^* / H_1(X, \mathbb{Z})_{\mathrm{free}},$$

and the Albanese map is given by (pick a base point $p_0 \in X$)

$$\alpha : X \rightarrow \mathrm{Alb}(X) \quad p \mapsto \left[\omega \mapsto \int_{\gamma: p_0 \sim p} \omega \right].$$

One classical question is the dimension of $\alpha(X)$. For this, consider the cotangent map of α at $p \in X$:

$$T_p^* \alpha : H^0(X, \Omega_X) \rightarrow T_p^* X = H^0(X, \Omega_X|_p).$$

Consider the short exact sequence of coherent sheaves on X (For convenient, we write $\Omega_X(-p) := \Omega_X \otimes \mathcal{I}_p$)

$$0 \longrightarrow \Omega_X(-p) \longrightarrow \Omega_X \longrightarrow \Omega_X|_p \longrightarrow 0$$

which induces a long exact sequence

$$\begin{array}{ccccccc} & & H^1(X, \Omega_X(-p)) & \longrightarrow & H^1(X, \Omega_X) & \longrightarrow & 0 \\ & \swarrow & & & & & \\ 0 & \longrightarrow & H^0(X, \Omega_X(-p)) & \longrightarrow & H^0(X, \Omega_X) & \xrightarrow{T_p^* \alpha} & H^0(X, \Omega_X|_p) \end{array}$$

The proposition below follows from standard arguments in homological algebra:

Proposition 2.7. *For a general point $p \in X$,*

$$\begin{aligned} \dim_{\mathbb{C}} \alpha(X) &= \operatorname{rank} T_p^* \alpha \\ &= n - h^0(X, \Omega_X(-p)) \\ &= h^1(X, \Omega_X(-p)) - h^1(X, \Omega_X). \end{aligned}$$

In particular,

$$\begin{aligned} \alpha \text{ is surjective} &\iff h^0(X, \Omega_X(-p)) = 0 \\ \alpha \text{ is finite onto image} &\iff h^0(X, \Omega_X(-p)) = n - r \\ \alpha \text{ is constant} &\iff h^0(X, \Omega_X(-p)) = n \\ &\iff h^1(X, \Omega_X(-p)) = h^1(X, \Omega_X) \\ &\iff n = 0. \end{aligned}$$

We will concentrate on the case where α is finite onto its image.⁵ Under this assumption, we set $Z = X$ and $A = \operatorname{Alb}(Z)$. The corresponding Gauss map is then a rational map:

$$\begin{array}{ccc} \phi_Z : Z & \dashrightarrow & \operatorname{Gr}(r, T^0 A) \cong \operatorname{Gr}(n-r, H^0(X, \Omega_X)) \\ p & \longmapsto & H^0(X, \Omega_X(-p)) \end{array}$$

and we have the isomorphisms

$$\begin{aligned} \mathbb{P}T_0^* A &\cong \mathbb{P}H^0(X, \Omega_X), \\ \mathbb{P}\Lambda_Z &= \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^* A \mid \omega(p) = 0 \} \\ &\cong \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^* A \mid \omega \in H^0(X, \Omega_X(-p)) \} \\ &\cong (\phi_Z, \operatorname{Id})^{-1} I_{n-r,1}, \end{aligned}$$

where

$$I_{n-r,1} := \{ (V, [\omega]) \in \operatorname{Gr}(n-r, n) \times \operatorname{Gr}(1, n) \mid \omega \in V \}$$

is the incidence variety relating $\operatorname{Gr}(n-r, n)$ and $\operatorname{Gr}(1, n)$. In that case,

$$\gamma_Z^{-1}([\omega]) = \{ p \in X \mid \omega(p) = 0 \}$$

is the zero set of section $\omega \in H^0(X, \Omega_X)$. The number $(-1)^r \deg \gamma_Z$ is the index in the Poincaré–Hopf index formula, and the monodromy group $\operatorname{Gal}(\gamma_Z)$ serves as a more refined invariant, encoding subtler aspects of the geometry.

The next proposition shows when the Gauss map ϕ_Z is not generic injective.

Proposition 2.8. *When α is finite onto its image,*

$$\begin{aligned} \phi_Z \text{ is not generic injective} &\iff \text{For general } p \in X, \text{ exists } q \neq p \text{ such that } h^0(X, \Omega_X(-p-q)) = n-r \\ &\stackrel{\text{6}}{\iff} \text{For general } p \in X, \text{ exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r \\ &\iff \text{For all } p \in X, \text{ exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r. \end{aligned}$$

⁵The general method remains valid in the broader setting, but the Gauss map then takes values in a different space.

⁶When $n = r$, the map φ_Z has a point as its target, so the equivalence is trivial. When $n > r$, the implication “ \Rightarrow ” is immediate. For the converse “ \Leftarrow ”, it suffices to show that the image of

$$\left\{ S \in X^{[2]} \mid h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r \right\}$$

under the map $\pi_X : X^{[2]} \rightarrow X^{(2)}$ does not include the diagonal. Indeed, we can choose a section $s \in H^0(X, \Omega_X)$ that cuts out finitely many reduced points p_1, \dots, p_d in X . (Well this maybe not so true, since ϕ_Z is not always regular, such a section s may vanish along some indeterminacy locus. However, for a general s , there will always be

This last step relies on the following lemma, combined with the closedness of the tautological correspondence in $X \times X^{[2]}$.

Lemma 2.9. *Let $m \in \mathbb{Z}_{>0}$. The function*

$$h^\Omega : X^{[m]} \longrightarrow \mathbb{Z}_{\geq 0} \quad S \longmapsto h^0(X, \Omega_X \otimes \mathcal{I}_S)$$

is Zariski upper semicontinuous.

Sketch of proof. Consider the coherent sheaf $\mathcal{F} \in \text{Coh}(X^{[m]} \times X)$ characterized by the property

$$\mathcal{F}|_{\{S\} \times X} \cong \Omega_X \otimes \mathcal{I}_S,$$

The proposition then follows directly from the semicontinuity theorem [14, 28.1.1]. \square

The stratification of $X^{[m]}$ by h^Ω offers a natural generalization of Brill–Noether theory beyond the setting of curves.

At the end of this subsection, let us turn to the setting of a general abelian variety A and a smooth subvariety $\iota_Z : Z \hookrightarrow A$, where $n = \dim_{\mathbb{C}} A$ and $r = \dim_{\mathbb{C}} Z$. Observe that ι_Z factors through the Albanese variety of Z :

$$\iota_Z : Z \xrightarrow{\alpha_Z} \text{Alb}(Z) \xrightarrow{\pi} A$$

We shall also assume that Z generates A ; it then follows that the map π is surjective. The cotangent map of ι_Z at a point $p \in Z$ factors through $H^0(Z, \Omega_Z)$:

$$T_p^* \iota_Z : T_p^* A \hookrightarrow H^0(Z, \Omega_Z) \longrightarrow T_p^* Z$$

For convenience, abbreviate $V := T_0^* A \cong T_p^* A$, and view V as a subspace of $H^0(Z, \Omega_Z)$.

Proposition 2.10. *Assume that Z is embedded in A and generates A , and let $V := T_0^* A$. Then*

$$\dim_{\mathbb{C}} H^0(Z, \Omega_Z(-p)) \cap V = n - r \quad \text{for all } p \in Z$$

It follows that the Gauss map is a regular morphism

$$\begin{array}{ccc} \phi_Z : Z & \longrightarrow & \text{Gr}(r, T^0 A) \cong \text{Gr}(n - r, V) \\ p & \longmapsto & H^0(Z, \Omega_Z(-p)) \cap V \end{array}$$

Furthermore,

ϕ_Z is not generic injective

$$\iff \text{For general } p \in Z, \text{ exists } q \neq p \text{ such that } h^0(Z, \Omega_Z(-p - q)) \cap V = n - r$$

$$\iff \text{For all } p \in Z, \text{ exists } S \in Z^{[2]} \text{ such that } p \in S \text{ and } h^0(Z, \Omega_Z \otimes \mathcal{I}_S) \cap V = n - r.$$

Question 2.11. *For an r -dimensional smooth variety Z whose Albanese map α_Z is an embedding, which subspaces $V \subset H^0(Z, \Omega_Z)$ can arise as the cotangent space of some abelian quotient variety of $\text{Alb}(Z)$?*

The precise formulation of this question is somewhat ambiguous, and depending on the interpretation, it may turn out to be either relatively straightforward or extremely difficult. On the one hand, the classification of abelian subvarieties of A via symmetric idempotents in $\text{End}_{\mathbb{Q}}(A)$ is well established [2, Theorem 5.3.2]. On the other hand, explicitly computing $\text{End}_{\mathbb{Q}}(A)$ can be highly nontrivial when A is a non-simple Albanese variety—even in the case where A arises from a curve.

at least one isolated zero p_1 , which is sufficient for our purposes.) Then for any $S \in \pi_X^{-1}(p_1)$, we have

$$H^0(X, \Omega_X \otimes \mathcal{I}_S) \subsetneq H^0(X, \Omega_X(-p_1)).$$

3. FAMILIES OF SUBVARIETIES

In this section, we move from the study of monodromy groups to a more direct analysis of the subvarieties themselves. Given an initial subvariety, one can naturally generate a family of subvarieties. Our goal here is to define these families and investigate their properties.

Proposition 3.1. *All irreducible conic Lagrangian cycles in T^*A are of the form Λ_Z for some irreducible subvariety $Z \subset A$. This yields a one-to-one correspondence between irreducible conic Lagrangian cycles in T^*A and irreducible subvarieties of A :*

$$\{ \text{irreducible conic Lagrangian cycles in } T^*A \} \longleftrightarrow \{ \text{irreducible subvarieties in } A \}$$

Sketch of proof. For any irreducible conic Lagrangian cycle $\Lambda \subset T^*A$, let Z denote the image of Λ under the natural projection $T^*A \rightarrow A$. Our goal is to show that $\Lambda = \Lambda_Z$.

- By definition, $\Lambda \subset T^*A|_Z$.
- Since Λ is conic, we have $s(Z) \subset \Lambda$, where $s : A \rightarrow T^*A$ denotes the zero section.
- Since Λ is Lagrangian and $s(Z) \subset \Lambda$, we have $\Lambda \subset \Lambda_Z$.
- Since Λ_Z is irreducible with $\dim_{\mathbb{C}} \Lambda = \dim_{\mathbb{C}} \Lambda_Z = n$, we have $\Lambda = \Lambda_Z$.

□

Why do we shift attention from Z to Λ_Z as the main object of study? One reason is the uniformity of Λ_Z : it always has dimension n , and in most cases, the natural map $\Lambda_Z \rightarrow T_0^*A$ is generically finite, with fibers lying inside A .

Definition 3.2 (Clean cycle). *An irreducible Lagrangian cycle $\Lambda \subset T^*A$ is called clean if the composed projection*

$$\Lambda \longrightarrow T^*A \twoheadrightarrow T_0^*A$$

is generically finite.

Another important reason is that the space of weighted clean conic Lagrangian cycles naturally acquires a convolution structure, arising from the group law on A , which plays a central role in the analysis.

Proposition 3.3. *The group of weighted clean conic Lagrangian cycles*

$$\begin{aligned} \mathcal{L}(A) &:= \{ \text{weighted clean conic Lagrangian cycles in } T^*A \} \\ &= \left\{ \sum_{\substack{Z_i \subset A \\ \text{irr clean}}} n_i \Lambda_{Z_i} \mid n_i \in \mathbb{Z} \right\} \end{aligned}$$

has a natural convolution structure as follows:

$$\begin{aligned} \Lambda_{Z_1} \circ \Lambda_{Z_2} &= \text{the clean part of } (a, \text{Id}_{T_0^*A})_* (\Lambda_{Z_1} \times_{T_0^*A} \Lambda_{Z_2}) \\ &= \overline{(a, \text{Id}_U)_* (\Lambda_{Z_1}|_U \times_U \Lambda_{Z_2}|_U)} \end{aligned}$$

where

$$U := \left\{ \xi \in T_0^*A \mid \deg \phi_{Z_i} = \# \phi_{Z_i}^{-1}(\xi) \text{ for } i = 1, 2 \right\}$$

and $a : A \times A \rightarrow A$ is the addition map in A . The general convolution is defined by \mathbb{Z} -linear extension.

Sketch of proof. To establish the claim, it suffices to show that $\Lambda_{Z_1} \circ \Lambda_{Z_2}$ defines a weighted conic Lagrangian cycle. The conic property follows directly from the definition, while the Lagrangian condition can be verified at a general point $(p_1 + p_2, \xi) \in \Lambda_{Z_1} \circ \Lambda_{Z_2}$. □

We now consider the projective versions of all objects involved, so that we may make use of properness. To simplify notation, we abbreviate $\mathbb{P}T_0^*A$ by \mathbb{P}^\vee .

Lemma 3.4.

(1) Suppose that $\mathbb{P}\Lambda_{Z_1}, \mathbb{P}\Lambda_{Z_1} \subset \mathbb{P}T^*A$ admit monodromy representations

$$\rho_{\gamma_{Z_i}} : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_i}^{-1}(\xi_0)),$$

then $\mathbb{P}\Lambda_{Z_1} \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_{Z_2} \subset A \times A \times \mathbb{P}^\vee$ admits monodromy representation given by

$$(\rho_{\gamma_{Z_1}}, \rho_{\gamma_{Z_2}}) : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_1}^{-1}(\xi_0) \times \gamma_{Z_2}^{-1}(\xi_0)),$$

(2) When $Z_1 = Z_2 = Z$, we obtain an one-to-one correspondence:

$$\left\{ \begin{array}{l} \text{irr components of } \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \end{array} \right\}$$

Sketch of proof. Statement (1) holds by definition. The proof of (2) reduces to the following purely topological statement:

Claim 3.5. *Let $\pi : E \longrightarrow B$ be a (unramified) covering space over a manifold B with deck transformation group G , then*

$$\{ \text{connected components of } E \times_B E \} \longleftrightarrow \{ G\text{-orbits of } \pi^{-1}(b_0) \times \pi^{-1}(b_0) \}.$$

The claim follows directly from the correspondence between covering spaces over B and $\pi_1(B)$ -sets; see [6, Theorem 1.38]. \square

Generalizing the argument of Lemma 3.4, we arrive at the following lemma.

Lemma 3.6. *For $d = \deg \gamma_Z$, $\xi_0 \in T_0^*A$ a general point, write*

$$\begin{aligned} \mathbb{P}\Lambda_Z^{\times d} &:= \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \cdots \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \subset A^d \times \mathbb{P}^\vee \\ \gamma_Z^{-1}(\xi_0)^d &:= \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \times \cdots \times \gamma_Z^{-1}(\xi_0) \subset A^d \end{aligned}$$

(1) we obtain an one-to-one correspondence:

$$\left\{ \begin{array}{l} \text{irr components of } \mathbb{P}\Lambda_Z^{\times d} \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0)^d \end{array} \right\}$$

(2) Write

$$\begin{aligned} \Delta_d &:= \{ (p_1, \dots, p_d) \in A^d \mid p_i = p_j \text{ for some } i \neq j \} \subset A^d \\ \mathbb{P}\Lambda_Z^{[d]} &:= \overline{(\mathbb{P}\Lambda_Z^{\times d} \setminus (\Delta_d \times \mathbb{P}^\vee))} \big|_U \subset A^d \times \mathbb{P}^\vee \end{aligned}$$

Fix a general point $\xi_0 \in \mathbb{P}^\vee$ and a well-order for $\gamma_Z^{-1}(\xi_0)$, one can identify $S_d \cong \gamma_Z^{-1}(\xi_0)^d \setminus \Delta_d$, and

$$\left\{ \text{irr components of } \mathbb{P}\Lambda_Z^{[d]} \right\} \longleftrightarrow \left\{ \text{Gal}(\gamma_Z)\text{-orbits of } S_d \right\}$$

In reference, Δ_d is usually called the big diagonal.

From this point on, we fix an irreducible component of $\mathbb{P}\Lambda_Z^{[d]}$, denoted by $\mathbb{P}\Lambda_Z^{\text{univ}}$. As we will see in Definition A, this variety generates all subvarieties within the families under consideration.

Definition 3.7 (The subvariety $Z^{(m)}$). *For any tuple $(m) = (m_1, \dots, m_d) \in \mathbb{Z}^d$, we define the weighted sum map*

$$a^{(m)} : A^d \longrightarrow A \quad (p_1, \dots, p_d) \longmapsto \sum_{i=1}^d m_i p_i.$$

We also define

$$\mathbb{P}\Lambda_Z^{(m)} := \left(a^{(m)}, \text{Id}_{\mathbb{P}^\vee} \right)_* \mathbb{P}\Lambda_Z^{\text{univ}}.$$

as the (projectivized) weighted Lagrangian cycle in $\mathbb{P}T^*A$. The projective cycle $\mathbb{P}\Lambda_Z^{(m)}$ is irreducible but may appear with multiplicities. We can therefore write

$$\mathbb{P}\Lambda_Z^{(m)} = c_Z^{(m)} \mathbb{P}\Lambda_{Z^{(m)}}$$

where $c_Z^{(m)} \in \mathbb{Z}_{>0}$ and $Z^{(m)} \subset A$ are uniquely determined. This gives rise to a family of subvarieties parametrized by \mathbb{Z}^d .

The next lemma gathers some basic properties of $Z^{(m)}$. Observe that $S_d = \text{Aut}(\gamma_Z^{-1}(\xi_0))$ acts naturally on \mathbb{Z}^d via

$$g(m) = (m_{g(1)}, \dots, m_{g(d)}) \in \mathbb{Z}^d.$$

Lemma 3.8.

- (1) For all $g \in \text{Gal}(\gamma_Z)$, we have $Z^{g(m)} = Z^{(m)}$, $c_Z^{g(m)} = c_Z^{(m)}$;
- (2) For $(m) = (1, 0, \dots, 0) \in \mathbb{Z}^d$, $Z^{(m)} = Z$;
- (3) For all $(m), (m') \in \mathbb{Z}^d$, we have $\mathbb{P}\Lambda_Z^{(m)} \circ \mathbb{P}\Lambda_Z^{(m')} = \mathbb{P}\Lambda_Z^{(m+m')}$;
- (4) The group $\langle \mathbb{P}\Lambda_{Z^{(m)}} \rangle_{\text{Abel}}$ is closed under the convolution product.

Question 3.9. *Can we compute the dimension and the homology class for these subvarieties $Z^{(m)}$?*

At first glance, this question may appear naive, but it turns out to be extremely challenging. The main difficulty lies in determining the homology class of $\mathbb{P}\Lambda_Z^{\text{univ}}$. Once this is known, one can compute the pushforward to obtain the homology class of $\mathbb{P}\Lambda_Z^{(m)}$, from which both $\dim_{\mathbb{C}} Z^{(m)}$ and the homology class of $Z^{(m)}$ (up to a scalar factor $c_Z^{(m)}$) can be extracted.

However, standard intersection theory yields only the homology class of $\mathbb{P}\Lambda_Z^{[d]}$, via the Chern–Mather class of Z . At present, there is no known method for computing the homology class of a single irreducible component of $\mathbb{P}\Lambda_Z^{[d]}$. Although we can distinguish these components topologically, we lack tools to compute their topological invariants, which presents an unfortunate limitation.

Nevertheless, Question 3.9 can still be addressed in certain special situations. For instance, when $\text{Gal}(\gamma_Z) = S_n$, the space $\mathbb{P}\Lambda_Z^{[d]}$ is known to be irreducible, enabling explicit computation. Similarly, when $Z = -Z$ and $\text{Gal}(\gamma_Z) = W(C_{d/2})$, one can impose additional conditions (e.g., $p_{2l-1} = -p_{2l}$) to guarantee the uniqueness of $\mathbb{P}\Lambda_Z^{\text{univ}}$, which again allows us to perform the computation.

Question 3.10. *Suppose $d = 2d_0$ with d_0 even, $(m) \in \mathbb{Z}^d$, and $g \in W(C_{d/2}) \setminus W(D_{d/2})$. Is it possible to construct an example with $Z = -Z$ and $\text{Gal}(\gamma_Z) = W(D_{d/2})$, yet $\dim_{\mathbb{C}} Z^{g(m)} \neq \dim_{\mathbb{C}} Z^{(m)}$?*

Question 3.11. *Suppose that Z is the Fano surface of a smooth cubic threefold, and let $A := \text{Alb}(Z)$. How can we compute the homology class of $Z^{(m)}$?*

4. WHEN IS MONODROMY GROUP SMALL

In this section, we discuss examples drawn from my ongoing work, focusing on the construction of subvarieties and the computation of their monodromy groups.

Broadly speaking, there are two approaches to constructing examples. One may start with a given variety Z and attempt to embed it into an abelian variety—this always factors through its Albanese $\text{Alb}(Z)$. Alternatively, one may begin with an abelian variety A and construct subvarieties by intersecting suitable divisors. Unfortunately, even in the case where Z is a curve, we have not fully resolved Question 6.1.

4.1. Curves, basic results. A complete answer to Question 6.3 is known when $Z = C$ is a curve. A curve $C \subseteq \mathbb{P}^{n-1}$ is said to be nondegenerate if it is not contained in any hyperplane $H \subseteq \mathbb{P}^{n-1}$.

Proposition 4.1 (See [1, p111] for a detailed proof). *Suppose that $\iota_C : C \subseteq \mathbb{P}^{n-1}$ is an irreducible nondegenerate curve of degree d , then $\text{Mon}(\iota_C) \cong S_d$.*

Sketch of proof. Because S_d is generated by its transpositions, we are reduced to verifying that:

- $\text{Mon}(\iota_C)$ acts doubly transitively on the fiber;
- $\text{Mon}(\iota_C)$ contains a transposition.

□

Proposition 4.2. *Let $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$ be an irreducible nondegenerate curve of degree $d/2$, and let $h : C \rightarrow C'$ be a degree 2 ramified covering, with ramification occurring at at least one smooth point of C' . Then $\text{Mon}(\iota_{C'} \circ h) \cong S_2^{\oplus d/2} \rtimes S_{d/2}$ is the hyperoctahedral group/signed symmetric group.*

Sketch of proof. By Lemma 2.6 we know that $\text{Mon}(\iota_{C'} \circ h) \subseteq S_2^{\oplus d/2} \rtimes S_{d/2}$. By Lemma 4.3, we are reduced to verifying that:

- The quotient map $\text{Mon}(\iota_{C'} \circ h) \rightarrow \text{Mon}(\iota_{C'}) \cong S_{d/2}$ is surjective;
- $\text{Mon}(\iota_{C'} \circ h)$ contains a transposition of a pair of points in the fiber of h .

□

Lemma 4.3. *Let G be a subgroup of $S_2^{\oplus m} \rtimes S_m$, acting naturally on the set $\pm 1, \dots, \pm m$. If the projection $G \rightarrow S_m$ is surjective and the transposition σ_0 of ± 1 lies in G , then $G = S_2^{\oplus m} \rtimes S_m$.*

Sketch of proof. (with help from Chenji Fu) Let ε_i denote the transposition of $\pm i$. For any $\sigma \in S_m$, choose a lift $\tilde{\sigma} \in G$, then

$$\varepsilon_{\sigma(1)} = \tilde{\sigma} \circ \sigma_0 \circ \tilde{\sigma}^{-1} \in G.$$

Thus, $S_2^{\oplus m} \subset G$, and since G maps onto S_m , we obtain $G = S_2^{\oplus m} \rtimes S_m$. □

Example 4.4. *Let C be a smooth curve of genus g embedded in its Jacobian $A := \text{Jac}(C)$ via the Abel–Jacobi map $\text{AJ}_C : C \hookrightarrow A$.*

When C is non-hyperelliptic, the corresponding Gauss map

$$|\omega_C| : C \rightarrow \mathbb{P}^{g-1}$$

makes C as an irreducible nondegenerate curve of degree $2g - 2$, by Proposition 4.1 we get

$$\text{Gal}(\gamma_C) \cong S_{2g-2}.$$

When C is hyperelliptic, the corresponding Gauss map is $2 : 1$ onto a rational normal curve $R \subset \mathbb{P}^{g-1}$:

$$|\omega_C| : C \xrightarrow{2:1} R \hookrightarrow \mathbb{P}^{g-1}$$

By Proposition 4.2 we get

$$\text{Gal}(\gamma_C) \cong S_2^{\oplus g-1} \rtimes S_{g-1}.$$

In fact, a degree $2 : 1$ map does not give rise to any exceptional monodromy groups beyond those listed in Table 1.

Definition 4.5 (big monodromy group). *We refer to the big monodromy group as any group of the following types:*

notation	name	alias
$W(A_{m+1}) = S_m$	full symmetric group	
$W(C_m) = S_2^{\oplus m} \rtimes S_m$	signed symmetric group	hyperoctahedral group
$W(D_m) = (S_2^{\oplus m})_0 \rtimes S_m$	even-signed symmetric group	demihyperoctahedral group

TABLE 1. big monodromy group

In practice, the term “big monodromy group” refers to the full symmetric group S_n when the subset $Z \subset A$ is not symmetric, and to the (even-)signed symmetric group when $Z \subset A$ is symmetric.

Proposition 4.6. *Let $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$ be an irreducible nondegenerate curve of degree $d/2$, and let $h : C \rightarrow C'$ be a degree 2 ramified covering. Then*

$$\text{Mon}(\iota_{C'} \circ h) \cong W(C_{d/2}) \text{ or } W(D_{d/2}).$$

Sketch of proof. By Lemma 2.6 we know that $\text{Mon}(\iota_{C'} \circ h) \subseteq W(C_{d/2})$. By Lemma 4.7, we are reduced to verifying that:

- The quotient map $\text{Mon}(\iota_{C'} \circ h) \longrightarrow \text{Mon}(\iota_{C'}) \cong S_{d/2}$ is surjective;
- (signed doubly transitive) $\text{Mon}(\iota_{C'} \circ h)$ acts transitively on pairs (x, y) with $x \neq \pm y$.

□

Lemma 4.7. *Let G be a subgroup of $W(C_m)$, acting naturally on the set $\pm 1, \dots, \pm m$. If the projection $G \rightarrow S_m$ is surjective then*

$$G \cong W(C_m) \text{ or } W(D_m) \text{ or } S_2 \times S_m \text{ or } S_m.$$

Sketch of proof. (with help from Chenji Fu) Let H denote the kernel of the natural quotient map $G \rightarrow S_m$. Then $H \subseteq (S_2)^{\oplus m}$ is stable under the action of S_n . There are only four possible forms that H can take:⁷

- $H = 0$. Then $G \cong S_m$.
- $H = \langle (-1, \dots, -1) \rangle \cong S_2$. Then $G \cong S_2 \times S_m$.
- $H = (S_2^{\oplus m})_0$. Then G is a index 2 subgroup of $W(C_m)$, so $G \cong W(D_m)$.⁸
- $H = S_2^{\oplus m}$. Then $G = W(C_m)$.

□

We are particularly interested in identifying cases where the monodromy group is small (i.e., not big), which can occur only when $k > 2$, where k denotes the degree of the map $\phi : C \rightarrow \mathbb{P}^{n-1}$. On the other hand, when both $k > 2$ and $n > 2$, Lemma 2.6 ensures that the resulting monodromy group is indeed small. Thus, in the case of curves, Question 6.1 is resolved except in the following three cases:

- (1) $n > 2$ and $k = 2$. By Proposition 4.6, the monodromy group is known to be large, although its precise structure remains undetermined.
- (2) $n = 2$. Here, A is an abelian surface and C corresponds to a divisor.
- (3) $n > 2$ and $k > 2$. Finding an example of this would guarantee a case of small monodromy.

Question 4.8. *Can we find any curve $C \subset A$ not stable under any translation on A , and for which the associated monodromy group is small?*

Assume that $n > 2$. Can we find any curve $C \subset A$ not stable under any translation on A , whose Gauss map has degree $d > 2$?

Example 4.9. *If we replace A with \mathbb{C}^3 in Question 4.8, we can construct curves $C \subset \mathbb{C}^3$ that are not preserved under any linear transformation but whose Gauss maps fail to be generically injective. Here is one such example:*

Consider the curve $C \subset \mathbb{C}^3$ defined by the parametrization

$$r(t) = \left(t, (t-1)e^t, \frac{1}{4}(2t^2 - 2t + 1)e^{2t} \right).$$

Its derivative is

$$r'(t) = (1, te^t, t^2 e^{2t}),$$

so the Gauss map fails to be generically injective.

⁷Here is a brief argument showing that H must be one of 0 , S_2 , $(S_2^{\oplus m})_0$, or $S_2^{\oplus m}$. If H contains an element $h = (a_1, \dots, a_m)$ with $a_i \neq a_j$ for some $i \neq j$, then

$$(ij)h + h = (1, \dots, \underset{\substack{\uparrow \\ i\text{-th}}}{-1}, \dots, \underset{\substack{\uparrow \\ j\text{-th}}}{-1}, \dots, 1) \in H,$$

implying $(S_2^{\oplus m})_0 \subseteq H$. Hence, H must be either $(S_2^{\oplus m})_0$ or $S_2^{\oplus m}$. Otherwise, if all $h \in H$ have identical coordinates, then H is either 0 or S_2 .

⁸Check [stackexchange discussions](#)

To answer Question 4.8, the starting point is to seek curves with decomposable Jacobians. A classical source of such examples is a covering $h : C \rightarrow C'$ of $\deg(h) > 1$ where $g(C') > 0$, in which case $\text{Jac}(C)$ decomposes. Nevertheless, this does not exhaust all possibilities: Paweł Borówka pointed out to me an example, proved in [3, Exercise 11.7], where $\text{Jac}(C)$ is decomposable although C does not cover a curve of positive genus.

4.2. Curves in Prym variety. To address Question 4.8, we seek explicit examples, and the case of a Prym curve mapping to its associated Prym variety offers a natural starting point for investigation.

Notations 4.10. Suppose C' is a smooth projective curve of genus $g(C')$, and let B be an effective (possibly zero) divisor on C' . For any line bundle $\eta \in \text{Pic}(C')$ such that $\eta^{\otimes 2} \cong \mathcal{O}_{C'}(B)$, one obtains a double cover $h : C \rightarrow C'$ of smooth projective curves, ramified precisely over B . The associated involution on C is denoted by ι , and the triple (C, C', h) is referred to as the Prym pair.

Recall that the Prym variety $A := \text{Prym}(C/C')$ is defined as the connected component of the identity in

$$\ker \left[\text{Nm} : \text{Jac}(C) \longrightarrow \text{Jac}(C') \right].$$

The Abel–Prym map is defined by

$$\text{AP}_{C/C'} : C \longrightarrow A \quad p \longmapsto \mathcal{O}_C(p - \iota(p)).$$

The classical theory treats the case where the Prym variety A is principally polarized—this occurs exactly when $B = \emptyset$ or $B = \{p_0, q_0\}$, that is, when h is either unramified or ramified at two points.⁹ For clarity and focus, we restrict our attention to these cases. Table 2 lists the relevant numerical data.¹⁰

	$g(C)$	$g(C')$	$\deg \eta$
$B = \emptyset$	$2n + 1$	$n + 1$	0
$B = \{p_0, q_0\}$	$2n$	n	1

TABLE 2. numerical data of Prym pair

The Abel–Prym map $\text{AP}_{C/C'}$ does not always behave as nicely as the Abel–Jacobi map; it may fail to be an embedding. When C is hyperelliptic, the map $\text{AP}_{C/C'}$ fails to be generically injective, and therefore falls outside the scope of our discussion. When C is non-hyperelliptic, we may regard it as a subvariety of A for our purposes, although it may be not strictly embedded.¹¹ However, it remains unclear whether C is stable under any translation in A .

In the Prym setting, the corresponding Gauss map γ_C factors through h :

$$\begin{aligned} \gamma_C : C &\xrightarrow{h} C' \xrightarrow{|\omega_{C'} \otimes \eta|} \mathbb{P}^{n-1} = \mathbb{P}(\text{H}^0(\omega_{C'} \otimes \eta)^*) \\ &\cong \text{Gr}(n-1, \text{H}^0(\omega_{C'} \otimes \eta)) \\ \tilde{p} &\longmapsto p \longmapsto \text{H}^0(\omega_{C'} \otimes \eta(-p)) \end{aligned}$$

Lemma 4.11. When $\text{gon}(C') > 4$, $|\omega_{C'} \otimes \eta|$ is injective. As a result, the monodromy group is big.

⁹See [4, Theorem 3.2.6] for the proof.

¹⁰As usual, $n = \dim_{\mathbb{C}} A$ is the dimension of the abelian variety.

¹¹For a detailed description of the map $\text{AP}_{C/C'}$, see [2, Proposition 12.5.2, Corollary 12.5.6]. Although $\text{AP}_{C/C'}$ may collapse p_0 and q_0 , this does not affect the relevant computation of $\text{Gal}(\gamma_Z)$.

Proof. Suppose that $|\omega_{C'} \otimes \eta|$ is not injective, we need to find a line bundle of degree 4 and rank ≥ 1 . In fact, for $p \neq q$,

$$\begin{aligned} |\omega_{C'} \otimes \eta|(p) &= |\omega_{C'} \otimes \eta|(q) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-p - q)) &= 1 && \text{By [14, 19.2.8]} \\ \iff h^0(\omega_{C'} \otimes \eta(-p - q)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^{\vee}(p + q)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

When $B = \emptyset$, write $\eta^{\vee}(p + q) = \mathcal{O}_{C'}(p' + q')$, then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^{\vee})^{\otimes 2}(2p + 2q) = \mathcal{O}_{C'}(2p' + 2q') \in g_4^1;$$

When $B = \{p_0, q_0\}$, write $\eta^{\vee}(p + q) = \mathcal{O}_{C'}(p')$, then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^{\vee})^{\otimes 2}(2p + 2q + p_0 + q_0) = \mathcal{O}_{C'}(2p' + p_0 + q_0) \in g_4^1.$$

□

Remark 4.12. Based on the strategy used in the proof of Lemma 2.10, we can in fact obtain stronger results. For $p \in C'$,

$$\begin{aligned} |\omega_{C'} \otimes \eta| \text{ is ramified at } p &&& \text{(i.e., the tangent map is 0 at } p) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-2p)) &= 1 && \text{By [14, 19.2.9]} \\ \iff h^0(\omega_{C'} \otimes \eta(-2p)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^{\vee}(2p)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

For the remainder of this discussion, we focus exclusively on the case $B = \emptyset$. Combining both,

$$\begin{aligned} |\omega_{C'} \otimes \eta| \text{ is not generically injective} \\ \iff \text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } h^0(\eta^{\vee}(p + q)) &= 1 \\ \iff \text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } \eta^{\vee} + p \in C' + C' - q \\ \iff \eta^{\vee} + C' \subseteq C' + C' - C'. \end{aligned}$$

It is expected that

$$\eta^{\vee} + C' \subseteq C' + C' - C' \iff \eta^{\vee} \in C' - C'. \quad (4.1)$$

If so, writing $\eta^{\vee} = \mathcal{O}_{C'}(p - q)$, we obtain

$$\mathcal{O}_{C'}(2p) = \mathcal{O}_{C'}(2q) \in g_2^1,$$

which implies that C' is hyperelliptic. By Fact 4.13, this forces C to be hyperelliptic as well, and hence it lies outside the scope of our discussion.

Unfortunately, (4.1) does not hold in general. A counterexample, communicated to me by Gavril Farkas, will be presented in Proposition 4.15.

Fact 4.13 (From [13, p346]). *We have an explicit classification of $2 : 1$ unramified covers of a hyperelliptic curve C' . Let $\pi : C' \rightarrow \mathbb{P}^1$ be the double cover ramified at $\{p_1, \dots, p_{2g+2}\}$, then every such unramified double cover of C' arises from some line bundle in C of the form*

$$\eta := \mathcal{O}_{C'}(p_{i_1} + \dots + p_{i_l} - p_{i_{l+1}} - \dots - p_{i_{2l}}) = \mathcal{O}_{C'}\left(\sum_{j=1}^{2l} p_{i_j} - 2l \cdot p_1\right).$$

When $l = 1$, the associated covering curve C is also hyperelliptic.

Question 4.14. *Can one find a non-hyperelliptic curve C' and a nontrivial line bundle $\eta \in \text{Jac}(C')$ satisfying $\eta^{\otimes 2} \cong \mathcal{O}_{C'}$, such that*

$$\eta^{\vee} + C' \subseteq C' + C' - C'?$$

If such a pair exists, can we find a case where the resulting curve $C \subset A$ is not preserved under any translation on A ?

As a key example relevant to Question 4.14, I include here a result of Gavril Farkas.

Proposition 4.15 (See [5, Proposition 2.8]). *Let C be a general smooth curve of genus $g = 2i$, and suppose $D \in W_{i+1}^1(C)$ is a pencil of minimal degree—i.e., a line bundle $\mathcal{O}_C(D) \in \text{Pic}^{i+1}(C)$ with $h^0(C, \mathcal{O}_C(D)) = 2$. Define*

$$\mathcal{L} := K_C - 2D \in \text{Pic}^{2i-4}(C),$$

then \mathcal{L} satisfies

$$\mathcal{L} + C \subset C_{2i-2} - C, \quad \text{but} \quad \mathcal{L} \notin C_{2i-3} - C.$$

Remark 4.16. Take $i = 2$, the line bundle \mathcal{L} provides a counterexample to (4.1). However, this example does not directly yield a counterexample to Question 4.14, due to Proposition 4.17.

Proposition 4.17. *The line bundle \mathcal{L} in Proposition 4.15 never satisfies*

$$\mathcal{L}^{\otimes 2} \cong \mathcal{O}_C \quad \text{but} \quad \mathcal{L} \not\cong \mathcal{O}_C,$$

i.e.,

$$2K_C = 4D \quad \text{but} \quad K_C \neq 2D.$$

Proof. Since C is a smooth non-hyperelliptic curve of genus 4, its canonical embedding realizes C as a complete intersection of an irreducible quadric \tilde{Q} and an irreducible cubic surface \tilde{K} in \mathbb{P}^3 . According to [12, Lemma 2.1], every divisor $D \in W_3^1(C)$ arises as the intersection $D = L \cap \tilde{K}$, where L is a line contained in the quadric surface \tilde{Q} .

When \tilde{Q} is a quadric cone, there exists a hyperplane H such that $H \cap \tilde{Q} = 2L$, implying $K_C = 2D$; when \tilde{Q} is smooth, we have $H \cap \tilde{Q} = L + L'$, with L' being the residual line. Interpreting $2K_C \in |\mathcal{O}_{\mathbb{P}^3}(2)|$ as a quadric $Q \subset \mathbb{P}^3$, we obtain

$$2K_C = 4D$$

$$\iff \text{there exists a quadric } Q \subset \mathbb{P}^3 \text{ such that } Q \cap C = 4D$$

$$\iff \text{there exists a quadric } Q \subset \mathbb{P}^3 \text{ such that } Q \cap \tilde{Q} \cap \tilde{K} = 4D$$

$$\iff \text{there exists a quadric } Q \subset \mathbb{P}^3 \text{ such that } Q \cap \tilde{Q} = 4L \quad \text{by Lemma 4.18}$$

However, via Lemma 4.19, such case won't happen at all. \square

Lemma 4.18 was inspired by a discussion with Søren Gammelgaard.

Lemma 4.18. *Let Q and \tilde{Q} be reduced quadric surfaces in \mathbb{P}^3 , and let $\tilde{K} \subset \mathbb{P}^3$ be a cubic surface. Assume that \tilde{Q} and \tilde{K} are smooth and that $\tilde{Q} \cap \tilde{K}$ is a complete intersection. Let $L \subset \tilde{Q}$ be a line, and set $D := L \cap \tilde{K}$. Then,*

$$Q \cap \tilde{Q} \cap \tilde{K} = 4D \iff Q \cap \tilde{Q} = 4L.$$

Proof. We only need to prove the “ \implies ” direction. Suppose that $Q \cap \tilde{Q} \cap \tilde{K} = 4D$. Then:¹²

- Since Q intersects L in at least three points, we have $L \subset Q$.
- Write $Q \cap \tilde{Q} = L + R$, where R is the residual curve. Then $\deg R = 3$ and $R \cap \tilde{K} = 3D$.
- If $R \subset \mathbb{P}^3$ is non-degenerate, then R is a rational normal curve of degree 3, which cannot pass through three distinct collinear points. This contradicts $3D \subset R \cap L$.
- If R is degenerate, then it lies in a hyperplane $H \cong \mathbb{P}^2$. Since $R \subset H \cap \tilde{Q}$ is contained in a conic, we can write $R = L' + R'$, where L' is a line and R' is a conic. From the inclusion $3D \subset (L' + R') \cap L$, it follows that $L' = L$ and $R' = 2L$.
- Therefore, $Q \cap \tilde{Q} = L + R = 4L$.

\square

¹²The previous argument is incorrect. For example, let $X = \{F = 0\}$ be a smooth cubic plane curve and let $L = \{G = 0\}$ be a line meeting X at three points p_1, p_2, p_3 . Then for some nonzero $a \in \mathbb{C}$, the cubic $Y_a := \{F + aG^3 = 0\}$ is smooth, and $X \cap Y_a = 3(p_1 + p_2 + p_3)$.

Lemma 4.19. *Let Q and \tilde{Q} be two reduced quadric surfaces in \mathbb{P}^3 , and assume that \tilde{Q} is smooth. Then Q and \tilde{Q} cannot intersect a line with multiplicity 4.*

Proof. We prove by contradiction. Without loss of generality, assume that $Q \cap \tilde{Q} = 4L$, where $L = \{[x : y : 0 : 0]\} \subset \mathbb{P}^3$ is a line. We can write down the equations of the quadrics:

$$\begin{aligned} Q : f(z_0, z_1, z_2, z_3) &= 0 \\ \tilde{Q} : \tilde{f}(z_0, z_1, z_2, z_3) &= 0 \end{aligned}$$

where

$$\begin{aligned} f(z_0, z_1, z_2, z_3) &= \sum_{i+j \leq 2} a_{ij}(z_2, z_3) z_0^i z_1^j \\ &= a_{01}(z_2, z_3) z_0 + a_{10}(z_2, z_3) z_1 + a_{00}(z_2, z_3) \\ &= (b_{02} z_2 + b_{03} z_3) z_0 + (b_{12} z_2 + b_{13} z_3) z_1 + a_{00}(z_2, z_3) \\ \tilde{f}(z_0, z_1, z_2, z_3) &= \sum_{i+j \leq 2} \tilde{a}_{ij}(z_2, z_3) z_0^i z_1^j \\ &= \tilde{a}_{01}(z_2, z_3) z_0 + \tilde{a}_{10}(z_2, z_3) z_1 + \tilde{a}_{00}(z_2, z_3) \\ &= (\tilde{b}_{02} z_2 + \tilde{b}_{03} z_3) z_0 + (\tilde{b}_{12} z_2 + \tilde{b}_{13} z_3) z_1 + \tilde{a}_{00}(z_2, z_3). \end{aligned}$$

Since \tilde{Q} is smooth,

$$\tilde{b}_{02}\tilde{b}_{03}, \tilde{b}_{12}\tilde{b}_{13}, \tilde{b}_{02}\tilde{b}_{12}, \tilde{b}_{03}\tilde{b}_{13} \neq 0.$$

Fix $t \in \mathbb{C}$. Restricting Q and \tilde{Q} to the hyperplane $\{z_1 = tz_0\}$, we obtain two conics:

$$\begin{cases} f(z_0, tz_0, z_2, z_3) = 0 \\ \tilde{f}(z_0, tz_0, z_2, z_3) = 0 \end{cases}$$

which intersect at the point $p_t := [1 : t : 0 : 0]$ with multiplicity 4. Writing down these equations explicitly, we get

$$\begin{cases} ((b_{02} + tb_{12})z_2 + (b_{03} + tb_{13})z_3) z_0 + a_{00}(z_2, z_3) = 0 \\ ((\tilde{b}_{02} + t\tilde{b}_{12})z_2 + (\tilde{b}_{03} + t\tilde{b}_{13})z_3) z_0 + \tilde{a}_{00}(z_2, z_3) = 0, \end{cases}$$

so the corresponding tangent lines at p_t are given by

$$\begin{cases} (b_{02} + tb_{12})z_2 + (b_{03} + tb_{13})z_3 = 0 \\ (\tilde{b}_{02} + t\tilde{b}_{12})z_2 + (\tilde{b}_{03} + t\tilde{b}_{13})z_3 = 0 \end{cases}$$

Since the conics intersect with multiplicity 4 at p_t , these tangent lines must coincide for every t , so

$$(b_{02} + tb_{12})(\tilde{b}_{03} + t\tilde{b}_{13}) = (b_{03} + tb_{13})(\tilde{b}_{02} + t\tilde{b}_{12}) \quad \text{for every } t \in \mathbb{C}.$$

By a change of coordinates, we may assume that $\tilde{a}_{10}(z_2, z_3) = z_3$. This implies that $\tilde{b}_{02} = 0$ and $\tilde{b}_{03} = 1$. Furthermore, we have $\tilde{b}_{12} \neq 0$, and

$$\begin{cases} b_{12}\tilde{b}_{13} = \tilde{b}_{12}b_{13} \\ b_{02} = 0 \\ b_{12} = b_{03}\tilde{b}_{12} \end{cases}$$

so $b_{13} = b_{03}\tilde{b}_{13}$.

Now choose $(z_2, z_3) \neq (0, 0)$ such that $a_{00}(z_2, z_3) = b_{03} \cdot \tilde{a}_{00}(z_2, z_3)$. Take $t \in \mathbb{C}$ such that

$$(\tilde{b}_{02} + t\tilde{b}_{12})z_2 + (\tilde{b}_{03} + t\tilde{b}_{13})z_3 = t\tilde{b}_{12}z_2 + (1 + t\tilde{b}_{13})z_3 \neq 0,$$

and take

$$z_0 := \frac{\tilde{a}_{00}}{t\tilde{b}_{12}z_2 + (1 + t\tilde{b}_{13})z_3} z_3.$$

Then the point $[z_0 : tz_0 : z_2 : z_3]$ lies in $Q \cap \tilde{Q}$ but not in L , yielding a contradiction. \square

Recently, we obtained a partial answer to Question 4.14, as illustrated in the following example.

Example 4.20. Let C' be a non-hyperelliptic bielliptic curve, and let $\text{pr} : C' \rightarrow E$ denote a $2 : 1$ covering onto an elliptic curve. For any nontrivial 2-torsion line bundle $\eta_0 \in \text{Pic}^0(E)[2]$, the pullback $\eta := \text{pr}^* \eta_0$ satisfies $\eta^{\otimes 2} \cong \mathcal{O}_{C'}$ and

$$\eta^\vee + C' \subseteq C' + C' - C'.$$

Proof of Example 4.20. For any $x \in C'$, write $x_0 : \text{pr}(x)$, then

$$\deg \eta_0^\vee(x_0) = 1 \implies \text{there exists } y_0 \in E \text{ such that } \eta_0^\vee(x_0) = \mathcal{O}_E(y_0).$$

Write $\text{pr}^{-1}(x_0) = \{x, x'\}$, $\text{pr}^{-1}(y_0) = \{y, y'\}$, we get

$$\eta^\vee(x + x') = \mathcal{O}_{C'}(y + y').$$

i.e.,

$$\eta^\vee(x) = \mathcal{O}_{C'}(y + y' - x).$$

□

The next proposition tells us, for the most time the resulting curve $C \subset A$ is not invariant under any non-trivial translation of A . Therefore, Example 4.20 serve as an example of Question 4.14 and 4.8.

Proposition 4.21. In Example 4.20, let C be the curve corresponding to $\eta \in \text{Pic}^0(C')[2]$. If $\text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$, then C admits three intermediate quotients over E . Denote them by C' , C_1 , and C_2 , where C_1 and C_2 are the two intermediate curves different from C' .

$$\begin{array}{ccc} \begin{array}{c} C \\ \downarrow h \\ C' \\ \downarrow \\ E \\ \downarrow \\ \mathbb{P}^1 \end{array} & & \begin{array}{ccccc} & C & & & \\ & \swarrow h_1 & \downarrow h & \searrow h_2 & \\ C_1 & & C' & & C_2 \\ & \searrow & \downarrow & \swarrow & \\ & E & & & \\ & \downarrow & & & \\ & \mathbb{P}^1 & & & \end{array} \\ \text{Gal}(C/E) \cong \mathbb{Z}/4\mathbb{Z} & & \text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \end{array}$$

If either $\text{Gal}(C/E) \cong \mathbb{Z}/4\mathbb{Z}$, or $\text{Gal}(C/E) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ with the coverings $h_i : C \rightarrow C_i$ ramified, then the curve $C \subset A$ is not fixed under any non-trivial translation on A .

Proof. We prove by contradiction. Assume there exists a non-trivial translation $\sigma : A \rightarrow A$ preserving C . Then the restriction $\sigma|_C : C \rightarrow C$ is an automorphism, and the induced map

$$h_\sigma : C \longrightarrow C/\sigma$$

is an unramified double covering. Since σ is a translation, the Gauss map γ_C necessarily factors through h_σ :

$$\begin{array}{ccccc} \gamma_C : C & \xrightarrow{2:1} & C' & \xrightarrow{2:1} & E \subset \mathbb{P}^{n-1} \\ & \searrow h_\sigma & & \nearrow \exists! & \\ & & C/\sigma & & \end{array}$$

Without loss of generality, we may assume that $\deg(h_\sigma) = 2$. By assumption, C/σ coincides with C' , and $\sigma|_C$ is the involution ι associated with the cover $h : C \rightarrow C'$, which extends to an involution $\tilde{\iota}$ of A . According to [2, Proposition 12.4.2], $\tilde{\iota}$ is a reflection. Hence $\sigma \circ \tilde{\iota}^{-1}$ is another reflection fixing C , which is impossible.

□

In fact, the Castelnuovo–Severi inequality, brought to my attention by Gavril Farkas, shows that Example 4.20 is the only example for Question 4.14 when $g(C') > 9$.

Fact 4.22 (Castelnuovo–Severi inequality, [8, p26, Corollary]). *Let C be a smooth projective curve equipped with two ramified coverings $f_i : C \rightarrow C_i$ of degrees d_i ($i = 1, 2$). Suppose that there is no morphism $h : C \rightarrow \tilde{C}$ with $\deg(h) > 1$ such that both f_1 and f_2 factor through h . Then*

$$g(C) \leq d_1 \cdot g(C_1) + d_2 \cdot g(C_2) + (d_1 - 1)(d_2 - 1).$$

Proposition 4.23. *If, in Question 4.14, we additionally require that C' be non-hyperelliptic and non-bielliptic, then any such curve must satisfy $g(C') \leq 9$.*

Proof. Assume that C' is an example of Question 4.14.

Step 1. we can find two distinct g_4^1 of C' .

For any $p \in C'$, there exist $q, p', q' \in C'$ such that

$$\eta^\vee(p + q) = \mathcal{O}_{C'}(p' + q').$$


This implies

$$\mathcal{O}_{C'}(2p + 2q) = \mathcal{O}_{C'}(2p' + 2q'),$$

which defines a degree-4 covering $f_1 : C' \rightarrow \mathbb{P}^1$ ramified at p, q, p', q' . Choosing $\tilde{p} \in C'$ outside the ramification locus of f_1 and repeating the construction yields another g_4^1 , denoted $f_2 : C' \rightarrow \mathbb{P}^1$.

Step 2. If f_1 and f_2 factor through a common map $h : C' \rightarrow \tilde{C}$ with $\deg(h) > 1$, then C' is hyperelliptic or bielliptic.

Indeed, in this case $\deg(h) = 2$, and the Castelnuovo–Severi inequality applied to $(\tilde{C}, \tilde{f}_1, \tilde{f}_2)$ gives $g(\tilde{C}) \leq 1$, so \tilde{C} is either \mathbb{P}^1 or an elliptic curve.



Step 3. If C' is neither hyperelliptic nor bielliptic, then the Castelnuovo–Severi inequality yields $g(C') \leq 9$. \square

5. TANNAKIAN FORMALISM

For simplicity, we work over the base field $\kappa = \mathbb{C}$. Let A denote a fixed complex abelian variety, and let $\text{Perv}(A)$ denote the category of perverse sheaves on A with coefficients in \mathbb{Q} . For any algebraic group G , we denote by $\text{Rep}(G)$ the category of algebraic representations of G .

Following the approach of [11], we work in the quotient category $\overline{\text{Perv}}(A) = \text{Perv}(A)/N(A)$, where $N(A) \subset \text{Perv}(A)$ is the Serre subcategory of negligible complexes. A complex \mathcal{F} is defined to be negligible if $\chi(A, \mathcal{F}) = 0$. This quotient category admits a natural convolution structure, and every finitely generated tensor subcategory of it is Tannakian, with a reductive Tannaka group G (see [11, Thm 7.1 & Cor 9.2]). In particular, for any perverse sheaf $\delta \in \overline{\text{Perv}}(A)$, the full subcategory generated by δ is categorically equivalent to the representation category of an algebraic group G :

$$\langle \delta, * \rangle \cong \text{Rep}(G).$$

6. DELETE LATER

Question 6.1. *Suppose that the subvariety $Z \subset A$ is not stable under any translation on A . Are there criterias to compute the monodromy group $\text{Gal}(\gamma_Z)$? Furthermore, what kinds of groups can appear as $\text{Gal}(\gamma_Z)$ for suitable choices of $Z \subset A$?*

Based on the discussion above, Question 6.1 reduces to Question 6.2 and Question 6.4, with Question 6.3 appearing as a special case of Question 6.2.

Question 6.2. *In the situation of Definition 2.5, how can we compute $\text{Mon}(\phi)$?*

Question 6.3. *Let $Z \subseteq \text{Gr}(r, n)$ be a subvariety of dimension r such that $[Z] \cdot [\text{Gr}(r, H)] \neq 0$ in $H_r(\text{Gr}(r, n); \mathbb{Z})$. What can be said about its monodromy group?*

Question 6.4. *For a map $\phi : Z \rightarrow \text{Gr}(r, n)$, when is it induced from some inclusion $Z \subset A$?*

REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Volume I*, volume 267 of *Grundlehren Math. Wiss.* Springer, Cham, 1985.
- [2] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 302 of *Grundlehren Math. Wiss.* Berlin: Springer, 2nd augmented ed. edition, 2004.
- [3] Paweł Borówka and Angela Ortega. Prym varieties and Prym map. *Rend. Ist. Mat. Univ. Trieste*, 54:34, 2022. Id/No 14.
- [4] Giulio Codogni and Thomas Krämer. Semicontinuity of Gauss maps and the Schottky problem. *Math. Ann.*, 382(1-2):607–630, 2022.
- [5] Gavril Farkas and Michael Kemeny. The generic Green-Lazarsfeld secant conjecture. *Invent. Math.*, 203(1):265–301, 2016.
- [6] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge University Press, 2002.
- [7] Ariyan Javanpeykar, Thomas Krämer, Christian Lehn, and Marco Maculan. The monodromy of families of subvarieties on abelian varieties. Preprint, arXiv:2210.05166 [math.AG] (2022), 2022.
- [8] Ernst Kani. On Castelnuovo’s equivalence defect. *J. Reine Angew. Math.*, 352:24–70, 1984.
- [9] Thomas Krämer. Cubic threefolds, Fano surfaces and the monodromy of the Gauss map. *Manuscr. Math.*, 149(3-4):303–314, 2016.
- [10] Thomas Krämer. Summands of theta divisors on Jacobians. *Compos. Math.*, 156(7):1457–1475, 2020.
- [11] Thomas Krämer and Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties. *J. Algebr. Geom.*, 24(3):531–568, 2015.
- [12] Eduard Looijenga. A ball quotient parametrizing trigonal genus 4 curves. *Nagoya Math. J.*, 254:366–378, 2024.
- [13] David Mumford. *The red book of varieties and schemes. Includes the Michigan lectures (1974) on “Curves and their Jacobians”*, volume 1358 of *Lect. Notes Math.* Berlin: Springer, 2nd, expanded ed. with contributions by Enrico Arbarello edition, 1999.
- [14] Ravi Vakil. *The rising sea: Foundations of algebraic geometry. preprint*, 2017.

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, BERLIN, 12489, GERMANY,
 Email address: email:xiaoxiang.zhou@hu-berlin.de