




Inclusion $\subseteq \rightarrow$
 dominance Δ




Lexicographic ordering



$\dim \chi_\lambda = \# \text{ irr compo}$
Hook length formula



dim of Springer fibration





Special Series

S_1

S_2

S_3



pt
trivial rep
reduced perm rep
standard rep

two row case

$B \cong G/B \cong \mathcal{F}(n)$
alternating rep

Q: What's the dim of $\{\text{Cartan subalg of } \mathfrak{sl}_n(\mathbb{C})\}$?

$$\begin{array}{ccc}
 \{\text{Borel subalg}\} & \longleftrightarrow & \{\text{Borel subgroup}\} \xrightarrow{\text{fix } B} G/B \\
 \downarrow \text{finite surj?} & & \downarrow \\
 \{\text{Cartan subalg}\} & \longleftrightarrow & \{\text{Maximal torus}\} \xrightarrow{\text{fix } T} G/N_G(T) \\
 & & \uparrow \text{G transitive?} \quad \text{finite?} \quad \text{structure?}
 \end{array}$$

Examples of Springer fiber. $E_{\lambda_1, \dots, \lambda_n} = \text{Springer fiber of } \begin{bmatrix} J_{\lambda_1} & & \\ & \ddots & \\ & & J_{\lambda_n} \end{bmatrix}$ $J_\lambda = \begin{bmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix}$


\square $[0]$ $E_1 = \{0 \in \mathbb{C}\} = \bullet$ $\mathcal{F} \mathcal{F} \mathcal{B} \mathcal{X} \mathcal{X} \mathcal{V} \mathcal{S} E$

\square $\begin{bmatrix} 0 & 1 \\ & 0 \end{bmatrix}$ $E_2 = \{0 \in \langle v_1 \rangle \subseteq \mathbb{C}^2\} = \bullet$

\square $\begin{bmatrix} 0 & & \\ & 0 & \\ & & 0 \end{bmatrix}$ $E_{1,1} = \{0 \in \langle ? \rangle \subseteq \mathbb{C}^2\} = \mathbb{P}^1$

\square $\begin{bmatrix} 0 & 1 & \\ & 0 & 1 \\ & & 0 \end{bmatrix}$ $E_3 = \{0 \in \langle v_1 \rangle \subseteq \langle v_1, v_2 \rangle \subseteq \mathbb{C}^3\} = \bullet$

\square $\begin{bmatrix} 0 & 1 & \\ & 0 & \\ & & 0 \end{bmatrix}$ $E_{2,1} = \{0 \in \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3\} = \mathbb{P}^1 \vee \mathbb{P}^1$



$E_{1,1} \rightarrow E_{2,1}$
 $E_2 \rightarrow E_{2,1}$
 \downarrow
 $\mathbb{P}^1 \subseteq \mathbb{P}^2$

$F_{e_1} = \{0 \in \langle e_1 \rangle \subseteq \langle e_1, ? \rangle \subseteq \mathbb{C}^3\} \hookrightarrow \begin{bmatrix} 0 & 1 & \\ & 0 & \\ & & 0 \end{bmatrix}$


\Downarrow

$\mathbb{P}^1 = E_{1,1} = \{0 \in \langle ? \rangle \subseteq \mathbb{C}^3\} \hookrightarrow \begin{bmatrix} 0 & \\ & 0 & \\ & & 0 \end{bmatrix}$

$F_{e_3} = \{0 \in \langle e_3 \rangle \subseteq \langle e_3, ? \rangle \subseteq \mathbb{C}^3\} \hookrightarrow \begin{bmatrix} 0 & 1 & \\ & 0 & \\ & & 0 \end{bmatrix}$

$\{pt\} = E_2 = \{0 \in \langle ? \rangle \subseteq \mathbb{C}^3\} \hookrightarrow \begin{bmatrix} 0 & 1 & \\ & 0 & \\ & & 0 \end{bmatrix}$

\square $\begin{bmatrix} 0 & & \\ & 0 & \\ & & 0 \end{bmatrix}$ $E_{1,1,1} = \{0 \in \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3\} = \mathcal{F}(3)$




$\mathcal{F}(1) - \mathcal{F}(2) - \mathcal{F}(3) - \mathcal{F}(4) - \mathcal{F}(5)$
 $\downarrow \quad \downarrow \quad \downarrow \quad \downarrow$
 $\mathbb{P}^1 \quad \mathbb{P}^2 \quad \mathbb{P}^3 \quad \mathbb{P}^4$

$E_1 - E_{1,1} - E_{1,1,1} - E_{1,1,1,1} - E_{1,1,1,1,1}$
 $\downarrow \quad \downarrow \quad \downarrow \quad \downarrow$
 $\bullet \quad \bullet \quad \bullet \quad \bullet$


$$\begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{bmatrix}$$

$$E_{3,1} = \{0 \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \mathbb{C}^4\} = \mathbb{P}^1 \vee \mathbb{P}^1 \vee \mathbb{P}^1$$




$$\begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{bmatrix}$$

$$\bar{E}_{2,2} = \{0 \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \mathbb{C}^4\} = \mathbb{P}^1\text{-bundle} \vee_{\mathbb{P}^1} \mathbb{P}^1\text{-bundle}$$



$$\begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{bmatrix}$$

$$\bar{E}_{2,1,1} = \{0 \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \langle ? \rangle \subseteq \mathbb{C}^4\}$$



Q: Can we construct an affine paving from this kind of fibration?

Q: How to understand the Weyl group action on Springer fibration?

1	3	4	6
2	5	7	8



" "

Bertrand's ballot theorem



"bracket pairing"

Define M^λ to be the complex vector space with basis the tabloids $\{T\}$ of shape λ , with λ a partition of n .

$$(3) \quad a_T = \sum_{p \in R(T)} p, \quad b_T = \sum_{q \in C(T)} \text{sgn}(q)q, \quad v_T = b_T \cdot \{T\} = \sum_{q \in C(T)} \text{sgn}(q)\{q \cdot T\}.$$

These elements, and the product

$$c_T = b_T \cdot a_T,$$

Define the **Specht module** S^λ to be the subspace of M^λ spanned by the elements v_T , as T varies over all numberings of λ .

Proposition 1 For each partition λ of n , S^λ is an irreducible representation of S_n . Every irreducible representation of S_n is isomorphic to exactly one S^λ .

Proposition 2 The elements v_T , as T varies over the standard tableaux on λ , form a basis for S^λ .

Young tabloid

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}$$

$$\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}$$

$$\begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}$$




$$X = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$0 \in \text{Im } X^2 \subseteq \text{Im } X \subseteq \mathbb{C}^3$$

$$B_{2,1} = \{0 \in \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \mathbb{C}^3\}$$


$$0 \in \left(\begin{array}{c} \text{Ker } X \subseteq \text{Ker } X^2 \\ \langle e_1, e_3 \rangle \quad \langle e_1, e_2, e_3 \rangle \end{array} \right) \subseteq \mathbb{C}^3$$



$$X = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$0 \in \quad 0 \in \quad \langle e_1, e_3 \rangle$$


$$B_{2,1} = \{0 \in \langle ? \rangle \subseteq \langle ?, ? \rangle \subseteq \langle ?, ?, ? \rangle \subseteq \mathbb{C}^4\}$$



$$\text{Im } \pi_2 = \{V \subseteq \mathbb{C}^4 \mid \dim_{\mathbb{C}} V = 2, X V \subseteq V\} \subseteq \text{Gr}(4,2)$$

↑ not smooth!

Global description?



Cohomology of B_λ