

# SUBVARIETIES IN COMPLEX ABELIAN VARIETIES

XIAOXIANG ZHOU

## CONTENTS

1. Basic setting	1
2. Searching for examples	4
3. Families of subvarieties	10
4. Tannakian formalism	13
References	13

This document is intended to collect the questions and doubts that arose during my research this year. For many of these problems, I have consulted my fellow students, my supervisor, and various other people I've met. However, most of them remain in the realm of folklore—problems that are likely known but for which I could not find a reference. On the other hand, some of the questions may not appear particularly interesting unless their underlying motivations are clearly explained. Therefore, I'll try to provide relevant background and outline some initial, perhaps naive, ideas while listing the problems along the way. Any responses, answers, or references are most welcome and will be added to keep this document updated.

## 1. BASIC SETTING

For simplicity, we work over the base field  $\kappa = \mathbb{C}$ , and by a variety we mean a integral separated scheme of finite type over  $\mathbb{C}$ . Let  $A/\mathbb{C}$  be an abelian variety of dimension  $n$ , and let  $Z \subseteq A$  be an irreducible closed subvariety of dimension  $r$ . We denote by  $\iota_Z : Z \hookrightarrow A$  the inclusion morphism.<sup>1</sup>

**1.1. Gauss map.** The goal of my research is to understand the geometry of  $Z$ , and the main tool for the subvariety geometry is the Gauss map. The Gauss map describe the tangent space information at each point:

$$\phi_Z : Z^{\text{sm}} \longrightarrow \text{Gr}(r, T_0 A) \quad p \longmapsto T_p Z \subseteq T_p A \cong T_0 A$$

Any map to the Grassmannian  $\text{Gr}(r, n)$  is induced by a rank  $r$  vector bundle together with  $n$  global sections. In this case, the map  $\phi_Z$  is induced by the tangent bundle  $\mathcal{T}_{Z^{\text{sm}}}$  and the sections

$$H^0(A, \mathcal{T}_{Z^{\text{sm}}}) \otimes_{\mathbb{C}} \mathcal{O}_{Z^{\text{sm}}} \rightarrow \mathcal{T}_{Z^{\text{sm}}}.$$

---

*Date:* June 10, 2025.

<sup>1</sup>I'm not sure whether we should consider the more general cases in the future—such as working over a field of characteristic  $p$ , letting  $A$  be a semiabelian variety or a complex torus, or allowing  $\iota$  to be a covering onto its image. For now, I will omit these possibilities from this document.

**1.2. Conormal variety.** This concept may already be familiar to many readers, so we briefly recall the definition. On the smooth locus, the normal and conormal bundles behave well as vector bundles:<sup>2</sup>

$$\mathcal{N}_{Z^{\text{sm}}/A} := \mathcal{T}_A|_{Z^{\text{sm}}} / \mathcal{T}_{Z^{\text{sm}}} \quad \Lambda_{Z^{\text{sm}}} := \mathcal{N}_{Z^{\text{sm}}/A}^* = \ker \left( \Omega_A|_{Z^{\text{sm}}} \rightarrow \Omega_{Z^{\text{sm}}} \right).$$

The conormal variety  $\Lambda_Z$  is just the closure of  $\Lambda_{Z^{\text{sm}}}$  viewed as a subvariety in  $T^*A$ :

$$\Lambda_Z := \overline{\Lambda_{Z^{\text{sm}}}} \subset T^*A \cong A \times T_0^*A$$

this is a conical Lagrangian cycle in  $T^*A$ .

Moreover, the projectivized conormal variety

$$\mathbb{P}\Lambda_Z := \overline{\mathbb{P}\Lambda_{Z^{\text{sm}}}} \subset \mathbb{P}T^*A \cong A \times \mathbb{P}T_0^*A$$

is a Legendrian cycle in the contact variety  $A \times \mathbb{P}T_0^*A$ .  $\mathbb{P}\Lambda_{Z^{\text{sm}}}$  is a  $\mathbb{P}^{r-1}$ -bundle over  $Z^{\text{sm}}$ , and the map

$$\gamma_Z : \mathbb{P}\Lambda_Z \subset A \times \mathbb{P}T_0^*A \longrightarrow \mathbb{P}T_0^*A$$

is generically finite (i.e., clean) when  $Z$  is (an integral variety) of general type, see [6, Theorem 2.8 (1)].

A lot of geometry of  $Z$  is encoded in the map  $\gamma_Z$ . For instance, if  $Z$  is smooth and lies inside  $A$ , then

$$\deg \gamma_Z = (-1)^r \chi(Z)$$

tells us the Euler characteristic of  $Z$ .

Further insight can be gained by analyzing the fibers of  $\gamma_Z$ . These fibers, though finite, are not arbitrary—they obey hidden structural rules. For instance, if  $Z$  is preserved by a translation  $t_v : A \rightarrow A$ , then each fiber  $\gamma_Z^{-1}(\xi)$  is also invariant under  $t_v$ . Likewise, if  $Z = -Z$ , then the fiber satisfies  $\gamma_Z^{-1}(\xi) = -\gamma_Z^{-1}(\xi)$ . Outside these special configurations, it becomes more challenging to identify further constraints.<sup>3</sup>

An important invariant arising from the fiber  $\gamma_Z^{-1}(\xi)$  is the monodromy group  $\text{Gal}(\gamma_Z)$ ; for completeness, we recall its definition below.

**Definition 1.1.** *Define*

$$U = \{ \xi \in \mathbb{P}T_0^*A \mid \# \gamma_Z^{-1}(\xi) = \deg \gamma_Z \}.$$

*Moving along a loop in  $U$  induces a permutation of the points in the fiber  $\gamma_Z^{-1}(\xi_0)$ , which defines the map*

$$\rho_{\gamma_Z} : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_Z^{-1}(\xi_0)) = S_{\deg \gamma_Z}.$$

*The monodromy group is then defined as the image of  $\rho_{\gamma_Z}$ , i.e.,*

$$\text{Gal}(\gamma_Z) := \text{Im } \rho_{\gamma_Z}.$$

**Question 1.2.** *Suppose that the subvariety  $Z \subset A$  is not stable under any translation on  $A$ . Are there known algorithms to compute the monodromy group  $\text{Gal}(\gamma_Z)$ ? Furthermore, what kinds of groups can appear as  $\text{Gal}(\gamma_Z)$  for suitable choices of  $Z \subset A$ ?*

We will try to compute  $\text{Gal}(\gamma_Z)$  for a number of specific cases in Section 2. Three special cases are already treated in [7, Theorem 9], and we will generalize the strategies there.

<sup>2</sup>This is more symmetric when writing them as short exact sequences:

$$\begin{aligned} 0 &\longrightarrow \mathcal{T}_{Z^{\text{sm}}} \longrightarrow \mathcal{T}_A|_{Z^{\text{sm}}} \longrightarrow \mathcal{N}_{Z^{\text{sm}}/A} \longrightarrow 0 \\ 0 &\longrightarrow \Lambda_{Z^{\text{sm}}} \longrightarrow \Omega_A|_{Z^{\text{sm}}} \longrightarrow \Omega_{Z^{\text{sm}}} \longrightarrow 0 \end{aligned}$$

<sup>3</sup>You can imagine the fiber  $\gamma_Z^{-1}(\xi)$  as a cluster of stars projected onto a celestial dome. As  $\xi$  varies, these points shift, tracing out paths much like stars drifting across the night sky. The constraints that govern them are subtle, like the imagined lines that shape constellations. And in the long arc of variation, monodromy emerges—like the slow turning that replaces Kochab with Polaris among the stars.

**1.3. Interpolation via hyperplanes.** Before delving into examples, we reinterpret  $\gamma_Z$  using a functorial and more transparent framework, enabling a decomposition of Question 1.2 into two primary subquestions.

Recognizing that each non-zero conormal vector  $\xi \in T_0^*A$  determines a hyperplane  $H_\xi \in \text{Gr}(n-1, T_0A)$ , we establish the isomorphisms

$$\begin{aligned} \mathbb{P}T_0^*A &\cong \text{Gr}(n-1, T_0A) \doteq (\mathbb{P}^{n-1})^\vee, \\ \mathbb{P}\Lambda_{Z^{\text{sm}}} &= \{ (p, \xi) \in Z^{\text{sm}} \times \mathbb{P}T_0^*A \mid \xi|_{T_pZ} \equiv 0 \} \\ &\cong \{ (p, H) \in Z^{\text{sm}} \times \text{Gr}(n-1, n) \mid \phi_Z(p) \subseteq H \} \\ &\cong (\phi_Z, \text{Id})^{-1} I_{r, n-1}, \end{aligned}$$

where

$$I_{r, n-1} := \{ (V, H) \in \text{Gr}(r, n) \times \text{Gr}(n-1, n) \mid V \subseteq H \}$$

is the incidence variety relating  $\text{Gr}(r, n)$  and  $\text{Gr}(n-1, n)$ . In that case,

$$\begin{aligned} \gamma_Z^{-1}(H) \cap Z^{\text{sm}} &= \{ p \in Z^{\text{sm}} \mid \phi_Z(p) \subseteq H \} \\ &\cong \phi_Z^{-1}(\text{Gr}(r, H)) \end{aligned}$$

is the collection of points whose tangent spaces lie entirely within  $H$ .

Geometrically, the monodromy can be described as follows: given a general hyperplane  $H$ , its preimage consists of  $d$  points  $p_1, \dots, p_d$ . Moving  $H$  continuously along a loop causes these points to permute, and the monodromy group  $\text{Gal}(\gamma_Z)$  consists of all permutations obtained this way. With this new formulation, it suffices to consider the Gauss map  $\phi_Z$  alone; the inclusion  $\iota_Z : Z \rightarrow A$  is no longer required for computing the monodromy group.

**Definition 1.3.** Let  $Z$  be an  $r$ -dimensional variety and  $\phi : Z \rightarrow \text{Gr}(r, n)$  a morphism. Suppose that for some  $d \in \mathbb{N}_{>0}$ , the set

$$U := \{ H \in \text{Gr}(n-1, n) \mid \#\phi^{-1}(\text{Gr}(r, H)) = d \}$$

is non-empty open in  $\text{Gr}(n-1, n)$ <sup>4</sup>. The monodromy group  $\text{Gal}(\phi)$  is defined as the image of

$$\rho_{\gamma_Z} : \pi_1(U, H_0) \rightarrow \text{Aut}(\phi^{-1}(\text{Gr}(r, H_0))) \cong S_d.$$

When  $\phi$  is not generically finite onto its image, the monodromy group is subject to additional constraints, as captured by the next lemma.

**Lemma 1.4.** When  $\phi : Z \rightarrow \text{Gr}(r, n)$  is generically  $k$ -to-1 onto its image, the monodromy group  $\text{Gal}(\phi)$  is contained in the wreath product

$$S_k \wr S_{d/k} := (S_k^{\oplus d/k}) \rtimes S_{d/k}.$$

*Proof.* Consider the diagram below:

$$\begin{array}{ccccc} \phi : & Z & \xrightarrow{k:1} & \text{Im } Z & \hookrightarrow & \text{Gr}(r, n) \\ & \cup & & \cup & & \cup \\ & \{p_1, \dots, p_d\} & \longrightarrow & \{q_1, \dots, q_{d/k}\} & \longrightarrow & \text{Gr}(r, H) \end{array}$$

The fiber  $\phi^{-1}(\text{Gr}(r, H_0))$  splits into  $d/k$  groups of points, with the monodromy group acting by permutations within each group and among the groups.  $\square$

Based on the discussion above, Question 1.2 reduces to Question 1.5 and Question 1.7, with Question 1.6 appearing as a special case of Question 1.5.

**Question 1.5.** In the situation of Definition 1.3, how can we compute  $\text{Gal}(\phi)$ ?

**Question 1.6.** Let  $Z \subseteq \text{Gr}(r, n)$  be a subvariety of dimension  $r$  such that  $[Z] \neq 0$  in  $H_r(\text{Gr}(r, n); \mathbb{Z})$ . What can be said about its monodromy group?

<sup>4</sup>Here, the fiber is understood set-theoretically; multiplicities are not taken into account.

**Question 1.7.** For a map  $\phi : Z \rightarrow \mathrm{Gr}(r, n)$ , when is it induced from some inclusion  $Z \subset A$ ?

## 2. SEARCHING FOR EXAMPLES

In this section, we discuss examples drawn from my ongoing work, focusing on the construction of subvarieties and the computation of their monodromy groups.

Broadly speaking, there are two approaches to constructing examples. One may start with a given variety  $Z$  and attempt to embed it into an abelian variety—this always factors through its Albanese  $\mathrm{Alb}(Z)$ . Alternatively, one may begin with an abelian variety  $A$  and construct subvarieties by intersecting suitable divisors. Unfortunately, even in the case where  $Z$  is a curve, we have not fully resolved Question 1.2.

**2.1. Curves, basic results.** A complete answer to Question 1.6 is known when  $Z = C$  is a curve.

**Proposition 2.1** (See [1, p111] for a detailed proof). *Suppose that  $\iota_C : C \subseteq \mathbb{P}^{n-1}$  is an irreducible nondegenerate curve of degree  $d$ , then  $\mathrm{Gal}(\iota_C) \cong S_d$ .*

*Sketch of proof.* Because  $S_d$  is generated by its transpositions, we are reduced to verifying that:

- $\mathrm{Gal}(\iota_C)$  acts doubly transitively on the fiber;
- $\mathrm{Gal}(\iota_C)$  contains a transposition.

□

**Proposition 2.2.** *Let  $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$  be an irreducible nondegenerate curve of degree  $d/2$ , and let  $h : C \rightarrow C'$  be a degree 2 ramified covering, with ramification occurring at at least one smooth point of  $C'$ . Then  $\mathrm{Gal}(\iota_{C'} \circ h) \cong S_2^{\oplus d/2} \rtimes S_{d/2}$  is the hyperoctahedral group/signed symmetric group.*

*Sketch of proof.* By Lemma 1.4 we know that  $\mathrm{Gal}(\iota_{C'} \circ h) \subseteq S_2^{\oplus d/2} \rtimes S_{d/2}$ . By Lemma 2.3, we are reduced to verifying that:

- The quotient map  $\mathrm{Gal}(\iota_{C'} \circ h) \rightarrow \mathrm{Gal}(\iota_{C'}) \cong S_{d/2}$  is surjective;
- $\mathrm{Gal}(\iota_{C'} \circ h)$  contains a transposition of a pair of points in the fiber of  $h$ .

□

**Lemma 2.3.** *Let  $G$  be a subgroup of  $S_2^{\oplus m} \rtimes S_m$ , acting naturally on the set  $\pm 1, \dots, \pm m$ . If the projection  $G \rightarrow S_m$  is surjective and the transposition  $\sigma_0$  of  $\pm 1$  lies in  $G$ , then  $G = S_2^{\oplus m} \rtimes S_m$ .*

*Sketch of proof.* (with help from Chenji Fu) Let  $\varepsilon_i$  denote the transposition of  $\pm i$ . For any  $\sigma \in S_m$ , choose a lift  $\tilde{\sigma} \in G$ , then

$$\varepsilon_{\sigma(1)} = \tilde{\sigma} \circ \sigma_0 \circ \tilde{\sigma}^{-1} \in G.$$

Thus,  $S_2^{\oplus m} \subset G$ , and since  $G$  maps onto  $S_m$ , we obtain  $G = S_2^{\oplus m} \rtimes S_m$ . □

**Example 2.4.** *Let  $C$  be a smooth curve of genus  $g$  embedded in its Jacobian  $A := \mathrm{Jac}(C)$  via the Abel–Jacobi map  $\mathrm{AJ}_C : C \hookrightarrow A$ .*

*When  $C$  is non-hyperelliptic, the corresponding Gauss map*

$$|\omega_C| : C \rightarrow \mathbb{P}^{g-1}$$

*makes  $C$  as an irreducible nondegenerate curve of degree  $2g - 2$ , by Proposition 2.1 we get*

$$\mathrm{Gal}(\gamma_C) \cong S_{2g-2}.$$

*When  $C$  is hyperelliptic, the corresponding Gauss map is  $2 : 1$  onto a rational normal curve  $R \subset \mathbb{P}^{g-1}$ :*

$$|\omega_C| : C \xrightarrow{2:1} R \hookrightarrow \mathbb{P}^{g-1}$$

*By Proposition 2.2 we get*

$$\mathrm{Gal}(\gamma_C) \cong S_2^{\oplus g-1} \rtimes S_{g-1}.$$

In fact, a degree  $2 : 1$  map does not give rise to any exceptional monodromy groups beyond those listed in Table 1.

**Definition 2.5** (big monodromy group). *We refer to the big monodromy group as any group of the following types:*

<i>notation</i>	<i>name</i>	<i>alias</i>
$W(A_{m+1}) = S_m$	<i>full symmetric group</i>	
$W(C_m) = S_2^{\oplus m} \rtimes S_m$	<i>signed symmetric group</i>	<i>hyperoctahedral group</i>
$W(D_m) = (S_2^{\oplus m})_0 \rtimes S_m$	<i>even-signed symmetric group</i>	<i>demihyperoctahedral group</i>

TABLE 1. big monodromy group

**Proposition 2.6.** *Let  $\iota_{C'} : C' \hookrightarrow \mathbb{P}^{n-1}$  be an irreducible nondegenerate curve of degree  $d/2$ , and let  $h : C \rightarrow C'$  be a degree 2 ramified covering. Then*

$$\mathrm{Gal}(\iota_{C'} \circ h) \cong W(C_{d/2}) \text{ or } W(D_{d/2}).$$

*Sketch of proof.* By Lemma 1.4 we know that  $\mathrm{Gal}(\iota_{C'} \circ h) \subseteq W(D_{d/2})$ . By Lemma 2.7, we are reduced to verifying that:

- The quotient map  $\mathrm{Gal}(\iota_{C'} \circ h) \rightarrow \mathrm{Gal}(\iota_{C'}) \cong S_{d/2}$  is surjective;
- (signed doubly transitive)  $\mathrm{Gal}(\iota_{C'} \circ h)$  acts transitively on pairs  $(x, y)$  with  $x \neq \pm y$ .

□

**Lemma 2.7.** *Let  $G$  be a subgroup of  $W(D_m)$ , acting naturally on the set  $\pm 1, \dots, \pm m$ . If the projection  $G \rightarrow S_m$  is surjective then*

$$G \cong W(C_m) \text{ or } W(D_m) \text{ or } S_m.$$

*Sketch of proof.* (with help from Chenji Fu) Let  $H$  denote the kernel of the natural quotient map  $G \rightarrow S_m$ . Then  $H$  is stable under the action of  $S_n$ . There are only three possible forms that  $H$  can take:

- $H = 0$ . Then  $G \cong S_m$ .
- $H = (S_2^{\oplus m})_0$ . Then  $G$  is a index 2 subgroup of  $W(C_m)$ , so  $G \cong W(D_m)$ .<sup>5</sup>
- $H = S_2^{\oplus m}$ . Then  $G = W(C_m)$ .

□

We are particularly interested in identifying cases where the monodromy group is small(i.e., not big), which can occur only when  $k > 2$ , where  $k$  denotes the degree of the map  $\phi : C \rightarrow \mathbb{P}^{n-1}$ . On the other hand, when both  $k > 2$  and  $n > 2$ , Lemma 1.4 ensures that the resulting monodromy group is indeed small. Thus, in the case of curves, Question 1.2 is resolved except in the following three cases:

- (1)  $n > 2$  and  $k = 2$ . By Proposition 2.6, the monodromy group is known to be large, although its precise structure remains undetermined.
- (2)  $n = 2$ . Here,  $A$  is an abelian surface and  $C$  corresponds to a divisor.
- (3)  $n > 2$  and  $k > 2$ . Finding an example of this would guarantee a case of small monodromy.

**Question 2.8.** *Can we find any curve  $C \subset A$  not stable under any translation on  $A$ , and for which the associated monodromy group is small?*

*Assume that  $n > 2$ . Can we find any curve  $C \subset A$  not stable under any translation on  $A$ , whose Gauss map has degree  $d > 2$ ?*

<sup>5</sup>Check [stackexchange discussions](#)

**2.2. Curves in Prym variety.** To address Question 2.8, we seek explicit examples, and the case of a Prym curve mapping to its associated Prym variety offers a natural starting point for investigation.

**Notations 2.9.** Suppose  $C'$  is a smooth projective curve of genus  $g(C')$ , and let  $B$  be an effective (possibly zero) divisor on  $C'$ . For any line bundle  $\eta \in \text{Pic}(C')$  such that  $\eta^{\otimes 2} \cong \mathcal{O}_{C'}(B)$ , one obtains a double cover  $h : C \rightarrow C'$  of smooth projective curves, ramified precisely over  $B$ . The associated involution on  $C$  is denoted by  $\iota$ , and the triple  $(C, C', h)$  is referred to as the Prym pair.

Recall that the Prym variety  $A := \text{Prym}(C/C')$  is defined as the connected component of the identity in

$$\ker [\text{Nm} : \text{Jac}(C) \rightarrow \text{Jac}(C')].$$

The Abel–Prym map is defined by

$$\text{AP}_{C/C'} : C \rightarrow A \quad p \mapsto \mathcal{O}_C(p - \iota(p)).$$

The classical theory treats the case where the Prym variety  $A$  is principally polarized—this occurs exactly when  $B = \emptyset$  or  $B = \{p_0, q_0\}$ , that is, when  $h$  is either unramified or ramified at two points.<sup>6</sup> For clarity and focus, we restrict our attention to these cases. Table 2 lists the relevant numerical data.<sup>7</sup>

	$g(C)$	$g(C')$	$\deg \eta$
$B = \emptyset$	$2n + 1$	$n + 1$	0
$B = \{p_0, q_0\}$	$2n$	$n$	1

TABLE 2. numerical data of Prym pair

The Abel–Prym map  $\text{AP}_{C/C'}$  does not always behave as nicely as the Abel–Jacobi map; it may fail to be an embedding. When  $C$  is hyperelliptic, the map  $\text{AP}_{C/C'}$  fails to be generically injective, and therefore falls outside the scope of our discussion. When  $C$  is non-hyperelliptic, we may regard it as a subvariety of  $A$  for our purposes, although it may be not strictly embedded.<sup>8</sup> However, it remains unclear whether  $C$  is stable under any translation in  $A$ .

In the Prym setting, the corresponding Gauss map  $\gamma_C$  factors through  $h$ :

$$\begin{aligned} \gamma_C : C &\xrightarrow{h} C' \xrightarrow{|\omega_{C'} \otimes \eta|} \mathbb{P}^{n-1} = \mathbb{P}(\text{H}^0(\omega_{C'} \otimes \eta)^*) \\ &\cong \text{Gr}(n-1, \text{H}^0(\omega_{C'} \otimes \eta)) \\ \tilde{p} &\longmapsto p \longmapsto \text{H}^0(\omega_{C'} \otimes \eta(-p)) \end{aligned}$$

Unfortunately, in the Prym setting, no example has been found for Question 2.8, and such an example is likely to exist only when the gonality  $\text{gon}(C)$  is sufficiently small.

**Lemma 2.10.** *When  $\text{gon}(C) > 4$ ,  $|\omega_{C'} \otimes \eta|$  is injective. As a result, the monodromy group is big.*

*Proof.* Suppose that  $|\omega_{C'} \otimes \eta|$  is not injective, we need to find a line bundle of degree 4 and rank  $\geq 1$ . In fact, for  $p \neq q$ ,

$$\begin{aligned} |\omega_{C'} \otimes \eta|(p) &= |\omega_{C'} \otimes \eta|(q) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-p - q)) &= 1 && \text{By [10, 19.2.8]} \\ \iff h^0(\omega_{C'} \otimes \eta(-p - q)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^{\vee}(p + q)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

<sup>6</sup>See [3, Theorem 3.2.6] for the proof.

<sup>7</sup>As usual,  $n = \dim_{\mathbb{C}} A$  is the dimension of the abelian variety.

<sup>8</sup>For a detailed description of the map  $\text{AP}_{C/C'}$ , see [2, Proposition 12.5.2, Corollary 12.5.6]. Although  $\text{AP}_{C/C'}$  may collapse  $p_0$  and  $q_0$ , this does not affect the relevant computation of  $\text{Gal}(\gamma_Z)$ .

When  $B = \emptyset$ , write  $\eta^\vee(p + q) = \mathcal{O}_{C'}(p' + q')$ , then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^\vee)^{\otimes 2}(2p + 2q) = \mathcal{O}_{C'}(2p' + 2q') \in g_4^1;$$

When  $B = \{p_0, q_0\}$ , write  $\eta^\vee(p + q) = \mathcal{O}_{C'}(p')$ , then

$$\mathcal{O}_{C'}(2p + 2q) = (\eta^\vee)^{\otimes 2}(2p + 2q + p_0 + q_0) = \mathcal{O}_{C'}(2p' + p_0 + q_0) \in g_4^1.$$

□

*Remark 2.11.* Based on the strategy used in the proof of Lemma 2.10, we can in fact obtain stronger results. For  $p \in C$ ,

$$\begin{aligned} |\omega_{C'} \otimes \eta| &\text{ is ramified at } p && \text{(i.e., the tangent map is 0 at } p) \\ \iff h^0(\omega_{C'} \otimes \eta) - h^0(\omega_{C'} \otimes \eta(-2p)) &= 1 && \text{By [10, 19.2.9]} \\ \iff h^0(\omega_{C'} \otimes \eta(-2p)) &= n - 1 && \text{Since } \dim_{\mathbb{C}} A = n \\ \iff h^0(\eta^\vee(2p)) &= 1 && \text{Via Riemann–Roch} \end{aligned}$$

For the remainder of this discussion, we focus exclusively on the case  $B = \emptyset$ . Combining both,

$$\begin{aligned} &|\omega_{C'} \otimes \eta| \text{ is not generically injective} \\ \iff &\text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } h^0(\eta^\vee(p + q)) = 1 \\ \iff &\text{For any } p \in C', \text{ there exist } q \in C' \text{ such that } \eta^\vee + p \in C' + C' - q \\ \iff &\eta^\vee + C' \subseteq C' + C' - C'. \end{aligned}$$

It is expected that

$$\eta^\vee + C' \subseteq C' + C' - C' \iff \eta^\vee \in C' - C'. \quad (2.1)$$

If so, writing  $\eta^\vee = \mathcal{O}_{C'}(p - q)$ , we obtain

$$\mathcal{O}_{C'}(2p) = \mathcal{O}_{C'}(2q) \in g_2^1,$$

which implies that  $C'$  is hyperelliptic. By Fact 2.12, this forces  $C$  to be hyperelliptic as well, and hence it lies outside the scope of our discussion.

Unfortunately, (2.1) does not hold in general. A counterexample, communicated to me by Gavril Farkas, will be presented in Proposition 2.14.

**Fact 2.12** (From [9, p346]). *We have an explicit classification of  $2 : 1$  unramified covers of a hyperelliptic curve  $C'$ . Let  $\pi : C' \rightarrow \mathbb{P}^1$  be the double cover ramified at  $\{p_1, \dots, p_{2g+2}\}$ , then every such unramified double cover of  $C'$  arises from some line bundle in  $C$  of the form*

$$\eta := \mathcal{O}_{C'}(p_{i_1} + \dots + p_{i_l} - p_{i_{l+1}} - \dots - p_{i_{2l}}) = \mathcal{O}_{C'}\left(\sum_{j=1}^{2l} p_{i_j} - 2l \cdot p_1\right).$$

When  $l = 1$ , the associated covering curve  $C$  is also hyperelliptic.

**Question 2.13.** *Can one find a non-hyperelliptic curve  $C'$  and a nontrivial line bundle  $\eta \in \text{Jac}(C')$  satisfying  $\eta^{\otimes 2} \cong \mathcal{O}_{C'}$ , such that*

$$\eta^\vee + C' \subseteq C' + C' - C'?$$

*If such a pair exists, can we find a case where the resulting curve  $C \subset A$  is not preserved under any translation on  $A$ ?*

As a key example relevant to Question 2.13, I include here a result of Gavril Farkas.

**Proposition 2.14** (See [4, Proposition 2.8]). *Let  $C$  be a general smooth curve of genus  $g = 2i$ , and suppose  $D \in W_{i+1}^1(C)$  is a pencil of minimal degree—i.e., a line bundle  $\mathcal{O}_C(D) \in \text{Pic}^{i+1}(C)$  with  $h^0(C, \mathcal{O}_C(D)) = 2$ . Define*

$$\mathcal{L} := K_C - 2D \in \text{Pic}^{2i-4}(C),$$

then  $\mathcal{L}$  satisfies

$$\mathcal{L} + C \subset C_{2i-2} - C, \quad \text{but} \quad \mathcal{L} \notin C_{2i-3} - C.$$

*Remark 2.15.* Take  $i = 2$ , the line bundle  $\mathcal{L}$  provides a counterexample to (2.1). However, this example does not directly yield a counterexample to Question 2.13. This is due to the uncertainty of whether there exist  $C$  and  $D$  satisfying  $\mathcal{L}^{\otimes 2} \cong \mathcal{O}_C$ , and whether the curve  $C$  is sufficiently general for Proposition 2.14 to hold.

I intend to revisit the proof in detail, especially the use of the base point free pencil trick and the Petri map. Should this yield any useful observations, they will be included here.

**2.3. Albanese variety.** Let  $X$  be a smooth complex projective variety of dimension  $r$ , and set  $n := \dim_{\mathbb{C}} H^0(X, \Omega_X) = h^{1,0}$ . Recall that the Albanese variety of  $X$  is defined as

$$\text{Alb}(X) := H^0(X, \Omega_X)^* / H_1(X, \mathbb{Z})_{\text{free}},$$

and the Albanese map is given by (pick a base point  $p_0 \in X$ )

$$\alpha : X \longrightarrow \text{Alb}(X) \quad p \longmapsto \left[ \omega \mapsto \int_{\gamma: p_0 \sim p} \omega \right].$$

One classical question is the dimension of  $\alpha(X)$ . For this, consider the cotangent map of  $\alpha$  at  $p \in X$ :

$$T_p^* \alpha : H^0(X, \Omega_X) \longrightarrow T_p^* X = H^0(X, \Omega_X|_p).$$

Consider the short exact sequence of coherent sheaves on  $X$

$$0 \longrightarrow \Omega_X(-p) \longrightarrow \Omega_X \longrightarrow \Omega_X|_p \longrightarrow 0$$

which induces a long exact sequence

$$\begin{array}{ccccccc} & & H^1(X, \Omega_X(-p)) & \longrightarrow & H^1(X, \Omega_X) & \longrightarrow & 0 \\ & \nearrow & & & & & \\ 0 & \longrightarrow & H^0(X, \Omega_X(-p)) & \longrightarrow & H^0(X, \Omega_X) & \xrightarrow{T_p^* \alpha} & H^0(X, \Omega_X|_p) \end{array}$$

The proposition below follows from standard arguments in homological algebra:

**Proposition 2.16.** *For a general point  $p \in X$ ,*

$$\begin{aligned} \dim_{\mathbb{C}} \alpha(X) &= \text{rank } T_p^* \alpha \\ &= n - h^0(X, \Omega_X(-p)) \\ &= h^1(X, \Omega_X(-p)) - h^1(X, \Omega_X). \end{aligned}$$

*In particular,*

$$\begin{array}{ll} \alpha \text{ is surjective} & \iff h^0(X, \Omega_X(-p)) = 0 \\ \alpha \text{ is finite onto image} & \iff h^0(X, \Omega_X(-p)) = n - r \\ \alpha \text{ is constant} & \iff h^0(X, \Omega_X(-p)) = n \\ & \iff h^1(X, \Omega_X(-p)) = h^1(X, \Omega_X) \\ & \iff n = 0. \end{array}$$

We will concentrate on the case where  $\alpha$  is finite onto its image.<sup>9</sup> Under this assumption, we set  $Z = X$  and  $A = \text{Alb}(Z)$ . The corresponding Gauss map is then a rational map:

$$\begin{array}{ccc} \phi_Z : Z & \dashrightarrow & \text{Gr}(r, T^0 A) \cong \text{Gr}(n - r, H^0(X, \Omega_X)) \\ p & \longmapsto & H^0(X, \Omega_X(-p)) \end{array}$$

<sup>9</sup>The general method remains valid in the broader setting, but the Gauss map then takes values in a different space.



and we have the isomorphisms

$$\begin{aligned}\mathbb{P}T_0^*A &\cong \mathbb{P}H^0(X, \Omega_X), \\ \mathbb{P}\Lambda_Z &= \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^*A \mid \omega(p) = 0 \} \\ &\cong \{ (p, [\omega]) \in Z \times \mathbb{P}T_0^*A \mid \omega \in H^0(X, \Omega_X(-p)) \} \\ &\cong (\phi_Z, \text{Id})^{-1} I_{n-r,1},\end{aligned}$$

where

$$I_{n-r,1} := \{ (V, [\omega]) \in \text{Gr}(n-r, n) \times \text{Gr}(1, n) \mid \omega \in V \}$$

is the incidence variety relating  $\text{Gr}(n-r, n)$  and  $\text{Gr}(1, n)$ . In that case,

$$\gamma_Z^{-1}([\omega]) = \{ p \in X \mid \omega(p) = 0 \}$$

is the zero set of section  $\omega \in H^0(X, \Omega_X)$ . The number  $(-1)^r \deg \gamma_Z$  is the index in the Poincaré–Hopf index formula, and the monodromy group  $\text{Gal}(\gamma_Z)$  serves as a more refined invariant, encoding subtler aspects of the geometry.

*Remark 2.17.* One can define the monodromy group in the real (or topological) setting in an analogous manner. Consider the set

$$U := \{ \omega \in \Gamma(\Omega_M) \mid \text{the zeros of } \omega \text{ are isolated with local index } 1 \}.$$

A loop in  $U$  induces a permutation of the zeros of  $\omega$ , and the monodromy group is defined as the group of all such permutations.

However, when  $\dim_{\mathbb{C}} M > 1$ , the corresponding monodromy group is always the full symmetric group (as all transpositions occur). As a result, it does not yield any further insight into the topology of the manifold.

The next proposition shows when the Gauss map  $\phi_Z$  is not generic injective.

**Proposition 2.18.** *When  $\alpha$  is finite onto its image,*

*$\phi_Z$  is not generic injective*

$$\iff \text{For general } p \in X, \text{ exists } q \neq p \text{ such that } h^0(X, \Omega_X(-p-q)) = n-r$$

$$\iff \text{For general } p \in X, \text{ exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r$$

$$\iff \text{For all } p \in X, \text{ exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) = n-r.$$

This last step relies on the following lemma, combined with the closedness of the tautological correspondence in  $X \times X^{[2]}$ .

**Lemma 2.19.** *Let  $m \in \mathbb{Z}_{>0}$ . The function*

$$h^\Omega : X^{[m]} \longrightarrow \mathbb{Z}_{\geq 0} \quad S \longmapsto h^0(X, \Omega_X \otimes \mathcal{I}_S)$$

*is Zariski upper semicontinuous.*

*Sketch of proof.* Consider the coherent sheaf  $\mathcal{F} \in \text{Coh}(X^{[m]} \times X)$  characterized by the property

$$\mathcal{F}|_{\{S\} \times X} \cong \Omega_X \otimes \mathcal{I}_S,$$

The proposition then follows directly from the semicontinuity theorem [10, 28.1.1].  $\square$

The stratification of  $X^{[m]}$  by  $h^\Omega$  offers a natural generalization of Brill–Noether theory beyond the setting of curves.

At the end of this subsection, let us turn to the setting of a general abelian variety  $A$  and a smooth subvariety  $\iota_Z : Z \hookrightarrow A$ , where  $n = \dim_{\mathbb{C}} A$  and  $r = \dim_{\mathbb{C}} Z$ . Observe that  $\iota_Z$  factors through the Albanese variety of  $Z$ :

$$\iota_Z : Z \xrightarrow{\alpha_Z} \text{Alb}(Z) \xrightarrow{\pi} A$$

We shall also assume that  $Z$  generates  $A$ ; it then follows that the map  $\pi$  is surjective. The cotangent map of  $\iota_Z$  at a point  $p \in Z$  factors through  $H^0(Z, \Omega_Z)$ :

$$T_p^* \iota_Z : T_p^* A \hookrightarrow H^0(Z, \Omega_Z) \longrightarrow T_p^* Z$$

For convenience, abbreviate  $V := T_0^* A \cong T_p^* A$ , and view  $V$  as a subspace of  $H^0(Z, \Omega_Z)$ .

**Proposition 2.20.** *Assume that  $Z$  is embedded in  $A$  and generates  $A$ , and let  $V := T_0^* A$ . Then*

$$\dim_{\mathbb{C}} H^0(Z, \Omega_Z(-p)) \cap V = n - r \quad \text{for all } p \in Z$$

*It follows that the Gauss map is a regular morphism*

$$\begin{array}{ccc} \phi_Z : Z & \longrightarrow & \text{Gr}(r, T_0^* A) \cong \text{Gr}(n - r, V) \\ p & \longmapsto & H^0(Z, \Omega_Z(-p)) \cap V \end{array}$$

Furthermore,

$\phi_Z$  is not generic injective

$$\iff \text{For general } p \in X, \text{ exists } q \neq p \text{ such that } h^0(X, \Omega_X(-p - q)) \cap V = n - r$$

$$\iff \text{For all } p \in X, \text{ exists } S \in X^{[2]} \text{ such that } p \in S \text{ and } h^0(X, \Omega_X \otimes \mathcal{I}_S) \cap V = n - r.$$

**Question 2.21.** *For an  $r$ -dimensional smooth variety  $Z$  whose Albanese map  $\alpha_Z$  is an embedding, which subspaces  $V \subset H^0(Z, \Omega_Z)$  can arise as the cotangent space of some quotient variety of  $\text{Alb}(Z)$ ?*

The precise formulation of this question is somewhat ambiguous, and depending on the interpretation, it may turn out to be either relatively straightforward or extremely difficult. On the one hand, the classification of abelian subvarieties of  $A$  via symmetric idempotents in  $\text{End}_{\mathbb{Q}}(A)$  is well established [2, Theorem 5.3.2]. On the other hand, explicitly computing  $\text{End}_{\mathbb{Q}}(A)$  can be highly nontrivial when  $A$  is a non-simple Albanese variety—even in the case where  $A$  arises from a curve.

### 3. FAMILIES OF SUBVARIETIES

In this section, we move from the study of monodromy groups to a more direct analysis of the subvarieties themselves. Given an initial subvariety, one can naturally generate a family of subvarieties. Our goal here is to define these families and investigate their properties.

**Proposition 3.1.** *All irreducible conic Lagrangian cycles in  $T^*A$  are of the form  $\Lambda_Z$  for some irreducible subvariety  $Z \subset A$ . This yields a one-to-one correspondence between irreducible conic Lagrangian cycles in  $T^*A$  and irreducible subvarieties of  $A$ :*

$$\{ \text{irreducible conic Lagrangian cycles in } T^*A \} \longleftrightarrow \{ \text{irreducible subvarieties in } A \}$$

*Sketch of proof.* For any irreducible conic Lagrangian cycle  $\Lambda \subset T^*A$ , let  $Z$  denote the image of  $\Lambda$  under the natural projection  $T^*A \rightarrow A$ . Our goal is to show that  $\Lambda = \Lambda_Z$ .

- By definition,  $\Lambda \subset T^*A|_Z$ .
- Since  $\Lambda$  is conic, we have  $s(Z) \subset \Lambda$ , where  $s : A \rightarrow T^*A$  denotes the zero section.
- Since  $\Lambda$  is Lagrangian and  $s(Z) \subset \Lambda$ , we have  $\Lambda \subset \Lambda_Z$ .
- Since  $\Lambda_Z$  is irreducible with  $\dim_{\mathbb{C}} \Lambda = \dim_{\mathbb{C}} \Lambda_Z = n$ , we have  $\Lambda = \Lambda_Z$ .

□

Why do we shift attention from  $Z$  to  $\Lambda_Z$  as the main object of study? One reason is the uniformity of  $\Lambda_Z$ : it always has dimension  $n$ , and in most cases, the natural map  $\Lambda_Z \rightarrow T_0^*A$  is generically finite, with fibers lying inside  $A$ .

**Definition 3.2** (Clean cycle). *An irreducible Lagrangian cycle  $\Lambda \subset T^*A$  is called clean if the composed projection*

$$\Lambda \longrightarrow T^*A \twoheadrightarrow T_0^*A$$

*is generically finite.*

Another important reason is that the space of weighted clean conic Lagrangian cycles naturally acquires a convolution structure, arising from the group law on  $A$ , which plays a central role in the analysis.

**Proposition 3.3.** *The group of weighted clean conic Lagrangian cycles*

$$\begin{aligned} \mathcal{L}(A) &:= \{ \text{weighted clean conic Lagrangian cycles in } T^*A \} \\ &= \left\{ \sum_{\substack{Z_i \subset A \\ \text{irr clean}}} n_i \Lambda_{Z_i} \mid n_i \in \mathbb{Z} \right\} \end{aligned}$$

has a natural convolution structure as follows:

$$\begin{aligned} \Lambda_{Z_1} \circ \Lambda_{Z_2} &= \text{the clean part of } (a, \text{Id}_{T_0^*A})_* (\Lambda_{Z_1} \times_{T_0^*A} \Lambda_{Z_2}) \\ &= \overline{(a, \text{Id}_U)_* (\Lambda_{Z_1}|_U \times_U \Lambda_{Z_2}|_U)} \end{aligned}$$

where

$$U := \left\{ \xi \in T_0^*A \mid \deg \phi_{Z_i} = \# \phi_{Z_i}^{-1}(\xi) \text{ for } i = 1, 2 \right\}$$

and  $a : A \times A \longrightarrow A$  is the addition map in  $A$ . The general convolution is defined by  $\mathbb{Z}$ -linear extension.

*Sketch of proof.* To establish the claim, it suffices to show that  $\Lambda_{Z_1} \circ \Lambda_{Z_2}$  defines a weighted conic Lagrangian cycle. The conic property follows directly from the definition, while the Lagrangian condition can be verified at a general point  $(p_1 + p_2, \xi) \in \Lambda_{Z_1} \circ \Lambda_{Z_2}$ .  $\square$

We now consider the projective versions of all objects involved, so that we may make use of properness.

**Lemma 3.4.**

- (1) Suppose that  $\mathbb{P}\Lambda_{Z_1}, \mathbb{P}\Lambda_{Z_2} \subset \mathbb{P}T^*A$  admit monodromy representations

$$\rho_{\gamma_{Z_i}} : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_i}^{-1}(\xi_0)),$$

then  $\mathbb{P}\Lambda_{Z_1} \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_{Z_2} \subset A \times A \times \mathbb{P}^\vee$  admits monodromy representation given by

$$(\rho_{\gamma_{Z_1}}, \rho_{\gamma_{Z_2}}) : \pi_1(U, \xi_0) \longrightarrow \text{Aut}(\gamma_{Z_1}^{-1}(\xi_0) \times \gamma_{Z_2}^{-1}(\xi_0)),$$

- (2) When  $Z_1 = Z_2 = Z$ , we obtain a one-to-one correspondence:

$$\left\{ \begin{array}{c} \text{irr components of } \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \end{array} \right\}$$

*Sketch of proof.* Statement (1) holds by definition. The proof of (2) reduces to the following purely topological statement:

**Claim 3.5.** *Let  $\pi : E \longrightarrow B$  be a (unramified) covering space over a manifold  $B$  with deck transformation group  $G$ , then*

$$\{ \text{connected components of } E \times_B E \} \longleftrightarrow \{ G\text{-orbits of } \pi^{-1}(b_0) \times \pi^{-1}(b_0) \}.$$

The claim follows directly from the correspondence between covering spaces over  $B$  and  $\pi_1(B)$ -sets; see [5, Theorem 1.38].  $\square$

Generalizing the argument of Lemma 3.4, we arrive at the following lemma.

**Lemma 3.6.** *For  $d = \deg \gamma_Z$ ,  $\xi_0 \in T_0^*A$  a general point, write*

$$\begin{aligned} \mathbb{P}\Lambda_Z^{\times d} &:= \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \times_{\mathbb{P}^\vee} \cdots \times_{\mathbb{P}^\vee} \mathbb{P}\Lambda_Z \subset A^d \times \mathbb{P}^\vee \\ \gamma_Z^{-1}(\xi_0)^d &:= \gamma_Z^{-1}(\xi_0) \times \gamma_Z^{-1}(\xi_0) \times \cdots \times \gamma_Z^{-1}(\xi_0) \subset A^d \end{aligned}$$

(1) we obtain an one-to-one correspondence:

$$\left\{ \begin{array}{l} \text{irr components of } \mathbb{P}\Lambda_Z^{\times d} \\ \text{with a surjection to } \mathbb{P}^\vee \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{Gal}(\gamma_Z)\text{-orbits of} \\ \gamma_Z^{-1}(\xi_0)^d \end{array} \right\}$$

(2) Write

$$\begin{aligned} \Delta_d &:= \{(p_1, \dots, p_d) \in A^d \mid p_i = p_j \text{ for some } i \neq j\} \subset A^d \\ \mathbb{P}\Lambda_Z^{[d]} &:= \overline{(\mathbb{P}\Lambda_Z^{\times d} \setminus (\Delta_d \times \mathbb{P}^\vee))} \big|_U \subset A^d \times \mathbb{P}^\vee \end{aligned}$$

Fix a general point  $\xi_0 \in \mathbb{P}^\vee$  and a well-order for  $\gamma_Z^{-1}(\xi_0)$ , one can identify  $S_d \cong \gamma_Z^{-1}(\xi_0)^d \setminus \Delta_d$ , and

$$\left\{ \text{irr components of } \mathbb{P}\Lambda_Z^{[d]} \right\} \longleftrightarrow \left\{ \text{Gal}(\gamma_Z)\text{-orbits of } S_d \right\}$$

In reference,  $\Delta_d$  is usually called the big diagonal.

From this point on, we fix an irreducible component of  $\mathbb{P}\Lambda_Z^{[d]}$ , denoted by  $\mathbb{P}\Lambda_Z^{\text{univ}}$ . As we will see in Definition A, this variety generates all subvarieties within the families under consideration.

**Definition 3.7** (The subvariety  $Z^{(m)}$ ). *For any tuple  $(m) = (m_1, \dots, m_d) \in \mathbb{Z}^d$ , we define the weighted sum map*

$$a^{(m)} : A^d \longrightarrow A \quad (p_1, \dots, p_d) \longmapsto \sum_{i=1}^d m_i p_i.$$

We also define

$$\mathbb{P}\Lambda_Z^{(m)} := \left( a^{(m)}, \text{Id}_{\mathbb{P}^\vee} \right)_* \mathbb{P}\Lambda_Z^{\text{univ}}.$$

as the (projectivized) weighted Lagrangian cycle in  $\mathbb{P}T^*A$ . The projective cycle  $\mathbb{P}\Lambda_Z^{(m)}$  is irreducible but may appear with multiplicities. We can therefore write

$$\mathbb{P}\Lambda_Z^{(m)} = c_Z^{(m)} \mathbb{P}\Lambda_{Z^{(m)}}$$

where  $c_Z^{(m)} \in \mathbb{Z}_{>0}$  and  $Z^{(m)} \subset A$  are uniquely determined. This gives rise to a family of subvarieties parametrized by  $\mathbb{Z}^d$ .

The next lemma gathers some basic properties of  $Z^{(m)}$ . Observe that  $S_d = \text{Aut}(\gamma_Z^{-1}(\xi_0))$  acts naturally on  $\mathbb{Z}^d$  via

$$g(m) = (m_{g(1)}, \dots, m_{g(d)}) \in \mathbb{Z}^d.$$

**Lemma 3.8.**

- (1) For all  $g \in \text{Gal}(\gamma_Z)$ , we have  $Z^{g(m)} = Z^{(m)}$ ,  $c_Z^{g(m)} = c_Z^{(m)}$ ;
- (2) For  $(m) = (1, 0, \dots, 0) \in \mathbb{Z}^d$ ,  $Z^{(m)} = Z$ ;
- (3) For all  $(m), (m') \in \mathbb{Z}^d$ , we have  $\mathbb{P}\Lambda_Z^{(m)} \circ \mathbb{P}\Lambda_Z^{(m')} = \mathbb{P}\Lambda_Z^{(m+m')}$ ;
- (4) The group  $\langle \mathbb{P}\Lambda_{Z^{(m)}} \rangle_{\text{Abel}}$  is closed under the convolution product.

**Question 3.9.** *Can we compute the dimension and the homology class for these subvarieties  $Z^{(m)}$ ?*

At first glance, this question may appear naive, but it turns out to be extremely challenging. The main difficulty lies in determining the homology class of  $\mathbb{P}\Lambda_Z^{\text{univ}}$ . Once this is known, one can compute the pushforward to obtain the homology class of  $\mathbb{P}\Lambda_Z^{(m)}$ , from which both  $\dim_{\mathbb{C}} Z^{(m)}$  and the homology class of  $Z^{(m)}$  (up to a scalar factor  $c_Z^{(m)}$ ) can be extracted.

However, standard intersection theory yields only the homology class of  $\mathbb{P}\Lambda_Z^{[d]}$ , via the Chern–Mather class of  $Z$ . At present, there is no known method for computing the homology class of a single irreducible component of  $\mathbb{P}\Lambda_Z^{[d]}$ . Although we can distinguish these components topologically, we lack tools to compute their topological invariants, which presents an unfortunate limitation.

Nevertheless, Question 3.9 can still be addressed in certain special situations. For instance, when  $\text{Gal}(\gamma_Z) = S_n$ , the space  $\mathbb{P}\Lambda_Z^{[d]}$  is known to be irreducible, enabling explicit computation. Similarly, when  $Z = -Z$  and  $\text{Gal}(\gamma_Z) = W(C_{d/2})$ , one can impose additional conditions (e.g.,  $p_{2l-1} = -p_{2l}$ ) to guarantee the uniqueness of  $\mathbb{P}\Lambda_Z^{\text{univ}}$ , which again allows us to perform the computation.

**Question 3.10.** *Suppose  $d = 2d_0$  with  $d_0$  even,  $(m) \in \mathbb{Z}^d$ , and  $g \in W(C_{d/2}) \setminus W(D_{d/2})$ . Is it possible to construct an example with  $Z = -Z$  and  $\text{Gal}(\gamma_Z) = W(D_{d/2})$ , yet  $\dim_{\mathbb{C}} Z^{g(m)} \neq \dim_{\mathbb{C}} Z^{(m)}$ ?*

**Question 3.11.** *Suppose that  $Z$  is the Fano surface of a smooth cubic threefold, and let  $A := \text{Alb}(Z)$ . How can we compute the homology class of  $Z^{(m)}$ ?*

#### 4. TANNAKIAN FORMALISM

For simplicity, we work over the base field  $\kappa = \mathbb{C}$ . Let  $A$  denote a fixed complex abelian variety, and let  $\text{Perv}(A)$  denote the category of perverse sheaves on  $A$  with coefficients in  $\mathbb{Q}$ . For any algebraic group  $G$ , we denote by  $\text{Rep}(G)$  the category of algebraic representations of  $G$ .

Following the approach of [8], we work in the quotient category  $\overline{\text{Perv}}(A) = \text{Perv}(A)/N(A)$ , where  $N(A) \subset \text{Perv}(A)$  is the Serre subcategory of negligible complexes. A complex  $\mathcal{F}$  is defined to be negligible if  $\chi(A, \mathcal{F}) = 0$ . This quotient category admits a natural convolution structure, and every finitely generated tensor subcategory of it is Tannakian, with a reductive Tannaka group  $G$  (see [8, Thm 7.1 & Cor 9.2]). In particular, for any perverse sheaf  $\delta \in \overline{\text{Perv}}(A)$ , the full subcategory generated by  $\delta$  is categorically equivalent to the representation category of an algebraic group  $G$ :

$$\langle \delta, * \rangle \cong \text{Rep}(G).$$

#### REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Volume I*, volume 267 of *Grundlehren Math. Wiss.* Springer, Cham, 1985.
- [2] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 302 of *Grundlehren Math. Wiss.* Berlin: Springer, 2nd augmented ed. edition, 2004.
- [3] Giulio Codogni and Thomas Krämer. Semicontinuity of Gauss maps and the Schottky problem. *Math. Ann.*, 382(1-2):607–630, 2022.
- [4] Gavril Farkas and Michael Kemeny. The generic Green-Lazarsfeld secant conjecture. *Invent. Math.*, 203(1):265–301, 2016.
- [5] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge University Press, 2002.
- [6] Ariyan Javanpeykar, Thomas Krämer, Christian Lehn, and Marco Maculan. The monodromy of families of subvarieties on abelian varieties. Preprint, arXiv:2210.05166 [math.AG] (2022), 2022.
- [7] Thomas Krämer. Cubic threefolds, Fano surfaces and the monodromy of the Gauss map. *Manuscr. Math.*, 149(3-4):303–314, 2016.
- [8] Thomas Krämer and Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties. *J. Algebr. Geom.*, 24(3):531–568, 2015.
- [9] David Mumford. *The red book of varieties and schemes. Includes the Michigan lectures (1974) on “Curves and their Jacobians”*, volume 1358 of *Lect. Notes Math.* Berlin: Springer, 2nd, expanded ed. with contributions by Enrico Arbarello edition, 1999.
- [10] Ravi Vakil. The rising sea: Foundations of algebraic geometry. *preprint*, 2017.

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, BERLIN, 12489, GERMANY,  
*Email address:* email:xiaoxiang.zhou@hu-berlin.de