

## Triangle Inequality

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

$$|z_2 + z_1| \geq |z_2| - |z_1|$$

$$f'(z) = \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z}$$

If  $f$  is differentiable in an open region we say  $f$  is analytic in that region

## Cauchy-Riemann Equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$$f'(z) = \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{f(z + iy) - f(z)}{iy}$$

$$\frac{\partial f}{\partial x} = -i \frac{\partial f}{\partial y}$$

### Example

$$f(z) = x + xy + i\sqrt{x^2+y^2} \text{ analytic.}$$

$$\frac{\partial u}{\partial x} = 1 + y = -\frac{\partial v}{\partial y} \quad -\frac{\partial u}{\partial y} = -x = \frac{\partial v}{\partial x}$$

$$\sqrt{x^2+y^2} = \int \frac{\partial v}{\partial y} dy = y + \frac{1}{2}y^2 + h(x)$$

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \Rightarrow h'(x) = -x \quad h(x) = -\frac{1}{2}x^2 + C$$

$$\sqrt{x^2+y^2} = y + \frac{1}{2}y^2 - \frac{1}{2}x^2 + C$$

### Consequences

Real + Imaginary parts of an analytic function are harmonic!

$$\nabla^2 \phi = \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0$$

### Wall

Ex 3: "Wall"  $\nabla^2 \phi = 0$  here

Sol'n  
Define two angles  
 $\theta_1 = 0, \theta_2 = \pi$   
 $\theta_3 = \pi, \theta_4 = 2\pi$   
 $\theta_5 = 2\pi, \theta_6 = 0$

$$\phi = A_1 \theta_1 + A_2 \theta_2 + B$$

$$= A_1 \operatorname{Arg}(z_{\text{in}}) + A_2 \operatorname{Arg}(z_{\text{out}}) + B$$

### Impose B.C.

For  $z = x + iy$ , have  $\theta_1 = \theta_2 = 0$   
 $\phi = A_1(0) + A_2(0) + B$   
 $\Rightarrow B = 0$

For  $-1 < x < 1$ :  $\theta_1 = 0, \theta_2 = \pi$   
 $\phi = A_1(0) + A_2(\pi) + B$

$\phi = A_1(0) + A_2(\pi) + B$   
 $A_2 = \frac{1}{\pi}$

For  $z = x + iy$ :  $\theta_1 = 0, \theta_2 = \pi$   
 $\phi = A_1(0) + A_2(\pi) + B$

$\phi = A_1(0) + A_2(\pi) + B$   
 $A_1 = -\frac{1}{\pi}$

Therefore,  
 $\phi = \frac{1}{\pi} [\operatorname{Arg}(z_{-1}) - \operatorname{Arg}(z_1)]$   
 $= \frac{1}{\pi} [\theta_2 - \theta_1]$

### Cauchy's Theorem

$\oint f(z) dz = 0$ ,  $f$  analytic in a simply connected domain  $D$ ,  $\gamma$  curve in  $D$

### Consequences

## Useful Stuff

- $\cos(z) = \cosh(y) \cos(x) + i \sinh(y) \sin(x)$
- $\sin(z) = \cosh(y) \sin(x) + i \sinh(y) \cos(x)$
- All polynomials are analytic

## Exponential Function $f(z) = e^z$

$$e^z = e^x e^{iy} = e^x [\cos(y) + i \sinh(y)]$$

periodic w/ period  $2\pi i$

One-to-one in this strip!

## Complex Power

$$f(z) = z^\alpha$$

$$z^\alpha = e^{\alpha \log z} = e^{\alpha \log(z)}$$

$\log$  has branch cuts so  $z^\alpha$  has branch cuts

## Newton's Method

$$z_{n+1} = z_n - \frac{f(z_n)}{f'(z_n)}$$

## Logarithms

$$\log(z) = \log(r) + i \arg(z)$$

Multivalued! Problem!  
Branch cut!

$$\operatorname{Log}(z) = \operatorname{Log}(r) + i \operatorname{Arg}(z)$$

$\operatorname{Arg}(z)$  is harmonic in  $\mathbb{C} - \{\text{branch cut}\}$


$$f(a) = \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{i\theta}) d\theta$$

$f$  analytic in a neighborhood of  $a$

## Washers

$$\phi(r) = A \ln(r) + B \text{ since it doesn't depend on } \theta$$

At end,  $\phi(z) = A \ln(|z-a|) + B$ ;  $A, B$  solved earlier



Solution of the four washers is generally

$$\phi = A \ln(r) + B$$

$$0 = A \ln(r') + B \Rightarrow B = 0$$

$$10 = A \ln(r) + B$$

$$A = \frac{10}{\ln(r/r')}$$

$$\phi(z) = \frac{10}{\ln(r)} \ln|z-z_0|$$

## Wedges



$\phi(\theta) = A\theta + B$  From  $(r_i, i)$  is the "new origin"

At  $1, \theta = 0$   
 $\phi(0) = 10 = B$   
 $\phi(\frac{3\pi}{2}) = 0 = 10 + A\frac{3\pi}{2}$

$A = -\frac{20}{3\pi}$

$\phi(\theta) = -\frac{20}{3\pi} \theta + 10$   
 ↓ as a function of  $z$

$\phi(z) = -\frac{20}{3\pi} \operatorname{Arg}(z-2_0) + 10$

## Antiderivative Property

Path dependent, let  $F'(z) = f(z)$   
 $f$  continuous on open, connected set


$$\int_{z_0}^{z_1} f(z) dz = F(z_1) - F(z_0)$$

### Proof

$$z = z(t), 0 \leq t \leq 1, z(0) = z_0, z(1) = z_1$$

$$\begin{aligned} \int_{z_0}^{z_1} f(z) dz &= \int_{t=0}^{t=1} f(z(t)) \frac{dz}{dt} dt = \int_{t=0}^{t=1} F'(z(t)) \frac{dz}{dt} dt \\ &= \int_{t=0}^{t=1} \frac{d}{dt} (F(z(t))) dt = F(z(1)) - F(z(0)) \\ &= F(z_1) - F(z_0) \end{aligned}$$

$\int_{\gamma} \frac{dz}{z-a}$  where  $\gamma$  is any contour which encircles  $a$  once positively



$\frac{1}{z-a}$  is analytic everywhere EXCEPT  $z=a$ .

By last result, we can deform contour to a small circle  $C_r$  around  $a$ .

$$\int_{\gamma} \frac{dz}{z-a} = \int_{C_r} \frac{dz}{z-a} \quad z = re^{it} + a \quad 0 \leq t \leq 2\pi$$

$$\Rightarrow \int_0^{2\pi} \frac{rie^{it}}{re^{it} + a} dt = 2\pi i$$

## ML Estimate

$$\left| \int_Y f(z) dz \right| \leq \max_{z \in Y} |f(z)| \cdot \text{length}(Y)$$

### Proof

$$\left| \sum_{k=1}^N f(z_k) \Delta z_k \right| \leq \left| \sum_{k=1}^N f(z_k) \right| \left\| \Delta z_k \right\| \leq M \sum_{k=1}^N |f(z_k)|$$

$$\int_{\gamma} (z-a)^n = 0, n \neq -1$$

Proof  $\frac{1}{n+1} (z-a)^{n+1}$  is single valued in  $\mathbb{C} - \{a\}$  (punctured plane)

## CIF Example

$$\int_{\gamma} \frac{\cos(z)}{(z-\pi)(z+5i)} dz$$

$$f(z) = \cos(z)/(z+5i) \rightarrow \int_{\gamma} \frac{f(z)}{z+5i} dz = 2\pi i f(5i)$$

z finitely differentiable

## CIF For Derivative

$$f^{(n)}(a) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz$$

## Bounds on Derivative

$$f(z) \text{ analytic on } \Gamma \text{ inside circle of radius } R \text{ about } a.$$

$$M = \max_{z \in \Gamma} |f(z)| \text{ then } |f^{(n)}(a)| \leq \frac{M^n |f(z)|}{R^n}$$

$$\text{Proof: } f^{(n)}(a) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz$$

$$\text{ML Estimate: } \left| f^{(n)}(a) \right| \leq \max_{z \in \Gamma} \left| \frac{f(z)}{(z-a)^{n+1}} \right| \frac{n!}{R^n}$$

$$\leq \max_{z \in \Gamma} |f(z)| \frac{n!}{R^n} R^n = M \frac{n!}{R^n}$$

$$\text{So, } \left| f^{(n)}(a) \right| \leq \frac{M n!}{R^n}$$

## Cauchy's Integral Formula

$$f(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z-a} dz$$

'Proof': Deform  $\gamma$  to  $C_r$

$$\int_{\gamma} \frac{f(z)}{z-a} dz = \int_{C_r} \frac{f(z)}{z-a} dz. \text{ On } C_r, f(z) = f(a) + \varepsilon(z)$$

Sub  $f$  in, ML estimate on  $\varepsilon$

## Louisville's Theorem

$f(z)$  is entire and bounded by some constant. If  $|f(z)| \leq M + z$ . Then  $f(z)$  must be constant.

**Proof**  
 $|f'(z)| \leq \frac{M}{R} + R$ . Since  $M$  is independent of  $R$  by boundedness. As  $R \rightarrow \infty$ ,  $|f'(z)| \rightarrow 0$

## Maximum Modulus Theorem

A function analytic in a bounded domain, continuous up to and including its boundary, attains its max or min on the boundary.

## Isolated Singularities

$f(z)$  has an isolated singularity at  $z_0$  if  $f(z)$  is analytic in punctured disk  $0 < |z - z_0| < R$

1) Removable  $|f(z)|$  meaning finite/doesn't blow up bounded as  $z \rightarrow z_0$ .

2) pole  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

3) Essential  $f(z)$  has no limit as  $z \rightarrow z_0$ . oscillates wildly as  $z \rightarrow z_0$

## Deformation Theorem

If a contour sweeps only through analytic points as it is deformed, the value of the integral does NOT change.

## Residues

### The $a_{-1}$ term!


If  $f$  has a pole of order  $m$  at  $z_0$ , then

$$\text{Res}(f; z_0) = \lim_{z \rightarrow z_0} \frac{1}{(m-1)!} \frac{d^{m-1}}{dz^{m-1}} [(z-z_0)^m f(z)]$$

If  $\gamma$  is a simple closed contour oriented CCW, and  $f$  analytic in and on  $\gamma$  except at the points  $z_1, z_2, \dots, z_n$  inside  $\gamma$ , then

$$\int_{\gamma} f(z) dz = 2\pi i \sum_{n=1}^N \text{Res}(f; z_n)$$

Example:



Given this in the  $w$ -plane.

We want diameter = distance from  $\frac{1}{r}$  to edge

$$\frac{1}{1-r} - \frac{1}{1+r} = \frac{1}{1+r} - \frac{1}{2}$$


$$\Rightarrow r^2 - 6r + 1 = 0 \\ r = \frac{6 \pm \sqrt{32}}{2} = 3 \pm 2\sqrt{2}$$

We know a geometry in the  $z$ -plane which maps to something similar.  
 Solve for  $\beta(u, v)$  in region outside circle  
 Use inversion map


$$w = \frac{1}{z} = \frac{x}{x+iy} = \frac{x}{x^2+y^2} - i \frac{y}{x^2+y^2}$$

Only want  $r < 1$ , so


$$r = 3 - 2\sqrt{2}$$




If we can find an  $r < 1$  s.t. the circle on the  $z$ -plane maps to the given physical circle




We can simplify further by shifting circle to the origin.



Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!



Washer!




Washer!




Washer!




Washer!




Washer!



Washer!




Washer!




Washer!




Washer!



Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!



Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!



Washer!




Washer!




Washer!



Washer!




Washer!




Washer!



Washer!




Washer!




Washer!




Washer!




Washer!




Washer!




Washer!



Washer!



Washer!



Washer!

