

# MAXIMUM LIKELIHOOD ESTIMATORS

Joseph Nelson, Data Science Immersive

---

# AGENDA

---

- Review of Bayes Theorem
- What is Maximum Likelihood Estimation?
- Calculating MLE
- MLE

---

## BAYES THEOREM EXAMPLE

---

- ▶ Imagine you have one fair coin, and one double-sided heads coin. Build a tree that describes the possible outcomes we may expect.
- ▶ What is the probability that you have the fair coin?

---

## BAYES THEOREM EXAMPLE

---

- ▶ Imagine you have one fair coin, and one double-sided heads coin. Build a tree that describes the possible outcomes we may expect.
- ▶ What is the probability that you have the fair coin?
- ▶ Now, after you've built the tree, you find that the coin flipped was actually heads. What is the probability that the coin is the fair coin?

---

## BAYES THEOREM EXAMPLE

---

- ▶ Imagine you have one fair coin, and one double-sided heads coin. Build a tree that describes the possible outcomes we may expect.
- ▶ What is the probability that you have the fair coin?
- ▶ Now, after you've built the tree, you find that the coin flipped was actually heads. What is the probability that the coin is the fair coin?
- ▶ You repeat this exercise. Draw the tree.

---

## BAYES THEOREM EXAMPLE

---

- ▶ Imagine you have one fair coin, and one double-sided heads coin. Build a tree that describes the possible outcomes we may expect.
- ▶ What is the probability that you have the fair coin?
- ▶ Now, after you've built the tree, you find that the coin flipped was actually heads. What is the probability that the coin is the fair coin?
- ▶ You repeat this exercise. Draw the tree.
- ▶ You find that the coin flipped was heads again. What is the probability that the coin is a fair coin?

---

## BAYES THEOREM

---

- ▶ Probability of a fair coin given a heads:  
probability of a fair coin divided by all the  
chances of having a heads

$$\Pr(A|X) = \frac{\Pr(X|A) \Pr(A)}{\Pr(X)}$$

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- ▶ Imagine we have a Bernoilli Distribution:
- ▶  $p^5(1-p)^4$
- ▶ The function would have to "peak" at some point; therefore, in some way, the function would have to look somewhat like a inverted parabola
- ▶ This peak would have to be a 'global' peak, i.e. it can have multiple peaks, but only one can be the 'largest'
- ▶ Once we find the peak, the value/level of the peak is not what we're actually interested in (the Y-value of the function). What we're actually interested in is to guess which X-value of the function (the independent variable), needs to be inputted to get a particular Y.



---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- ▶ Imagine we have a Bernoilli Distribution:
- ▶  $p^5(1-p)^4$
- ▶ The function would have to "peak" at some point; therefore, in some way, the function would have to look somewhat like a inverted parabola
- ▶ This peak would have to be a 'global' peak, i.e. it can have multiple peaks, but only one can be the 'largest'
- ▶ Once we find the peak, the value/level of the peak is not what we're actually interested in (the Y-value of the function). What we're actually interested in is to guess which X-value of the function (the independent variable), needs to be inputted to get a particular Y.

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- Code: Graph
- Imagine we have a Bernoilli Distribution:
  - $p^5(1-p)^4$
  - The function would have to "peak" at some point; therefore, in some way, the function would have to look somewhat like a inverted parabola
  - This peak would have to be a 'global' peak, i.e. it can have multiple peaks, but only one can be the 'largest'
  - Once we find the peak, the value/level of the peak is not what we're actually interested in (the Y-value of the function). What we're actually interested in is to guess which X-value of the function (the independent variable), needs to be inputted to get a particular Y.

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- Code: Graph
- Differentiate: maximize the function (FOC)  $p^5(1-p)^4 = 0$

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- Code: Graph
- Differentiate: maximize the function (FOC)  $p^5(1-p)^4 = 0$
- Apply the product rule:  $5p^4(1-p)^4 - 4(1-p)^3p^5 = 0$

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- ▶ Code: Graph
- ▶ Differentiate: maximize the function (FOC)  $p^5(1-p)^4 = 0$
- ▶ Apply the product rule:  $5p^4(1-p)^4 - 4(1-p)^3p^5 = 0$
- ▶ “Balance” the equation:  $5p^4(1-p)^4 = 4(1-p)^3p^5$

---

## MAXIMUM LIKELIHOOD ESTIMATION

---

- Code: Graph
- Differentiate: maximize the function (FOC)  $p^5(1-p)^4 = 0$
- Apply the product rule:  $5p^4(1-p)^4 - 4(1-p)^3p^5 = 0$
- “Balance” the equation:  $5p^4(1-p)^4 = 4(1-p)^3p^5$
- Algebra:  $p = .55556$
- Did it work? Check the graph...

---

## WHAT DID WE JUST DO?

---

► Class?

---

## WHAT DID WE JUST DO?

---

- Class?
- We determined what  $x$  (input) values provides the maximum chance of some scenario.
- Let's do an example



---

## WHAT DID WE JUST DO?

---

- ▶ Let's say we have a coin that comes up heads with some probability  $\theta$
- ▶ We see two heads come up. Our likelihood then becomes  $P(D|\theta)=\theta^2$
- ▶ How do we maximize this function?

---

## WHAT DID WE JUST DO?

---

- ▶ Let's say we have a coin that comes up heads with some probability  $\theta$
- ▶ We see two heads come up. Our likelihood then becomes  $P(D|\theta)=\theta^2$
- ▶ How do we maximize this function?
- ▶  $\theta=1$

---

## WHAT DID WE JUST DO?

---

- ▶ Let's say we have a coin that comes up heads with some probability  $\theta$
- ▶ We see two heads come up. Our likelihood then becomes  $P(D|\theta)=\theta^2$
- ▶ How do we maximize this function?
- ▶  $\theta=1$
- ▶ So, our MLE is that the coin always comes up heads, and so we predict future coins will all come up head

---

## WHAT DID WE JUST DO?

---

- ▶ Let's say we have a coin that comes up heads with some probability  $\theta$
- ▶ We see two heads come up. Our likelihood then becomes  $P(D|\theta)=\theta^2$
- ▶ How do we maximize this function?
- ▶  $\theta=1$
- ▶ So, our MLE is that the coin always comes up heads, and so we predict future coins will all come up head

---

## BAYES VS MLE: DOCTOR

---

- ▶ Imagine you are a doctor. You have a patient who shows an odd set of symptoms. You look in your doctor book and decide the disease could be either a common cold or lupus. Your doctor book tells you that if a patient has lupus then the probability that he will show these symptoms is 90%. It also states that if the patient has a common cold then the probability that he will show these symptoms is only 10%. Which disease is more likely?
- ▶ (Hint: does rarity of disease matter?)

---

## BAYES VS MLE: DOCTOR

---

- ▶ Imagine you are a doctor. You have a patient who shows an odd set of symptoms. You look in your doctor book and decide the disease could be either a common cold or lupus. Your doctor book tells you that if a patient has lupus then the probability that he will show these symptoms is 90%. It also states that if the patient has a common cold then the probability that he will show these symptoms is only 10%. Which disease is more likely?

---

## MLE: A BETTER PATH – POSTERIOR ESTIMATION

---

- Maximum A-Posterior Estimation (MAP)
- Estimate the maximum likelihood of the posterior rather than the prior
- $\theta_{\text{MAP}} = \operatorname{argmax}_{\theta} P(\theta|D)$
- $\theta_{\text{MLE}} = \operatorname{argmax}_{\theta} P(D|\theta)$

---

## MLE: A BETTER PATH – POSTERIOR ESTIMATION

---

- Maximum A-Posterior Estimation (MAP)
- Estimate the maximum likelihood of the posterior rather than the prior
- $\theta_{\text{MAP}} = \operatorname{argmax}_{\theta} P(\theta|D)$
- $\theta_{\text{MLE}} = \operatorname{argmax}_{\theta} P(D|\theta)$
  
- Code Example



---

## MLE: A BETTER PATH – POSTERIOR ESTIMATION

---

- Maximum A-Posterior Estimation (MAP)
- Estimate the maximum likelihood of the posterior rather than the prior
- $\theta_{\text{MAP}} = \operatorname{argmax}_{\theta} P(\theta|D)$
- $\theta_{\text{MLE}} = \operatorname{argmax}_{\theta} P(D|\theta)$
  
- Advantages:
- Easy and interpretable
- Avoids overfitting
- Tends to follow the same asymptotic distribution

---

## MLE: A BETTER PATH – POSTERIOR ESTIMATION

---

- ▶ Maximum A-Posterior Estimation (MAP)
- ▶ Estimate the maximum likelihood of the posterior rather than the prior
- ▶  $\theta_{\text{MAP}} = \operatorname{argmax}_{\theta} P(\theta|D)$
- ▶  $\theta_{\text{MLE}} = \operatorname{argmax}_{\theta} P(D|\theta)$
  
- ▶ Disdvantages:
- ▶ Must assume a prior on  $\theta$
- ▶ No representation of uncertainty in  $\theta$