Aggistic Regission > Classification Algorithm > In Reglession Bollom we used to pedicted a Centimorial Value. Whele as, in classification Problem we pledict whother it is a yes of no type. > In Llucal Replession we used distance flawclood to price of Col. In Classification we will use some other flatures also to pladict whether the Col is Costly of chapter (yes) (PD) > Consider Some flatures like > Kenk Size > Planuforthere Now these fentures will be feel to a mathematical function F(x) and that function will gives different plotabilities. Suppose, if the probability is > 0.5 then Cos is (settler). This can happen vice-versa also -> Function would be like -> Function would be like -> Sigmoid F(x) = (Ax, +1, +2, Az observer). Here A, Az Az observery.	_	1	-	-					
> In Reglession Problem we used to pledicted a Continuous value. Whele as, in classification Problem we pledict whother it is a yes of no type. > In Liveal Replession we used distance flowellod to plice of Get. In Classification we will list some other features also to pledict whether the Get is Costly of chapper (yes) (PD) > Consider Some flatures like > Kank Size > Klange > Howeforther will be fed to a mathematical function F(x) and that function will gives different plobabilities. Quepose, if the probability is > 0-5 then Cos is Gettier. This can hoppen vice-versa also. -> Function would be like -> Gigmoid F(x) = (A1x, +A2x2 +A3xX3) + Kee A, A2A Obe weights and x, x2 x3 ale				ogistic	Rogse	88.ion		1 .	
> In Reglession Problem we used to pledicted a Continuous value. Whele as, in classification Problem we pledict whother it is a yes of no type. > In Liveal Replession we used distance flowellod to plice of Get. In Classification we will list some other features also to pledict whether the Get is Costly of chapper (yes) (PD) > Consider Some flatures like > Kank Size > Klange > Howeforther will be fed to a mathematical function F(x) and that function will gives different plobabilities. Quepose, if the probability is > 0-5 then Cos is Gettier. This can hoppen vice-versa also. -> Function would be like -> Gigmoid F(x) = (A1x, +A2x2 +A3xX3) + Kee A, A2A Obe weights and x, x2 x3 ale		01-00:1:	c la /	10 -00 10			• • • • • • • • • • • • • • • • • • • •	94 - 1	-
Value. Whele as, in classification Problem we pledict whather it is a yes of no type. In Lineal Replession we used distance flevelled to peice of Col. In Classification we will use some other features also to pledict whether the Col is Costly of chaques like \(\) (yes) (Po) Consider Some features like \(\) kenk Size \(\) Kink Size \(\) Kink Size \(\) Kink Size \(\) Hileage \(\) A mathematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0.5 then Cos is (88 then.) This can happen vice-versa also. Frunction Coould be like \(\) Sigmoid \(\) F(x) = \(\) (Axx, Axx = Ax = Ole weights ord x, xx, xx, xx = Ole \(\)		Cassi ji	ation t	1gouth	<u></u>				,
Value. Whele as, in classification Problem we pledict whather it is a yes of no type. In Lineal Replession we used distance flevelled to peice of Col. In Classification we will use some other features also to pledict whether the Col is Costly of chaques like \(\) (yes) (Po) Consider Some features like \(\) kenk Size \(\) Kink Size \(\) Kink Size \(\) Kink Size \(\) Hileage \(\) A mathematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0.5 then Cos is (88 then.) This can happen vice-versa also. Frunction Coould be like \(\) Sigmoid \(\) F(x) = \(\) (Axx, Axx = Ax = Ole weights ord x, xx, xx, xx = Ole \(\)		To Ros	Passino	Poollow	2 150	100A 1- 00	الدوا ادا	- Co.15	100 / 100)
> In Liveal Replession we used distance flevelled to flice of Gel. In Classification we will like Some other flectures also to pledict whether the Gel is Costly of chaquel (yes) (PO) > Consider Some flectures like > Kenk Size > Kiloage > Llouge > Llouge > Long these fentules will be feel to a mathematical function F(x) and that function will gives different plobabilities. Suppose, if the plobability is > 0.5 then Cas is Gettler. This can happen vice-velsa also. > Frunction coould be like F(x) = (A, x, +A, x, +A, x, x, +A, x,		Volue	whole	1,x000m	Class i	ication Po	alle-a	a Conna	<u>uous</u> ': c+
> In Liveal Replession we used distance flevelled to flice of Gel. In Classification we will like Some other flectures also to pledict whether the Gel is Costly of chaquel (yes) (PO) > Consider Some flectures like > Kenk Size > Kiloage > Llouge > Llouge > Long these fentules will be feel to a mathematical function F(x) and that function will gives different plobabilities. Suppose, if the plobability is > 0.5 then Cas is Gettler. This can happen vice-velsa also. > Frunction coould be like F(x) = (A, x, +A, x, +A, x, x, +A, x,		(Shother	it is	a 40	8 8 00	1.00	UVUITI (se preu	161
flice of lot. In Clessification we will like some other features also to predict whether the Cal is Costly of charges (yes) (ro) Consider some features like ⇒ Kenk Size ⇒ Milange ⇒ Many Jochese Now these fentules will be fed to a mashematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0.5 then Cas is Costlier. This can happen vice-versa also. → Function would be like → Signnoid F(x) = (+e^{-(A_1x_1+A_2x_2+A_3x_3x_3)}) Here A, A, A, a see weights and x, x, x, x, x, x, a ale		in in	hal-	1		9/4	J .		
flice of lot. In Clessification we will like some other features also to predict whether the Cal is Costly of charges (yes) (ro) Consider some features like ⇒ Kenk Size ⇒ Milange ⇒ Many Jochese Now these fentules will be fed to a mashematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0.5 then Cas is Costlier. This can happen vice-versa also. → Function would be like → Signnoid F(x) = (+e^{-(A_1x_1+A_2x_2+A_3x_3x_3)}) Here A, A, A, a see weights and x, x, x, x, x, x, a ale	->	In Lin	eal Rep	Cession	we u	red disto	ncelleve	elledto	
(yes) (Po) Consider Some Jeatures (ike ⇒ Kunk Size ⇒ Klange ⇒ Klange ⇒ Klange † Slangother Now these Jentures will be Jed to a mathematical function F(x) and that Junction will gives different plobabilities. Suppose, if the probability is > 0-5 then Cas is Costfied. This can happen vice-velsa also. → Function coould be Coke → Sigmoid F(x) = (+e^{-(A_1x_1+A_2x_2+A_3x_3)}) Here A, A, A, a sole weights and x, x, x, x, x, a ale		plice of	6l. I	a Clas	sifical	on we w	il Use	Some	•
(yes) (Po) Consider Some Jeatures (ike ⇒ Kunk Size ⇒ Klange ⇒ Klange ⇒ Klange † Slangother Now these Jentures will be Jed to a mathematical function F(x) and that Junction will gives different plobabilities. Suppose, if the probability is > 0-5 then Cas is Costfied. This can happen vice-velsa also. → Function coould be Coke → Sigmoid F(x) = (+e^{-(A_1x_1+A_2x_2+A_3x_3)}) Here A, A, A, a sole weights and x, x, x, x, x, a ale		other !	latules	also	to pee	dict wh	eshor s	he Cal	
(yes) (PO) Consider Some Jeatures like ⇒ Kank Size ⇒ Kileage ⇒ Llanufactures Now those Jentures will be fed to a mathematical Junction F(x) and that Junction will gives different plobabilities. Suppose, if the probability is > 0-5 then Cas is Costfiel. This Can happen vice-velsa also. → Function coould be like -> Signaid F(x)= (+e^{(A_1x_1+A_2x_2+A_3x_3x_3}) Here A, A, A, A, are the weights and x, x, x, x, x, x, and are the supposed to the suppose of the supposed to the supp		is Cos	tly of	hagee	R)	THE THE THE	16 10	14.	
> Consider Some Jeatwee like ⇒ Kenk Size ⇒ Mileage ⇒ Lanufockeles Now these Jenfules will be Jeal to a mathematical Junction F(x) and that Junction will gives different plobabilities. Suppose, if the probability is > 0-5 then Cas is Costfiel. This can happen vice-velsa also. → Function would be leke → Signnoid F(x) = 1 1 te CA, x, +A, A, a all weights and x, x, x, x, x, alle		Contract Co	jes)	(PO)) Mi	12 110	<u>s</u> • • •	, the gray	, 13
⇒ Kenk Size ⇒ Kileage ⇒ Llanufactures Now these sentures will be fed to a mathematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0-5 then Car is Corting. This can happen vice-versa also. → Function would be like -> Signoid F(x)= (+e^-(A_1x,+A_2x_2+A_3x_3)) Here A, A, A, A, a are weights and x, x, x, x, x, x, alle		(tab)			DO BERTHA	ALL ALL	i a.	0-1	. î,
⇒ Kenk Size ⇒ Kileage ⇒ Llanufactures Now these sentures will be fed to a mathematical function F(x) and that function will gives different plobabilities. Suppose, if the probability is > 0-5 then Car is Corting. This can happen vice-versa also. → Function would be like -> Signoid F(x)= (+e^-(A_1x,+A_2x_2+A_3x_3)) Here A, A, A, A, a are weights and x, x, x, x, x, x, alle		Consider	Some	Patules	z like		24/3		idaa
Now those sentules will be fed to a mathematical function $F(x)$ and that function will gives different plobabilities. Suppose, if the probability is > 0.5 then Cas is Costfied. This can happen vice-versa also. First conduit be like First in coord be like Signaid F(x) = 1		> .	Kenk Si	"zl					
Junction F(x) and shot function will gives different phobabilities. Suppose, if the phobability is > 0-5 then Cas is Costfiel. This can happen vice-velsa also. F(x) = 1 (A, Y, +A, Y2 +A3 × M3) Here A, A2, A3 are weights and x, x2, x3 are									
Suppose, if the plobability is >0-5 then Cas is Costlied. This can happen vice-velsa also. Frenchian would be like F(x) = 1 1 te (A, x, +A, x,)) /	Conufoct	rills.	. 77. (F 1 1	0 1	V 1:0	1
Suppose, if the plobability is >0-5 then Cas is Costlied. This can happen vice-velsa also. Frenchian would be like F(x) = 1 1 te (A, x, +A, x,	/	NOW OF	ese je	a fulles	(e)// (e jed to	a mag	shomatila	+
Suppose, if the plobability is > 0-5 then Cas is Costlice. This can happen vice-velsa also. Function would be like F(x) = 1 1 + e^{-(A_1 x_1 + A_2 x_2 + A_3 x_3 x_3)} Here A, A, A, A, a ole weights and x, x, x, x, x, ale	0	on-lon-	<u> </u>	and mo	of fun	CHOA WILL	gres	<u>Aijxelln</u>)
F(x)= 1 I te A, A2, A3 Ole weights and x, x2, x3 ale		poul	<u> 125 · </u>					1 .	
F(x)= 1 I te A, A2, A3 Ole weights and x, x2, x3 ale		0,000%	il the r	280/50/50	ido is	>0-5 She	n Cas i	s Costfi	ol.
F(x)= 1 I te A, A2, A3 Ole weights and x, x2, x3 ale		This con	700000	Vi(0-V	osea al	\(\rac{1}{2}\).			
F(x)= 1 1+e ^{-(A₁x₁+A₂x₂+A₃xx₃) Here A₁, A₂, A₃ ple weights and x₁, x₂, x₃ ale}		1145	- response		,				
F(x)= 1 1+e ^{-(A₁x₁+A₂x₂+A₃xx₃) Here A₁, A₂, A₃ ple weights and x₁, x₂, x₃ ale}	7	Function	Coould	le like					8.
Here A, A2, A3 ple weights and x, x2, x3 ale							-> 5	ignoid	
. Here A, Az, Az ole weights and x, xz, xz ale			F(x)	=	A. X. LA	A HARKA		<u>J</u>	
Les A, Az, Az ole weights and x, xz, xz all features lespectively.				ite		A. A. 1. 13 17 15			
features lespectively.	•	Heee	A, Az,	13-0	de Wei	ohts on	(x, x)	2, ×3 Ql	le
	E Ste	featur	s les	pectively	5-				

