


THE COMPLEX LANGEVIN METHOD FOR ULTRACOLD FERMIONS

Lukas Rammelmüller, TU Darmstadt
Marburg, FOR 1807 Winter School 2018


[LR, Loheac, Drut, Braun *in preparation*]

[LR, Porter, Drut, Braun *Phys. Rev. D* 96, 094506, 2017]


[LR, Porter, Drut *Phys. Rev. A* 93, 033639, 2016]




(electrons in metals, nuclear physics,
neutron stars, controllable experiments, ...)




(electrons in metals, nuclear physics,
neutron stars, controllable experiments, ...)



(electrons in metals, nuclear physics,
neutron stars, controllable experiments, ...)



spin polarized systems
(inhomogeneous phases?)




mass imbalanced systems
(inhomogeneous phases?)

model: **contact interaction**

$$\hat{H} = - \sum_{s=\uparrow,\downarrow} \int d^d x \, \hat{\psi}_s^\dagger(\vec{x}) \left(\frac{\hbar^2 \vec{\nabla}^2}{2m_s} \right) \hat{\psi}_s(\vec{x}) \\ + g \int d^d x \, \hat{\psi}_\uparrow^\dagger(\vec{x}) \hat{\psi}_\uparrow(\vec{x}) \hat{\psi}_\downarrow^\dagger(\vec{x}) \hat{\psi}_\downarrow(\vec{x})$$

model: **contact interaction**

$$\hat{H} = - \sum_{s=\uparrow,\downarrow} \int d^d x \, \hat{\psi}_s^\dagger(\vec{x}) \left(\frac{\hbar^2 \vec{\nabla}^2}{2m_s} \right) \hat{\psi}_s(\vec{x}) \\ + g \int d^d x \, \hat{\psi}_\uparrow^\dagger(\vec{x}) \hat{\psi}_\uparrow(\vec{x}) \hat{\psi}_\downarrow^\dagger(\vec{x}) \hat{\psi}_\downarrow(\vec{x})$$



what do we need to compute?

$$\mathcal{Z} = \text{Tr}[e^{-\beta\hat{H}}] = \text{Tr}[e^{-\beta(\hat{T} + \hat{V})}]$$
$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \text{Tr}[\hat{\mathcal{O}} e^{-\beta\hat{H}}]$$

what do we need to compute?

$$\mathcal{Z} = \text{Tr}[e^{-\beta\hat{H}}] = \text{Tr}[e^{-\beta(\hat{T} + \hat{V})}]$$
$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \text{Tr}[\hat{\mathcal{O}} e^{-\beta\hat{H}}]$$

discretize space & time, decouple the interaction
with a **Hubbard Stratonovich transformation**

what do we need to compute?

$$\mathcal{Z} = \text{Tr}[e^{-\beta\hat{H}}] = \text{Tr}[e^{-\beta(\hat{T} + \hat{V})}]$$
$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \text{Tr}[\hat{\mathcal{O}} e^{-\beta\hat{H}}]$$

discretize space & time, decouple the interaction
with a **Hubbard Stratonovich transformation**

Rewrite the problem as a path-integral:

$$\mathcal{Z} = \int \mathcal{D}\phi \det M_{\phi}^{\uparrow} \det M_{\phi}^{\downarrow} \equiv \int \mathcal{D}\phi e^{-S[\phi]}$$

calculating observables

$$\langle \mathcal{O} \rangle = \int \mathcal{D}\phi P[\phi] \mathcal{O}[\phi]$$

produce a set of **random auxiliary-field configurations** $\{\phi_i\}$,
compute observables & average for expectation values

calculating observables

$$\langle \mathcal{O} \rangle = \int \mathcal{D}\phi \, P[\phi] \, \mathcal{O}[\phi]$$

produce a set of **random auxiliary-field configurations** $\{\phi_i\}$,
compute observables & average for expectation values

probability measure:

$$P[\phi] \propto \det M_{\phi}^{\uparrow} \det M_{\phi}^{\downarrow} = \begin{cases} \text{attractive, balanced case} \rightarrow \text{OK} \\ \text{otherwise} \rightarrow \text{sign problem} \end{cases}$$

Complex Langevin (in a nutshell)


stochastic quantization: equilibrium distribution of a $(d + 1)$ -dimensional random process is identified with the probability measure of our d -dimensional path integral

random walk governed by **Langevin equation** (Brownian motion):

$$\frac{\partial \phi}{\partial t} = - \frac{\partial S[\phi]}{\partial \phi} + \eta(t)$$

first step: compare to other methods



[LR, Porter, Drut, Braun '17]



[BA: Iida, Wadati '07; Tracy, Widom '16]
[DFT-RG: Kemler, Pospiech, Braun '16]
[HMC: LR, Porter, Loheac, Drut '15]

first step: compare to other methods

[LR, Porter, Drut, Braun '17]



repulsive side:
Outliers skew
expectation values!
(**'fat tail' problem**)

$$\gamma = g/n$$

[BA: Iida, Wadati '07; Tracy, Widom '16]




[DFT-RG: Kemler, Pospiech, Braun '16]

[HMC: LR, Porter, Loheac, Drut '15]

first step: compare to other methods

[LR, Porter, Drut, Braun '17]

attractive side:
no outliers,
no problems!




repulsive side:
Outliers skew
expectation values!
(*'fat tail' problem*)

[BA: Iida, Wadati '07; Tracy, Widom '16]
[DFT-RG: Kemler, Pospiech, Braun '16]
[HMC: LR, Porter, Loheac, Drut '15]


polarized 1D fermions: equation of state

[LR, Drut, Braun *in preparation*]



polarized 1D fermions: pair correlation

[LR, Drut, Braun *in preparation*]




$$\gamma = g/n$$
$$p = \frac{N_{\uparrow} - N_{\downarrow}}{N_{\uparrow} + N_{\downarrow}}$$

$$n_{pair}(x, x') = \langle \hat{\psi}_{\uparrow}^{\dagger}(x) \hat{\psi}_{\downarrow}^{\dagger}(x) \hat{\psi}_{\downarrow}(x') \hat{\psi}_{\uparrow}(x') \rangle$$

polarized 1D fermions: pair correlation

[LR, Drut, Braun *in preparation*]




$$\gamma = g/n$$

$$p = \frac{N_{\uparrow} - N_{\downarrow}}{N_{\uparrow} + N_{\downarrow}}$$

$$n_{pair}(x, x') = \langle \hat{\psi}_{\uparrow}^{\dagger}(x) \hat{\psi}_{\downarrow}^{\dagger}(x) \hat{\psi}_{\downarrow}(x') \hat{\psi}_{\uparrow}(x') \rangle$$

polarized 1D fermions: pair correlation

[LR, Drut, Braun *in preparation*]



signature of
FFLO type
pairing


$$\gamma = g/n$$

$$p = \frac{N_\uparrow - N_\downarrow}{N_\uparrow + N_\downarrow}$$

$$n_{pair}(x, x') = \langle \hat{\psi}_\uparrow^\dagger(x) \hat{\psi}_\downarrow^\dagger(x) \hat{\psi}_\downarrow(x') \hat{\psi}_\uparrow(x') \rangle$$

UFG at finite T: equation of state

[LR, Loheac, Drut, Braun *in preparation*]




[experiment/BDMC: van Houcke et al. '12]

[DHMC: Drut, Lähde, Wlazlowski, Magierski '12]

UFG at finite T: equation of state

[LR, Loheac, Drut, Braun *in preparation*]


$$T_C \sim 0.17 T_F$$



[experiment/BDMC: van Houcke et al. '12]
[DHMC: Drut, Lähde, Wlazlowski, Magierski '12]

UFG at finite T: equation of state

[LR, Loheac, Drut, Braun *in preparation*]



CL results:
finite lattice!
($V = 9^3$)

low temperatures: λ_T increases
($\lambda_T \ll V^{1/3}$ must be fulfilled)


[experiment/BDMC: van Houcke et al. '12]
[DHMC: Drut, Lähde, Wlazlowski, Magierski '12]

UFG at finite T: equation of state

[LR, Loheac, Drut, Braun *in preparation*]

$$T_C \sim 0.17 T_F$$

good agreement
with experiment
and other
methods!




CL results:
finite lattice!
($V = 9^3$)

low temperatures: λ_T increases
($\lambda_T \ll V^{1/3}$ must be fulfilled)

[experiment/BDMC: van Houcke et al. '12]
[DHMC: Drut, Lähde, Wlazlowski, Magierski '12]

UFG at finite T: phase diagram (mean field)




$$h = \frac{\mu_{\uparrow} - \mu_{\downarrow}}{2}$$

[mean-field phase diagram: e.g. Chevy, Mora '10; Braun et al. '13]

[beyond mean-field:: e.g. Boettcher et al. '14; Roscher, Braun, Drut '15]

UFG at finite T: phase diagram (mean field)



$$h = \frac{\mu_{\uparrow} - \mu_{\downarrow}}{2}$$

[mean-field phase diagram: e.g. Chevy, Mora '10; Braun et al. '13]

[beyond mean-field:: e.g. Boettcher et al. '14; Roscher, Braun, Drut '15]

UFG at finite T: equation of state

[LR, Loheac, Drut, Braun *in preparation*]

$$\beta h = 2.0$$


$$\beta h = 1.6$$

$$\beta h = 1.2$$

$$\beta h = 0.8$$

$$\beta h = 0.4$$

$$\beta h = 0.0$$



$$h = \frac{\mu_{\uparrow} - \mu_{\downarrow}}{2}$$

$$\text{virial expansion: } \ln \mathcal{Z} = Q_1 \sum_n z^n b_n$$

RECAP & OUTLOOK

imbalanced Fermi gases are hard to treat:
accessible with the **complex Langevin** method

CL compares well with other methods wherever possible

works in the ground state & at finite temperature
in any spatial dimension

Up next: investigation of pair correlations in $d > 1$,
mass imbalance at finite temperature, phase diagram in 2D/3D, ...