

Approaching Fermi-Fermi mixtures with complex Langevin: EoS and pairing

Lukas Rammelmüller^{1,2}, Joaquín E. Drut³, Jens Braun²

¹Ludwig-Maximilians Universität München, Arnold Sommerfeld Center, ²Technische Universität Darmstadt, IKP Theory Center, ³University of North Carolina at Chapel Hill, Department of Physics and Astronomy

Theoretical treatment of imbalanced Fermi systems is challenging. Exact analytic methods, if available, are limited to 1D setups and thus numerical treatment is often the only viable option. Among the most successful methods for balanced Fermi gases, in particular for systems beyond the few-body regime, are Quantum Monte Carlo (QMC) approaches. For imbalanced Fermi systems, however, these approaches suffer from an exponential scaling with system size: the infamous sign-problem. A way to circumvent this issue is provided by the complex Langevin method which we employ to 1D imbalanced fermions as well as the 3D unitary Fermi gas with finite spin asymmetry. [\[recent review: Berger, LR, Loheac, Ehmann, Braun, Drut '19\]](#)

complex Langevin in a nutshell

- after discretizing space and (imaginary) time and performing a Hubbard-Stratonovich transformation, we can write the partition sum \mathcal{Z} as a path integral over the auxiliary field ϕ :

$$\mathcal{Z} \equiv \text{Tr}[e^{-\beta\hat{H}}] \rightarrow \int \mathcal{D}\phi e^{-S[\phi]}$$

- similarly, we can compute observables:

$$\langle \hat{O} \rangle = \frac{\text{Tr}[\hat{O} e^{-\beta\hat{H}}]}{\text{Tr}[e^{-\beta\hat{H}}]} \rightarrow \frac{1}{\mathcal{Z}} \int \mathcal{D}\phi \mathcal{O}[\phi] e^{-S[\phi]}$$

- key idea of stochastic quantization: a $(d+1)$ -dimensional random process is used to sample the measure of a d -dimensional euclidean path integral

$$\frac{\partial \phi}{\partial t} = -\frac{\delta S[\phi]}{\delta \phi} + \eta$$

- with a discrete Langevin equation we can generate a Markov chain of complexified auxiliary fields ϕ that can be used to compute observables stochastically



HIC for FAIR

DFG Deutsche Forschungsgemeinschaft
German Research Foundation

HGS-HIRE for FAIR
Heinrich Heine Graduate School for Hadron and Ion Research

TECHNISCHE UNIVERSITÄT DARMSTADT

LMU LUDWIG-MAXIMILIANS-UNIVERSITÄT MÜNCHEN


erc European Research Council

MCQST

3D unitary fermions with finite spin asymmetry

[\[LR, Loheac, Drut, Braun '18; LR, Drut, Braun in preparation\]](#)

$$\mathcal{Z} = \text{Tr}[e^{-\beta(\hat{H} - \mu_{\uparrow}\hat{N}_{\uparrow} - \mu_{\downarrow}\hat{N}_{\downarrow})}] = \text{Tr}[e^{-\beta(\hat{H} - \mu\hat{N} - h\hat{M})}]$$




density equation of state & polarization

- balanced gas: excellent agreement with experimental results and state-of-the-art numerical data
[\[experiment/BDMC: Van Houcke et al. '12; DHMC: Drut, Lähde, Wlazowski, Magierski '12\]](#)
- low temperatures: thermal wavelength λ_T increases, finite volume effects visible (currently $V = 11^3$)
- virial expansion (VE): valid at high temperature ($\beta\mu < 0$), provides an important benchmark for polarized systems
- CL shows excellent agreement with 3rd order VE for all polarizations studied


isothermal compressibility $\kappa = \frac{1}{n} \left(\frac{\partial n}{\partial P} \right)_{T,V,h}$

- agrees well with experiment & VE in the balanced case
[\[experiment: Ku et al. '12\]](#)
- peak shows weak dependence on polarization, suggests a flat phase-boundary near the balanced limit




spin susceptibility $\chi = \left(\frac{\partial m}{\partial h} \right)_{T,V,\mu}$

- functional form similar to non-interacting Fermi gas (Pauli susceptibility) but rescaled
- zero polarization: theoretical methods agree at high-temperature, experimental values disagree
[\[experiment: Sommer et al. '11; T-matrix: Pantel et al. '14, Palestini et al. '12; ETMA: Kashimura et al. '12; Luttinger-Ward: Enns et al. '12; AFQMC: Wlazowski et al. '13, Jensen et al. '18\]](#)
- low temperature: CL suggests absence of pseudogap, however, accuracy needs to be increased



shot noise for spin-and mass imbalanced systems: subleading pairing

- peak in pair-momentum distribution remains constant despite increasing mass imbalance, shot noise reveals emergence of subleading contribution
- mechanism: heavy component may be scattered from either deep inside (heavy majority, left) or far above (heavy minority, right) the Fermi surface




1D fermions (ground state): polarization & mass imbalance

[\[LR, Porter, Drut, Braun '17; LR, Drut, Braun '20\]](#)

$$\hat{H} = - \sum_{s=\uparrow,\downarrow} \int d^d x \hat{\psi}_s^\dagger(\vec{x}) \left(\frac{\hbar^2 \vec{\nabla}^2}{2m_s} \right) \hat{\psi}_s(\vec{x}) + g \int d^d x \hat{\psi}_\uparrow^\dagger(\vec{x}) \hat{\psi}_\uparrow(\vec{x}) \hat{\psi}_\downarrow^\dagger(\vec{x}) \hat{\psi}_\downarrow(\vec{x})$$

energy equation of state


- excellent agreement of GS energies with exact solution in the thermodynamic limit (Bethe Ansatz)
[\[BA: Iida, Wadati '07; Tracy, Widom '16\]](#)



on-site pair-correlation function:

$$\rho_{pair}(|x-x'|) = \langle \hat{\psi}_\uparrow^\dagger(x) \hat{\psi}_\downarrow^\dagger(x') \hat{\psi}_\downarrow(x') \hat{\psi}_\uparrow(x) \rangle$$

- off-center peak in pair-momentum distribution at $q = |k_F^\uparrow - k_F^\downarrow|$
- spatially oscillating "order parameter" (inhomogeneous pairing)



density-density correlations (momentum space):

$$n_{\uparrow\downarrow}(k, k') = \langle \hat{\psi}_{k\uparrow}^\dagger \hat{\psi}_{k\uparrow} \hat{\psi}_{k'\downarrow}^\dagger \hat{\psi}_{k'\downarrow} \rangle - \langle \hat{\psi}_{k\uparrow}^\dagger \hat{\psi}_{k\uparrow} \rangle \langle \hat{\psi}_{k'\downarrow}^\dagger \hat{\psi}_{k'\downarrow} \rangle$$

- clean signal of FFLO-type pairing at $(\pm k_F^\uparrow, \mp k_F^\downarrow)$

