





- deletions/duplications by low coverage massively parallel sequencing. *Prenatal diagnosis*, **33**(6), 584–590.
- Chiu, R. W., Chan, K. A., Gao, Y., Lau, V. Y., Zheng, W., Leung, T. Y., Foo, C. H., Xie, B., Tsui, N. B., Lun, F. M., *et al.* (2008). Noninvasive prenatal diagnosis of fetal chromosomal aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Proceedings of the National Academy of Sciences*, **105**(51), 20458–20463.
- Douglas, W. R., Langlois, S., and Johnson, J.-A. (2007). Mid-trimester amniocentesis fetal loss rate (committee opinion). *Journal of Obstetrics and Gynaecology Canada*, **29**(7), 586–590.
- Fan, H. C., Blumenfeld, Y. J., Chitkara, U., Hudgins, L., and Quake, S. R. (2008). Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proceedings of the National Academy of Sciences*, **105**(42), 16266–16271.
- Kitzman, J., Snyder, M., Ventura, M., Lewis, A., Qiu, R., Simmons, L., Gammill, H., *et al.* (2012). Noninvasive whole-genome sequencing of a human fetus. *Science Translational Medicine*, **4**(137), 137ra76–137ra76.
- Lo, Y., Corbetta, N., Chamberlain, P. F., Rai, V., Sargent, I. L., Redman, C. W., and Wainscoat, J. S. (1997). Presence of fetal DNA in maternal plasma and serum. *The Lancet*, **350**(9076), 485–487.
- Rampasek, L., Arbabi, A., and Brudno, M. (2014). Probabilistic method for detecting copy number variation in a fetal genome using maternal plasma sequencing. *In publication process, accepted for ISMB2014/Bioinformatics*.
- Srinivasan, A., Bianchi, D. W., Huang, H., Sehnert, A. J., and Rava, R. P. (2013). Noninvasive detection of fetal subchromosome abnormalities via deep sequencing of maternal plasma. *The American Journal of Human Genetics*, **92**(2), 167–176.