

1.5 Greenhouse gas effect

a) Calculate the unknown temperatures T_A and T_E .

Balance around area A:

$$-2\sigma T_A^4 + \sigma T_E^4 = 0 \quad (1.31)$$

Balance around area E:

$$\sigma T_A^4 - \sigma T_E^4 + (1 - \rho_{E,K})\dot{q}_S'' = 0 \quad (1.32)$$

Subtract the balances:

$$T_A = \left(\frac{(1 - \rho_{E,K})\dot{q}_S''}{\sigma} \right)^{1/4} = \left(\frac{0,7 \cdot 343,25 \text{ W m}^{-2}}{5,76 \cdot 10^{-8} \text{ W m}^{-2} \text{ K}^{-1}} \right)^{1/4} = 255,14 \text{ K} \quad (1.33)$$

$$T_E = (2 T_A^4)^{1/4} = 303,4 \text{ K} \quad (1.34)$$

1.6 Grilling pig

a) *Determine the surface temperature of the pig T_P in function of the given variables.*

The outer energy balance around the pig provides:

$$\dot{Q}_P = \dot{Q}_C \Phi_{CP} \quad (1.35)$$

The pig acts as a grey body, which gives the following surface brightness:

$$\dot{Q}_P = \varepsilon_P \sigma A_P T_P^4 + (1 - \varepsilon_P) (\dot{Q}_C \Phi_{CP}) \quad (1.36)$$

The coal is a black radiator:

$$\dot{Q}_C = \sigma A_C T_C^4 \quad (1.37)$$

Inserting the surface brightness yields:

$$\varepsilon_P \sigma A_P T_P^4 + (1 - \varepsilon_P) (\dot{Q}_C \Phi_{CP}) = \Phi_{CP} \dot{Q}_C \quad (1.38)$$

The equation corresponds to an inner balance, which can be established directly.

$$\Longleftrightarrow \sigma A_P \varepsilon_P T_P^4 = \sigma A_C \varepsilon_C T_C^4 \Phi_{CP} \Longleftrightarrow T_P^4 = \frac{A_C \Phi_{CP}}{A_P} T_C^4 \quad (1.39)$$

Using the reciprocity gives:

$$T_P = T_C \Phi_{PC}^{\frac{1}{4}} \quad (1.40)$$

1.7 Radiation within a wedge-shaped opening

a) Determine the view factors $\Phi_{1,2}, \Phi_{2,1}, \Phi_{1,o}, \Phi_{2,o}$.

Sum rule (where $\Phi_{1,1} = \Phi_{2,2} = \Phi_{o,o} = 0$):

$$\Phi_{1,2} + \Phi_{1,\ddot{o}} = 1 \quad (1.41)$$

$$\Phi_{2,2} + \Phi_{2,\ddot{o}} = 1 \quad (1.42)$$

$$\Phi_{\ddot{o},1} + \Phi_{\ddot{o},2} = 1 \quad (1.43)$$

Reciprocity rule:

$$a \cdot \Phi_{1,2} = a\sqrt{2} \cdot \Phi_{2,1} \quad (1.44)$$

$$a \cdot \Phi_{1,\ddot{o}} = a \cdot \Phi_{\ddot{o},1} \quad (1.45)$$

$$a \cdot \Phi_{\ddot{o},2} = a\sqrt{2} \cdot \Phi_{2,\ddot{o}} \quad (1.46)$$

Given Equations 1.41 - 1.46, we have 6 unknowns and 6 equations. Rearranging, substituting and filling in numerical values results in:

$$\Phi_{1,2} = \frac{1}{2} \cdot \sqrt{2} \quad (1.47)$$

$$\Phi_{1,\ddot{o}} = 1 - \frac{1}{2} \cdot \sqrt{2} \quad (1.48)$$


$$\Phi_{2,1} = \frac{1}{2} \quad (1.49)$$

$$\Phi_{2,\ddot{o}} = \frac{1}{2} \quad (1.50)$$

$$\Phi_{\ddot{o},1} = 1 - \frac{1}{2} \cdot \sqrt{2} \quad (1.51)$$

$$\Phi_{\ddot{o},2} = \frac{1}{2} \cdot \sqrt{2} \quad (1.52)$$

b) Determine the energy loss of surface (1) $\dot{q}'_{1,V}$ and the opening $\dot{q}'_{\ddot{o},V}$ for a unit length of the opening.



For the opening:

$$\tau_{\ddot{o}} = 1 \Rightarrow \epsilon_{\ddot{o}} = \rho_{\ddot{o}} = 0 \quad (1.53)$$

Surface brightness of the opening:

$$\dot{Q}_{\ddot{o},a} = \dot{Q}_1 \Phi_{1,\ddot{o}} + \dot{Q}_2 \Phi_{2,\ddot{o}} \quad (1.54)$$

For body 1 (black body):


$$\tau_1 = 0 \text{ and } \epsilon_1 = \alpha_1 = 1 \quad (1.55)$$

Surface brightness of body 1:

$$\dot{Q}_1 = A_1 \sigma T_1^4 \quad (1.56)$$

Surface brightness of body 2:

$$\dot{Q}_2 = A_2 \sigma T_2^4 \epsilon_2 + \dot{Q}_1 \Phi_{12} \rho_2 \quad (1.57)$$



Energy balance around body 2:

$$\dot{Q}_1\Phi_{12} + \dot{Q}_{\ddot{o}}\Phi_{\ddot{o}2} - \dot{Q}_2 = 0 \quad (1.58)$$

Rate of heat transfer from the opening to the environment:

$$\dot{Q}_{\ddot{o},a} = \dot{Q}_1\Phi_{1\ddot{o}} + \dot{Q}_2\Phi_{2\ddot{o}} \quad (1.59)$$

Substitution of \dot{Q}_2 (Equation 1.58) into Equation 1.59 and dividing by L results in:

$$\begin{aligned} \dot{q}'_{\ddot{o},a} &= \dot{q}'_1 (\Phi_{1\ddot{o}} + \Phi_{12}\Phi_{2\ddot{o}}) = a\sigma T_1^4 (\Phi_{1\ddot{o}} + \Phi_{12}\Phi_{2\ddot{o}}) = \\ 0.3 \cdot 5.67 \text{ [m]} \cdot 10^{-8} \text{ [W/m}^2\text{K}^4] 1000^4 \text{ [K]} &\left(1 - \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{4}\right) = 11 \text{ [kW/m]} \end{aligned} \quad (1.60)$$

c) Determine the temperature T_2 of surface (2)

Substitution of Equation 1.57 into 1.58 results in:

$$\dot{Q}_1\Phi_{12} = A_2\sigma T_2^4\epsilon_2 + \dot{Q}_1\Phi_{12}\rho_2 \quad (1.61)$$

Rearranging (note that from Kirchhoff and the fact that we are dealing with a grey body it results that $(1 - \rho_2) = \alpha_2 = \epsilon_2$):

$$T_2 = \sqrt[4]{\frac{\dot{Q}_1\Phi_{12}(1 - \rho_2)}{A_2\sigma\epsilon_2}} = \sqrt[4]{\frac{T_1^4\Phi_{12}}{\sqrt{2}}} = \sqrt[4]{\frac{1000^4}{2}} = 840.9 \text{ [K]} \quad (1.62)$$