


# Solutions lecture 4

## 4.1 Cooling of a hot water pipe

### Analysis

In this situation, natural convection will occur. Given data are:  $T_s = 70^\circ\text{C}$ ,  $T_\infty = 18^\circ\text{C}$ , length is 6 metres and the diameter 8.0 cm. A sketch is presented below:



### Approach

#### Assumptions

#### Route to solution

This natural convection problem can be solved by using the following steps:

1. Determine the average temperature.
2. Determine the Grashof number.
3. Determine the Rayleigh number.
4. Choose the right correlation based on geometry and the Rayleigh number.
5. Determine the Nusselt number.
6. Derive the value of  $h$ .
7. Substitute in Newton's cooling law.

### Elaboration

We start with determining the average temperature:

$$T_f = \frac{T_s + T_\infty}{2} = \frac{70 + 18}{2} = \frac{88}{2} = 44^\circ\text{C}$$

For natural convection, the Grashof number needs to be calculated:

$$\text{Gr} = \frac{g\beta(T_s - T_\infty)L_c^3}{\nu^2}$$

where  $9.81 \text{ m s}^{-2}$  is the gravitational constant,  $\beta$  is the thermal expansion coefficient:

$$\beta = \frac{2}{T_s + T_\infty} = \frac{2}{342.15 + 291.15} = 3.15 \times 10^{-3} \text{ K}^{-1}$$

Note that the temperatures need to be substituted in **Kelvin**, not Celsius. Furthermore,  $D$  is the aforementioned diameter, and  $\nu$  is the kinematic viscosity. The value for this is taken at 45 °C, to be  $1.750 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Substituting these values:

$$\text{Gr} = \frac{g\beta(T_s - T_\infty)D^3}{\nu^2} = \frac{9.81 \cdot (3.15 \cdot 10^{-3}) \cdot (70 - 18) \cdot 0.08^3}{(1.750 \cdot 10^{-5})^2} = 2.67 \cdot 10^6 [-]$$

At 45 °C, the Prandtl number is  $\text{Pr} = 0.7241[-]$ . With this, the Rayleigh number can be determined:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = 2.67 \cdot 10^6 \cdot 0.7241 = 1.94 \cdot 10^6 [-]$$

Now, since the pipe is horizontal and cylindrical, we can use the following correlation:

$$\text{Nu} = \left( 0.6 + \frac{0.387 \cdot \text{Ra}_D^{\frac{1}{4}}}{\left( 1 + \left( \frac{0.559}{\text{Pr}} \right)^{\frac{9}{16}} \right)^{\frac{8}{27}}} \right)^2$$

Substitution of all variables:

$$\text{Nu} = \left( 0.6 + \frac{0.387 \cdot (1.94 \cdot 10^6)^{\frac{1}{4}}}{\left( 1 + \left( \frac{0.559}{0.7241} \right)^{\frac{9}{16}} \right)^{\frac{8}{27}}} \right)^2 = 17.58 [-]$$

The Nusselt number is defined as:

$$\text{Nu} = \frac{hD}{k}$$

At 45 °C, the value of  $k$  is  $k = 0.02699 \text{ W m}^{-1} \text{ K}^{-1}$ . Rewriting and substituting gives:

$$h = \frac{\text{Nu}_D k}{D} = \frac{17.58 \cdot 0.02699}{0.08} = 5.93 \text{ W m}^{-2} \text{ K}^{-1}$$

We can now substitute all values in Newton's cooling law:

$$\dot{Q} = hA\Delta T$$

where the area is  $A = \pi DL = \pi \cdot 0.08 \cdot 6 = 1.51 \text{ m}^2$

$$\dot{Q} = h \cdot A(T_s - T_\infty) = 5.93 \cdot 1.51 \cdot (70 - 18) = 466 \text{ W}$$

## Evaluation

Check your answer:

- Does the answer have the correct dimensions?
- Is the answer in the right order of magnitude?
- Did you answer all the questions that were asked?

## 4.2 Convection of heat from a coffee machine

### Analysis

This plate loses 90 W of heat, of which is 52.4% due to radiation and the other 47.6% due to natural convection. The situation is sketched below.




Figure 4.1: Heat loss of a coffee machine plate

This gives a total of  $0.476 \cdot 90 = 42.84$  W of natural convection. For natural convection, the following relation is valid:

$$\dot{Q} = hA\Delta T$$

However,  $h$  is dependent on the average temperature, which is subsequently dependent on the temperature  $T_s$  of the plate, considering the fact that this is codependent for the film temperature of which the value will be based

$$T_f = \frac{T_s + T_\infty}{2}$$

We need to determine the temperature by means of iteration. The maximum temperature of the plate is given, 250 °C. A first initial guess for the temperature of the plate is 180 °C.

### Approach

#### Assumptions

#### Route to solution

1. Determine the average temperature

2. Determine the Grashof number
3. Determine the Rayleigh number
4. Choose the right correlation based on geometry and the Rayleigh number
5. Determine the Nusselt number
6. Derive the value of  $h$
7. Substitute in Newton's cooling law
8. If necessary, iterate

## Elaboration

The average temperature is:

$$T_f = \frac{T_s + T_\infty}{2} = \frac{180 + 20}{2} = 100^\circ\text{C}$$

The Grashof number is defined as:

$$\text{Gr} = \frac{g\beta(T_s - T_\infty)L_c^3}{\nu^2}$$

in which  $g = 9.81 \text{ m s}^{-2}$  is the gravitational constant,  $\beta$  is the thermal expansion coefficient:

$$\beta = \frac{1}{T_f} = \frac{1}{100 + 273} = 2.68 \times 10^{-3} \text{ K}^{-1}$$

Note that the temperatures in  $\beta$  has the unit of Kelvin, not Celsius. Furthermore  $L_c$  is the characteristic length, which, for a horizontal cylindrical flat plate with diameter  $D$ , is

$$L_c = \frac{A_s}{p} = \frac{\frac{\pi}{4}D^2}{\pi D} = \frac{D}{4} = 0.04\text{m}$$

At  $100^\circ\text{C}$ , air has the following properties:  $k = 0.03095 \text{ W m}^{-1} \text{ K}^{-1}$  and  $\nu = 2.306 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Substituting all these values give:

$$\text{Gr} = \frac{9.81 \cdot (5.38 \cdot 10^{-3} \cdot (180 - 20) \cdot 0.04^3)}{(2.306 \cdot 10^{-5})^2} = 5.063 \cdot 10^5 [-]$$

At  $100^\circ\text{C}$ , the Prandtl number is  $\text{Pr} = 0.7111 [-]$ . With this, the Rayleigh number can be determined to be:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = 5.063 \cdot 10^5 \cdot 0.7111 = 3.60 \cdot 10^5 [-]$$

Now, the situation concerns an upper part of a horizontal hot flat plate with  $10^4 < \text{Ra} < 10^7$ . For this, the Nusselt number is

$$\text{Nu} = 0.54 \text{Ra}^{\frac{1}{4}} = 0.54 \cdot (3.60 \cdot 10^5)^{\frac{1}{4}} = 13.2 [-]$$

The Nusselt number is defined as

$$\text{Nu} = \frac{hL_c}{k}$$

Substitution of all variables gives

$$h = \frac{\text{Nu} \cdot k}{L_c} = \frac{13.2 \cdot 0.03095}{0.04} = 10.2 \text{ W m}^{-2} \text{ K}^{-1}$$

Substitution of this result into Newton's cooling law, where the surface is  $A = \frac{\pi}{4}D^2$ :

$$\dot{Q} = hA\Delta T = 12.1 \cdot \frac{\pi}{4} \cdot 0.16^2 \cdot (180 - 20) = 32.8 \text{ W}$$

As stated earlier, this 32.8 W is not in accordance with the 42.84 W we calculated earlier. This means that the plate will be hotter than  $180^\circ\text{C}$ . This means we need to reiterate.

Let's say that the plate is  $220^\circ\text{C}$ . The average temperature is then:

$$T_f = \frac{T_s + T_\infty}{2} = \frac{220 + 20}{2} = 120^\circ\text{C}$$

The Grashof number is defined as:

$$\text{Gr} = \frac{g\beta(T_s - T_\infty)L_c^3}{\nu^2}$$

in which  $g = 9.81 \text{ m s}^{-2}$  is the gravitational constant,  $\beta$  is the thermal expansion coefficient:

$$\beta = \frac{1}{T_f} = \frac{1}{120 + 273} = 2.54 \times 10^{-3} \text{ K}^{-1}$$

At 120 °C, air has the following properties:  $k = 0.03235 \text{ W m}^{-1} \text{ K}^{-1}$  and  $\nu = 2.522 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Substituting all these values give:

$$\text{Gr} = \frac{9.81 \cdot (2.54 \cdot 10^{-3}) \cdot (220 - 20) \cdot 0.04^3}{(2.522 \cdot 10^{-5})^2} = 5.014 \cdot 10^5 [-]$$

At 120 °C, the Prandtl number is  $\text{Pr} = 0.7073[-]$ . With this, the Rayleigh number can be determined to be:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = 5.014 \cdot 10^5 \cdot 0.7073 = 3.55 \cdot 10^5 [-]$$

Now, the situation concerns an upper part of a horizontal hot flat plate with  $10^4 < \text{Ra} < 10^7$ . For this, the Nusselt number is

$$\text{Nu} = 0.54 \text{Ra}^{\frac{1}{4}} = 0.54 \cdot (3.55 \cdot 10^5)^{\frac{1}{4}} = 13.2 [-]$$

The Nusselt number is defined as

$$\text{Nu} = \frac{hL_c}{k}$$

Substitution of all variables gives

$$h = \frac{\text{Nu} \cdot k}{L_c} = \frac{13.2 \cdot 0.03235}{0.04} = 10.7 \text{ W m}^{-2} \text{ K}^{-1}$$

Substitution of this result into Newton's cooling law:

$$\dot{Q} = hA\Delta T = 10.7 \cdot \frac{\pi}{4} \cdot 0.16^2 \cdot (220 - 20) = 42.9 \text{ W}$$

This value is quite in accordance with the earlier presented 42.84 W. This indicates that the plate will be around 220 °C.

## Evaluation

Check your answer:

- Does the answer have the correct dimensions?
- Is the answer in the right order of magnitude?
- Did you answer all the questions that were asked?

## 4.3 Heat convection parameters

For forced convection, it was provided in lecture 3, that  $Nu$  is a function of  $Re$  and  $Pr$ :

$$Nu = f(Re, Pr)$$

$$\frac{hL_c}{k} = f\left(\frac{\rho UL_c}{\mu}, Pr\right)$$

This means that the group of parameters in which the heat transfer coefficient  $h$  is processed, is a function of  $\rho, U, L_c, \mu$  and  $Pr$ . We can rewrite this to:

$$h = f(\rho, U, L_c, \mu, Pr, k)$$


These variables will be discussed from here:

- Density  $\rho$ : Combined with the velocity  $U$ , the density forms the impulse of the fluid. The impulse can be seen as an indicator for the impact force of the oncoming particles. Is this value high, the fluid particles just above the surface will be hardly affected by the surface they are flowing above and by the particles close to the surface area. The particles close to the surface will stagnate by the surface and by the viscosity and will affect particles further away from the surface.

Having a high density will result in a high impulse and therefore the boundary layer will grow less fast and remain thinner. A small boundary layer means a small distance for the heat to cover and subsequently an easier heat transfer.  $h$  will increase if the density increases.

At a certain point the impulse will be that much larger than the viscous effects that the laminar flow will transform into turbulent flow. In that case the  $h$  will increase drastically as the gradient close to the surface will be a lot larger.

For a turbulent flow, looking very close to the surface a very small boundary layer will appear, meaning a large value of  $h$ . It can be concluded that a turbulent flow results in a larger  $h$ .



- Velocity  $U$  : Just like the density, the velocity is determining the impulse ( $\rho U$ ). So all statements for above are valid as well if  $\rho$  is replaced by  $U$ .
- Characteristic length  $L_c$ : the larger the surface area, the more the boundary layer has opportunity to grow. At the end of the characteristic length, the boundary layer will be thick with locally low  $h$  values, especially compared with the leading edge. The larger the  $L_c$ , the smaller the average value of  $h$ .
- Dynamic viscosity  $\mu$ : the dynamic viscosity ensures the no slip boundary condition at the surface is felt in the fluid above the surface. The higher the viscosity, the larger the influence and the more thick the boundary layer will grow. The viscosity has an inverted effect compared to the density and the velocity. This is also quite obvious if the definition of the Reynolds number is considered.

If the dynamic viscosity is high enough, the impulse will be dominated by the viscous effects and this will result in a laminar flow. The Reynolds number will be under the critical value for that specific geometry.

- Prandtl number  $Pr$ : the Prandtl number indicates how, given a certain velocity boundary layer (as well present without thermal effects), the thermal boundary layer will look like as a temperature difference is present between the surface and the free flow. A thermal boundary layer larger than ( $Pr < 1$ ), thinner than ( $Pr > 1$ ) or as thick as ( $Pr = 1$ ) the velocity boundary layer.

If in a flow the Prandtl number is enhanced, without altering velocity profile, the thermal boundary layer will get thinner. Using the statement at the density variable, the value of  $h$  will increase.

- Thermal conductivity  $k$ : If a fluid conducts heat better, the value of  $k$  will be higher. Convection can only be present if the particles are able to conduct heat. Convection can be stated as conduction being enhanced by the continuous supply of fresh particles and the take away of old particles. Therefore, if you calculate the convective heat transfer, the conductive heat transfer is already included and there is no need to calculate the conductive heat transfer independently.

For natural convection, the following correlations are valid:

$$\text{Nu} = f(\text{Ra}) = g(\text{Gr}, \text{Pr})$$

$$\frac{hL_c}{k} = f(g, \beta, (T_s - T_\infty), L_c, \nu, \text{Pr}, k)$$

Giving for  $h$ :

$$h = f(g, \beta, (T_s - T_\infty), L_c, \nu, \text{Pr}, k)$$

The gravitational constant  $g$  is always the same, and therefore has no influence on  $h$ .  $L_c$ ,  $\text{Pr}$  and  $k$  have the same effect as in the case of forced convection. The temperature difference can be seen as the driver behind the phenomenon and therefore fulfills the same role as  $U$  in the case of forced convection. A higher value will result in a higher value for  $h$ .

- Thermal expansion coefficient  $\beta$ : This coefficient is larger the more a fluid expands with a certain increase in temperature. The larger the expansion, the more the decrease in density and therefore a larger flow velocity and a higher value for  $h$ .
- Kinematic viscosity  $\nu$ : the kinematic viscosity is the dynamic viscosity divided by the density. Therefore it has the same effect as the effects of dynamic viscosity and density.

A last important remark is the influence of the geometry on the value of  $h$ . The geometry determines how the flow pattern will look like and which correlation is valid. Also, the roughness of the surface plays a large role: the rougher the surface the quicker the transition from laminar to turbulent flow.

## 4.4 Light bulb temperature

### Analysis

Consider a 25 W lightbulb with a light-efficiency of 10 %. The lightbulb has a diameter of 8.0 cm, and an outside temperature of 25 °C. When assuming all heat is lost due to natural convection, determine the surface temperature of the lightbulb.

### Approach

#### Assumptions

- Only natural convection
- Light bulb can be modelled as a sphere

#### Route to solution

1. Determine the average temperature
2. Determine the Grashof number
3. Determine the Rayleigh number
4. Choose the right correlation based on geometry and the Rayleigh number
5. Determine the Nusselt number
6. Derive the value of  $h$
7. Substitute in Newton's cooling law
8. If necessary, iterate

### Elaboration

We take an initial guess for the surface temperature,  $T_s = 100$  °C. The average temperature is:

$$T_f = \frac{T_s + T_\infty}{2} = \frac{100 + 25}{2} = 62.5 \text{ °C}$$

The Grashof number is defined as:

$$\text{Gr} = \frac{g\beta(T_s - T_\infty)L_c^3}{\nu^2}$$

in which  $g = 9.81 \text{ m s}^{-2}$  is the gravitational constant,  $\beta$  is the thermal expansion coefficient:

$$\beta = \frac{2}{T_f} = \frac{2}{298 + 373} = 2.98 \times 10^{-3} \text{ K}^{-1}$$

Note that the temperatures in  $\beta$  has the unit of Kelvin, not Celsius.

At 60 °C, air has the following properties:  $k = 0.02808 \text{ W m}^{-1} \text{ K}^{-1}$  and  $\nu = 1.896 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Substituting all these values give:

$$\text{Gr} = \frac{9.81 \cdot (2.98 \cdot 10^{-3}) \cdot (100 - 25) \cdot 0.08^3}{(1.896 \cdot 10^{-5})^2} = 3.12 \cdot 10^6 [-]$$

At 60 °C, the Prandtl number is  $\text{Pr} = 0.7202[-]$ . With this, the Rayleigh number can be determined to be:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = 3.12 \cdot 10^6 \cdot 0.7202 = 2.25 \cdot 10^6 [-]$$

Now, the situation concerns a sphere with  $\text{Ra}_D \leq 10^{11}$ , where  $\text{Pr} \geq 0.7$ . For this, the Nusselt number is:

$$\text{Nu} = 2 + \frac{0.589\text{Ra}^{1/4}}{[1 + (0.469/\text{Pr})^{9/16}]^{4/9}} = 19.6$$



The Nusselt number is defined as

$$\text{Nu} = \frac{hL_c}{k}[-]$$

Substitution of all variables gives

$$h = \frac{\text{Nu} \cdot k}{D} = \frac{19.6 \cdot 0.02808}{0.08} = 6.89 \text{ W m}^{-2} \text{ K}^{-1}$$

Substitution of this result into Newton's cooling law, where the surface is  $A = \pi D^2$ :

$$\dot{Q} = hA\Delta T = 6.89 \cdot \pi \cdot D^2 \cdot (100 - 25) = 10.4 \text{ W}$$

As stated earlier, this 10.4 W is not in accordance with the 22.5 W we need to have earlier. This means that the surface is hotter than 100°C. Reverse calculation shows a  $\Delta T$  of 162 °C. After some iterations, a value of  $T_s = 168^\circ\text{C}$  is found. With this value, the following results are obtained:

$$T_f = 96.5^\circ\text{C} \approx 100^\circ\text{C}$$

$$\beta = 2.7 \times 10^{-3} \text{ K}^{-1}$$

At 100 °C, air has the following properties:  $k = 0.03095 \text{ W m}^{-1} \text{ K}^{-1}$ ,  $\nu = 2.306 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$  and  $\text{Pr} = 0.7202[-]$ .

$$\text{Gr} = 3.66 \cdot 10^6[-]$$

$$\text{Ra} = 2.60 \cdot 10^6[-]$$

$$\text{Nu} = 20.25[-]$$

$$h = 7.84 \text{ W m}^{-2} \text{ K}^{-1}$$

$$\dot{Q} = 22.53 \text{ W}$$


$$T = 167.8^\circ\text{C}$$

## 4.5 Cooling of a head - Hand-in

- a) Determine the average surface temperature of the head.

To determine the rate of heat loss by convection, a surface temperature for the head should be determined.

The thermal resistance network looks as follows:



The skin temperature  $T_s$  can be determined by rewriting the relationship for the thermal resistance theorem. To do so, first  $R_{\text{conv}}$  should be determined.

It will be assumed for now that the skin  $T_s$  will be approximately  $33^\circ\text{C}$ . Which yields a film temperature of:

$$T_f = \frac{T_s + T_{\text{amb}}}{2} = \frac{33 + 21}{2} = 27^\circ\text{C}$$

Taking the air properties at  $27^\circ\text{C}$  (which are approximately equal to those of  $25^\circ\text{C}$  from the table):

$$k = 0.02551 \text{ W/mK}$$

$$\nu = 1.562 \cdot 10^{-5} \text{ m}^2/\text{s}$$

$$\text{Pr} = 0.7296$$

$$\beta = \frac{1}{T_f} = \frac{1}{273 + 27} = 0.0033 \text{ K}^{-1}$$

The characteristic length in this case is equal to the diameter of the head, which yields:

$$L_c = d = 0.3 \text{ m}$$

Now the Grashof number can be calculated:

$$\text{Gr} = \frac{g \cdot \beta \cdot (T_s - T_{\text{amb}}) \cdot L_c^3}{\nu^2} = \frac{9.81 \cdot 0.0033 \cdot (33 - 21) \cdot 0.3^3}{(1.562 \cdot 10^{-5})^2} = 4.3402 \cdot 10^7$$

With this known, the Rayleigh number yields:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = (4.3402 \cdot 10^7) \cdot (0.7296) = 3.1666 \cdot 10^7$$

Using the Nusselt correlation valid for natural convection around spheres gives:

$$\text{Nu} = 2 + \frac{0.589 \cdot \text{Ra}^{1/4}}{\left[1 + \left(\frac{0.469}{\text{Pr}}\right)^{9/16}\right]^{4/9}} = 2 + \frac{0.589 \cdot (3.1666 \cdot 10^7)^{1/4}}{\left[1 + \left(\frac{0.469}{0.7296}\right)^{9/16}\right]^{4/9}} = 36.2$$

Using the definition of the Nusselt number to find the average heat transfer coefficient:

$$h = \frac{\text{Nu} \cdot k}{L_c} = \frac{36.2 \cdot 0.02551}{0.3} = 3.08 \text{ W/m}^2\text{K}$$

Which yields the convective resistance:


$$R_{\text{conv}} = \frac{1}{h \cdot A} = \frac{1}{3.08 \cdot \pi 0.3^2} = 1.15 \text{ K/W}$$

From the thermal resistance theorem, we can check whether our guess of  $T_s = 33^\circ\text{C}$  was indeed correct:

$$\dot{Q} = \frac{T_s - T_{\text{amb}}}{R_{\text{conv}}} = \frac{33 - 21}{1.15} = 10.4 \text{ W}$$

- b) Determine again the average surface temperature of the head while standing still. To determine the rate of heat loss by convection, a surface temperature for the head should be determined.

The thermal resistance network looks as follows:



The skin temperature  $T_s$  can be determined by rewriting the relationship for the thermal resistance theorem. To do so, first  $R_{\text{cond}}$  and  $R_{\text{conv}}$  should be determined.

The thermal resistance can be approximated to be a flat plate as  $x \ll d$ , thus:

$$R_{\text{cond}} = \frac{x}{k \cdot A} = \frac{0.005}{0.03 \cdot \pi \cdot 0.3^2} = 0.6932 \text{ K/W}$$

It will be assumed for now that the temperature of the outer surface of the balaclava  $T_i$  will be approximately  $0^\circ\text{C}$ . Which yields a film temperature of:

$$T_f = \frac{T_i + T_{\text{amb}}}{2} = \frac{0 + -10}{2} = -5^\circ\text{C}$$

Taking the air properties at  $-5^\circ\text{C}$ , the values were taken from engineersedge.com since the slides do not include them, you could've taken the properties of air at  $20^\circ\text{C}$ :

$$k = 0.02364 \text{ W/mK}$$

$$\nu = 1.295 \cdot 10^{-5} \text{ m}^2/\text{s}$$

$$\text{Pr} = 0.7375$$

$$\beta = \frac{1}{T_f} = \frac{1}{273 + -5} = 0.00373 \text{ K}^{-1}$$

The characteristic length in this case is equal to sum of the diameter of the head and twice the thickness of the balaclava hat, which yields:

$$L_c = d + 2x = 0.31 \text{ m}$$

Now the Grashof number can be calculated:

$$\text{Gr} = \frac{g \cdot \beta \cdot (T_s - T_{\text{amb}}) \cdot L_c^3}{\nu^2} = \frac{9.81 \cdot 0.00373 \cdot (-5 + 10) \cdot 0.31^3}{(1.295 \cdot 10^{-5})^2} = 1.048 \cdot 10^8$$

With this known, the Rayleigh number yields:

$$\text{Ra} = \text{Gr} \cdot \text{Pr} = (1.048 \cdot 10^8) \cdot (0.7375) = 7.732 \cdot 10^7$$

Using the Nusselt correlation valid for natural convection around spheres gives:

$$\text{Nu} = 2 + \frac{0.589 \cdot \text{Ra}^{1/4}}{\left[1 + \left(\frac{0.469}{\text{Pr}}\right)^{9/16}\right]^{4/9}} = 2 + \frac{0.589 \cdot (7.732 \cdot 10^7)^{1/4}}{\left[1 + \left(\frac{0.469}{0.7375}\right)^{9/16}\right]^{4/9}} = 50.6$$

Using the definition of the Nusselt number to find the average heat transfer coefficient:

$$h = \frac{\text{Nu} \cdot k}{L_c} = \frac{50.6 \cdot 0.02364}{0.31} = 3.86 \text{ W/m}^2\text{K}$$

Which yields the convective resistance:

$$R_{\text{conv}} = \frac{1}{h \cdot A} = \frac{1}{3.86 \cdot \pi \cdot 0.31^2} = 0.8581 \text{ K/W}$$

From the thermal resistance theorem, we can check whether our guess of  $T_s = 0\text{ }^{\circ}\text{C}$  was indeed correct:

$$\dot{Q} = \frac{T_s - T_{\text{amb}}}{R_{\text{cond}} + R_{\text{conv}}} = \frac{0 - -10}{0.6932 + 0.8581} = 6.45\text{ W}$$

The calculated  $\dot{Q}$  lies close to the actual value (7.5% of 100 W), which makes  $0\text{ }^{\circ}\text{C}$  a good guess. A closer answer would be:

$$T_s = \dot{Q} \cdot (R_{\text{cond}} + R_{\text{conv}}) + T_{\text{amb}} = 7.5 \cdot (0.6932 + 0.8581) - 10 = 1.63\text{ }^{\circ}\text{C}$$

Since the new guess is really close to the guess of  $0\text{ }^{\circ}\text{C}$ , and therefore the properties of air will not change much, the answer of  $1.6\text{ }^{\circ}\text{C}$  is satisfactory.

- c) Sketch the temperature profile showing how the temperature changes from the center of the head to the surrounding cold air.

