


# Falling Marbles



Marbles fall from rest through an opening at a steady rate of three per second. Determine the vertical separation  $h$  of two consecutive marbles when the lower one has dropped 7 meters. Neglect air resistance and take  $g = 10 \text{ m}^2$ .

*Using known expressions:*

$$a = \frac{dv}{dt} \Rightarrow dv = a dt \quad (1)$$

$$\int_{v_0}^v dv = a \int_0^t dt \quad (2)$$

$$v(t) = a \cdot t + v_0 \quad (3)$$

$$v = \frac{ds}{dt} \Rightarrow ds = v dt = (v_0 + at) dt \quad (4)$$

$$\int_{s_0}^s ds = \int_0^t (v_0 + at) dt \quad (5)$$

$$s(t) = \frac{1}{2}a \cdot t^2 + v_0 \cdot t + s_0 \quad (6)$$

*Given:*

Distance:  $h = 7m$

Gravitational acceleration:  $g = 10m/s^2$

The marbles fall from rest, thus  $v_0 = 0$ . Seen from the top  $s_0 = 0$ , thus Equation 6 simplifies to:

$$s(t) = \frac{1}{2}a \cdot t^2 + v_0 \cdot t + s_0 \Rightarrow s(t) = \frac{1}{2}a \cdot t^2 \quad (7)$$

Where  $a = -g = -10m/s^2$  (since it points in the negative y-direction). The time it takes for the ball to drop 7m ( $s = -7$ ) can be calculated using:

$$s(t) = \frac{1}{2}a \cdot t^2 \Rightarrow t = \sqrt{\frac{2 \cdot s}{a}} = \sqrt{\frac{2 \cdot -7}{-10}} \approx 1.18s \quad (8)$$

Three balls are dropped every second, thus every  $\frac{1}{3}s$  a ball is dropped. It took 1.18 seconds for the first ball to reach 7m, and after  $\frac{1}{3}s$  a second ball was dropped. The distance the second ball travelled is thus:

$$s(t) = \frac{1}{2}a \cdot t^2 \Rightarrow s(t_1 - t_2) = \frac{1}{2} \cdot -10 \cdot (1.18 - \frac{1}{3})^2 = -3.61m \quad (9)$$

Thus the distance between two consecutive marbles is  $h = -3.61 - -7 = 3.39m$