




## EB - Cond. - IE 7

Derive the energy balance and boundary conditions required to calculate the one-dimensional radial steady-state temperature profile inside the spherical wall.

### 1 Setting up the balance:

To derive the one-dimensional steady-state temperature profile, an energy balance around an infinitesimal element is needed. Heat is conducted in and out of the element.



Hence, the steady-state balance reads:

$$\dot{Q}_{r,in} - \dot{Q}_{r,out} = 0,$$

the sum of the in- and outgoing fluxes should equal zero, because of steady-state conditions.

### 2 Defining the elements within the balance:

The ingoing flux described by use of Fourier's law:

$$\dot{Q}_{r,in} = -\lambda \cdot 4\pi r^2 \cdot \frac{\partial T}{\partial r},$$

and the outgoing flux is approximated by the use of the Taylor series expansion.

$$\begin{aligned}\dot{Q}_{r,out} &= \dot{Q}_{r,in} + \frac{\partial \dot{Q}_{r,in}}{\partial r} \cdot dr \\ &= -\lambda \cdot 4\pi r^2 \cdot \frac{\partial T}{\partial r} + \frac{\partial}{\partial r} \left( -\lambda \cdot 4\pi r^2 \cdot \frac{\partial T}{\partial r} \right) \cdot dr.\end{aligned}$$

### 3 Inserting and rearranging:

$$\frac{\partial}{\partial r} \left( r^2 \frac{\partial T}{\partial r} \right) = 0.$$





## 4 Defining the boundary and/or initial conditions:

The boundary conditions yield from the given temperatures

$$T(r = r_1) = T_1,$$

and:

$$T(r = r_2) = T_2.$$