Relative arithmetic content in symmetric odd-zeta forms

Exact reductions, higher-weight decay, and a conditional irrationality criterion

28 October 2026

Abstract

We separate analytic decay from integral normalization in symmetric Padé forms for odd zeta values. Smith reduction gives an exact formula for the content of the target and rational rows on the cancellation lattice. A direct kernel comparison extends the weight-five decay estimate to every fixed higher weight. An elementary logarithm inequality gives a uniform exponential margin greater than one third for weights at least seven. Consequently, nonzero integral forms with subexponential relative arithmetic cost imply irrationality. The Lean companion proves this conditional implication, including eventual bounds and arbitrary unbounded indexing, and an exact integral-projector divisibility criterion. Signed combinations are also allowed. A target-eliminating cofactor combination gives an integer lower bound that rules out an overstrong cofactor-norm saving proposed in the supplied note. Individual primitive normalization gives a sharper signed-pair criterion, using each form's own coefficient gcd instead of the common content. The construction-specific relative-cost bound and the existence of suitable positive integral cancellation vectors in every higher weight are hypotheses, not conclusions. This paper does not prove the unconditional irrationality of every odd zeta value at least seven, or of $\zeta(5)$.

1 Construction and the arithmetic quantity

Fix $s = 2r + 1 \geq 5$. Let $n \geq 2r$ be even and select $m_j = n - j$, $0 \leq j \leq r$. Write

$$d_n = \operatorname{lcm}(1, \ldots, n), \qquad T_{s,n,m}(t) = \frac{(n!)^s}{(m!)^2} \frac{\prod_{v=1}^{m}(t - v)(t + n + v)}{\prod_{h=0}^{n}(t + h)^s}.$$

The convergent positive sums are

$$\mathcal{A}_{s,n,m} = \sum_{k=m+1}^{\infty} T_{s,n,m}(k).$$

Every summand is positive in this range. Partial fractions, reflection $t \mapsto -n - t$, and the vanishing total simple-pole residue give

$$\mathcal{A}_{s,n,m} = b_{n,m} + \sum_{q=1}^{r} a_{q,n,m}\zeta(2q + 1). \tag{1}$$

The coefficients are rational. Weighted Taylor expansion at each pole shows

$$d_n^{s-(2q+1)} a_{q,n,m} \in \mathbb{Z}, \qquad d_n^s b_{n,m} \in \mathbb{Z}.$$

For completeness, factor $T = P_m Q_n^s$, where

$$P_m(t) = \binom{t-1}{m}\binom{t+n+m}{m}, \qquad Q_n(t) = \frac{n!}{\prod_{h=0}^{n}(t+h)}.$$

Vandermonde expansion shows that d_n^ℓ clears the coefficient of x^ℓ in $P_m(-j+x)$. The same statement holds for $[xQ_n(-j+x)]^s$, since its factors are $(1+x/(h-j))^{-s}$, apart from an integral leading coefficient. Thus d_n^{s-k} clears each coefficient of $(t+j)^{-k}$. Summing the partial fractions proves the displayed denominator bounds.

Clear all rows by $D_n = d_n^s$. Let $C_{s,n}$ be the $(r-1)$-by-$(r+1)$ integer matrix of coefficients of the lower periods $\zeta(3), \dots, \zeta(s-2)$, and let $Y_{s,n}, Z_{s,n}$ be the cleared target and constant rows. Whenever $C_{s,n}$ has full row rank, its integral kernel

$$\Lambda_{s,n} = \ker_{\mathbb{Z}} C_{s,n}$$

is a saturated lattice of rank two. For an integral basis a, b, define

$$\Gamma_{s,n} = \gcd(Ya, Yb, Za, Zb). \tag{2}$$

Here and below gcds are nonnegative; we require $\Gamma_{s,n} > 0$ when dividing by it. This gcd is independent of the lattice basis.

Proposition 1.1 (Kernel rank and saturation)**.** *If* $\operatorname{rank}_{\mathbb{Q}} C = r-1$*, then* $\ker_{\mathbb{Z}} C \cong \mathbb{Z}^2$*, and this kernel is saturated in* $\mathbb{Z}^{r+1}$*.*

Proof. Smith normal form supplies unimodular matrices U, V with

$$UCV = (\operatorname{diag}(d_1, \dots, d_{r-1}) \mid 0 \mid 0), \qquad d_i > 0.$$

The equation $Cv = 0$ says precisely that the first $r-1$ coordinates of $V^{-1}v$ are zero. The final two coordinates are arbitrary integers, so the final two columns of V give an integral kernel basis. If kv belongs to this kernel, with $k \in \mathbb{Z} \setminus \{0\}$, then $0 = C(kv) = kCv$. Torsion-freeness of $\mathbb{Z}^{r-1}$ gives $Cv = 0$, which proves saturation. □

At weight five the established adjacent-sign theorem gives $x_{0,n}x_{1,n} < 0$ for the single lower-period row at every even $n \geq 2$. That row is nonzero and therefore has rational rank one; its integral kernel has rank two by the proposition. For higher weights, counting the $r-1$ rows is not a proof of their independence. The supplied rank argument explicitly retains that hypothesis.

Proposition 1.2 (Exact content by Smith reduction)**.** *Let* $\Delta_k(B)$ *be the nonnegative gcd of all* k*-minors of an integer matrix* B*, with* $\Delta_0(B) = 1$*. For a full-row-rank matrix* C *with* $r-1$ *rows and* $r+1$ *columns,*

$$\Gamma = \frac{\gcd(\Delta_r(C;Y), \Delta_r(C;Z))}{\Delta_{r-1}(C)}. \tag{3}$$

The notation $(C;Y)$ *means append the row* Y*.*

Proof. Choose unimodular U, V with $UCV = (\operatorname{diag}(d_1, \dots, d_{r-1}) \mid 0)$. The last two columns of V are a basis of the integral kernel. The gcd of the maximal minors of C is $\prod_i d_i$. The only possibly nonzero r-minors after appending Y are $(\prod_i d_i)Ya$ and $(\prod_i d_i)Yb$, up to sign. Unimodular row and column transformations preserve determinantal divisors. Repeat with Z and take the gcd. □

Consequently, with the convention $v_p(0) = +\infty$,

$$v_p(\Gamma) = \min\{v_p(\Delta_r(C;Y)), v_p(\Delta_r(C;Z))\} - v_p(\Delta_{r-1}(C)). \tag{4}$$

This is a divisibility problem, distinct from the real growth of the minors.

2 The exact integer-form bound

Suppose $w_{s,n} \in \Lambda_{s,n} \cap \mathbb{Z}_{\geq 0}^{r+1}$ is nonzero. Put $M_{s,n} = \|w_{s,n}\|_1$ and

$$h(w) = \gcd(Yw, Zw), \qquad L_{s,n} = \frac{(Yw)\zeta(s) + Zw}{h(w)}.$$

The numerator equals $D_n \sum_j w_j \mathcal{A}_{s,n,n-j} > 0$. In particular $h(w) > 0$. Since $\Gamma_{s,n} \mid h(w)$, we obtain

$$0 < L_{s,n} \leq d_n^s \frac{M_{s,n}}{\Gamma_{s,n}} S_{s,n}, \qquad S_{s,n} = \max_{0 \leq j \leq r} \mathcal{A}_{s,n,n-j}. \tag{5}$$

Both coefficients of $L_{s,n}$ are integers. Rational normalization of a kernel vector to mass one does not prove a bound for $M_{s,n}/\Gamma_{s,n}$, since clearing its denominator must still be included in (5).

3 A direct analytic extension to higher weights

The following argument makes the weight dependence explicit without a new saddle computation at each weight.

Lemma 3.1 (Extra powers of the kernel)**.** *For $n \geq 2r$, $0 \leq j \leq r$, and $k \geq n - j + 1$,*

$$Q_n(k) \leq 3^j 4^{-n}, \qquad T_{s,n,n-j}(k) = Q_n(k)^{s-5} T_{5,n,n-j}(k).$$

Proof. The binomial sum and maximality of the central binomial coefficient give $4^n \leq (2n+1)\binom{2n}{n}$, hence

$$Q_n(n+1) = \frac{1}{(2n+1)\binom{2n}{n}} \leq 4^{-n}.$$

The function $Q_n(k)$ decreases for positive k. Also $Q_n(k)/Q_n(k+1) = (k+n+1)/k$. For $k \geq n-j+1$, $n \geq 2j$, this ratio is at most three. At most j backward steps from $n+1$ suffice. The identity follows by cancellation of the definitions. □

For comparison of the fixed shifts at weight five, exact factorial cancellation gives

$$\begin{aligned} \frac{T_{5,n,n-j}(k)}{T_{5,n,n}(k+j)} &= \left(\frac{n!}{(n-j)!}\right)^2 \prod_{h=1}^{j} \left(\frac{k+n+h}{k+h-1}\right)^6 \left(\prod_{h=1}^{2j} (k+2n-j+h)\right)^{-1} \\ &\leq (3^6/4)^j. \end{aligned} \tag{6}$$

Indeed the first factorial ratio is at most n^j, each middle ratio is at most three, and each final factor is at least $2n$. The change $k \mapsto k+j$ carries the summation range onto $k+j \geq n+1$.

The elementary weight-five estimate, proved in Appendix A, is

$$\mathcal{A}_{5,n,n} \leq C_0 (n+1)^{12} 140^{-n}, \qquad C_0 = 12 \cdot 3^{12} \cdot 5^{10} e^{10}. \tag{7}$$

It is not a Lean theorem in the present companion. Appendix A also proves the sharp bound $\mathcal{A}_{5,n,n} \leq C_0 (n+1)^{12} e^{\lambda_5 n}$. The derivation locates the unique maximum of

$$6[x \log x - (x+1)\log(x+1)] + (x+2)\log(x+2) - (x-1)\log(x-1).$$

Writing $y = x(x+1)$, its stationary equation is $3y^3 - 4y^2 - 5y - 1 = 0$. The maximum occurs below $1057/1000$, where the resulting exponential base is below $1/140$. The complete summation and tail estimates in Appendix A supply the prefactor in (7) and certify the sharper interval $-4.970794 < \lambda_5 < -4.970793$.

Proposition 3.2 (Higher-weight raw decay). *For each fixed odd $s = 2r + 1 \geq 7$ and even $n \geq 2r$,*

$$S_{s,n} \leq C_s (n+1)^{12} \exp(-a_s n), \qquad a_s = \log 140 + (s-5)\log 4, \qquad C_s = C_0\, 3^{r(s+1)}. \tag{8}$$

Proof. Apply the lemma termwise, sum, and use (6). For the shift j, the constant obtained is $3^{j(s-5)}(3^6/4)^j \leq 3^{r(s+1)}$. The powers $140^{-n}4^{-(s-5)n}$ equal $\exp(-a_s n)$. □

Lemma 3.3 (Explicit margin, checked in Lean). *For every integer $s \geq 7$,*

$$a_s > s + \tfrac{1}{3}. \tag{9}$$

Proof. The elementary inequality $2x/(x+2) < \log(1+x)$, applied at $x = 1$, gives $\log 2 > 2/3$. Also $\log 140 > 7\log 2 > 14/3$ and $\log 4 = 2\log 2 > 4/3$. Thus

$$a_s > \frac{14}{3} + \frac{4}{3}(s-5) = \frac{4s-6}{3} \geq s + \frac{1}{3}.$$

□

4 The precise conditional irrationality theorem

All limits in this section may be restricted to an unbounded sequence of eligible even indices. Write $\log^+ x = \max(0, \log x)$ for $x > 0$.

Theorem 4.1 (Relative arithmetic cost). *Fix an odd $s \geq 7$. Suppose that the full-rank cancellation setup, the positive integral vectors, and the integer forms in (5) exist along an unbounded sequence of even indices. Suppose (8) holds and*

$$\log^+(M_{s,n}/\Gamma_{s,n}) = o(n). \tag{10}$$

Then $\zeta(s)$ is irrational.

Proof. The prime number theorem yields $\log d_n = n + o(n)$. For fixed s, the denominator term has upper exponential rate s. The polynomial factor in (8) has rate zero. By (10) the relative arithmetic cost also has upper rate at most zero. Taking logarithms in (5) therefore gives

$$\limsup \frac{\log L_{s,n}}{n} \leq s - a_s < -\tfrac{1}{3}.$$

Hence the nonzero integer linear forms tend to zero. If $\zeta(s) = p/q$, $p, q \in \mathbb{Z}$, $q \neq 0$, then $qL_{s,n}$ is a nonzero integer and has absolute value at least one, contradicting convergence to zero. □

For clarity, the Lean theorem uses eventual exponential envelopes instead of a quotient of logarithms. With $D_n = d_n^s$, $W_n = M_{s,n}/\Gamma_{s,n}$, its assumptions are, for every $\varepsilon > 0$, eventually

$$D_n \leq e^{(s+\varepsilon)n}, \qquad W_n \leq e^{\varepsilon n}, \qquad S_n \leq e^{(-a_s+\varepsilon)n}.$$

These are supplied as hypotheses to the theorem, together with (5) and arbitrarily late nonvanishing. Choosing $\varepsilon = 1/12$ proves the explicit eventual bound

$$|L_n| \leq e^{-n/12} \longrightarrow 0.$$

The formal theorem permits any index map tending to infinity, including the even subsequence; no finite-index extrapolation is used.

5 An integral projector for the divisibility obligation

At weight five the cancellation row is a nonzero integer row x. Put $g = \gcd(x_1, x_2, x_3)$, $\ell = x/g$, so that ℓ is primitive. Choose $u \in \mathbb{Z}^3$ with $\ell u = 1$. Define

$$\Pi = I - u\ell.$$

Then $\Pi^2 = \Pi$, $\ell\Pi = 0$, and $\Pi v = v$ whenever $\ell v = 0$. Its integral image is exactly $\Lambda = \ker_{\mathbb{Z}} \ell$.

Proposition 5.1 (Universal congruence test, checked in Lean)**.** *For any integer matrix F with three columns and any $c \in \mathbb{Z}$,*

$$c \mid (F\Pi)_{ij} \text{ for every } i, j \quad \Longleftrightarrow \quad c \mid (Fv)_i \text{ for every } v \in \Lambda \text{ and every } i. \tag{11}$$

The gcd of the entries of $F\Pi$ is independent of the choice of u, up to multiplication by a unit.

Proof. If the left side holds, write $v = \Pi v$ and expand $F\Pi v$ as an integral linear combination of its columns. Conversely apply the right side to each column Πe_j, which lies in Λ. The equivalence characterizes all divisors of the entry gcd and is independent of the section u. □

Taking F to have rows Y, Z identifies its entry gcd with Γ in (2). This formulation accounts for division by g before imposing higher-order congruences. For $c = p^e$, it is an exact reformulation of the local obligation, not a lower bound for its exponent.

6 Shape control and the weight-five boundary

The augmented-rank condition used here has the following exact proof. Choose an integral kernel basis a, b, under the lower-row rank condition, and put

$$B = \begin{pmatrix} Y(a) & Y(b) \\ Z(a) & Z(b) \end{pmatrix}.$$

Apply the Smith transformations from Section 1 to the augmented matrix. Its first rows have diagonal entries $d_1, \ldots, d_{r-1}$, and its lower right corner is B. Unimodular transformations change determinant only by a sign. Expansion along the first rows proves

$$|\det(C; Y; Z)| = \Delta_{r-1}(C) |\det B|. \tag{12}$$

Thus the augmented square matrix has full rational rank if and only if $\det B \neq 0$. This proves the equivalence; it does not assert determinant nonvanishing for every member of the higher-weight Padé family.

Let a, b be a Gauss-reduced basis of a saturated rank-two cancellation lattice, $\lambda_1 = \|a\| \leq \lambda_2 = \|b\|$, and $H = \|a \wedge b\|$. Set

$$U_Y = Y(a)b - Y(b)a, \qquad U_Z = Z(a)b - Z(b)a, \qquad \mathcal{M} = \max(\|U_Y\|, \|U_Z\|).$$

Assume the augmented square matrix $(C; Y; Z)$ has nonzero determinant.

Proposition 6.1 (Cofactor comparison)**.** *Under these assumptions,*

$$\lambda_2 \leq \frac{\mathcal{M}}{\Gamma}, \qquad \kappa = \frac{\lambda_2}{\lambda_1} \leq \frac{\mathcal{M}^2}{\Gamma^2 H}, \qquad \frac{\Gamma}{\lambda_2} \geq \frac{\Gamma^2}{\mathcal{M}}.$$

Proof. The vectors $U_Y/\Gamma, U_Z/\Gamma$ are independent lattice vectors. The second successive minimum is therefore at most $\mathcal{M}/\Gamma$; for a Gauss-reduced rank-two basis it equals λ_2. Since $H = \lambda_1\lambda_2 \sin\theta$ and $\sin\theta \le 1$, $\kappa \le \lambda_2^2/H$. Combine the two inequalities. □

In weight five, positive rounding gives the separate sufficient bound

$$M_{5,n} \le 3(1 + 2R_n)\lambda_{2,n}.$$

The row-balance result $\log R_n = o(n)$ removes this factor from the exponential budget, but not the factor $\lambda_{2,n}/\Gamma_{5,n}$. Using only the explicit base 140, a sufficient joint estimate is

$$\Gamma_{5,n} \ge e^{0.06n}\lambda_{2,n}.$$

The sharper saddle value $-4.970794 < \lambda_5 < -4.970793$, certified in Appendix A, reduces 0.06 to 0.03. The sharper constant enclosure and this sufficient joint estimate are not formalized in the new Lean modules. Neither joint arithmetic estimate has been proved for the constructed family. In particular this section supplies no unconditional theorem for $\zeta(5)$.

7 Signed cancellation and a cofactor restriction

For real weights of either sign, the triangle inequality gives

$$\left|\sum_j w_j \mathcal{A}_j\right| \le \left(\sum_j |w_j|\right) \max_j |\mathcal{A}_j|. \tag{13}$$

Thus nonnegativity is not essential to the upper estimate. Its role in Section 2 is to guarantee nonvanishing. A different guarantee is available for two coefficient pairs: if $ad - bc \neq 0$, the forms $a\alpha + c$ and $b\alpha + d$ cannot both vanish, since

$$a(b\alpha + d) - b(a\alpha + c) = ad - bc.$$

One may therefore work with a pair of independent signed forms, provided both satisfy the required arithmetic bounds. These two elementary assertions are proved in the signed Lean companion.

Proposition 7.1 (Individual primitive normalization)**.** *Let a, b be any integral kernel basis, and suppose $\delta = Y(a)Z(b) - Y(b)Z(a) \neq 0$. Put*

$$h_a = \gcd(Y(a), Z(a)), \quad h_b = \gcd(Y(b), Z(b)), \quad A = \|a\|_2^2,\ B = \langle a, b\rangle,\ C = \|b\|_2^2.$$

Both gcds are positive. The two primitive integer forms

$$L_a = \frac{Y(a)\zeta(s) + Z(a)}{h_a}, \qquad L_b = \frac{Y(b)\zeta(s) + Z(b)}{h_b}$$

cannot both vanish. Define

$$G^\sharp = \begin{pmatrix} A/h_a^2 & B/(h_a h_b) \\ B/(h_a h_b) & C/h_b^2 \end{pmatrix}, \qquad \tau^\sharp = A/h_a^2 + C/h_b^2.$$

Writing $N = r + 1$ and $\mu^\sharp$ for the larger eigenvalue,

$$\max(|L_a|, |L_b|) \le \sqrt{N}\, d_n^s S_{s,n} \sqrt{\mu^\sharp} \le \sqrt{N}\, d_n^s S_{s,n} \sqrt{\tau^\sharp}, \qquad \tau^\sharp \le \frac{A + C}{\Gamma^2}. \tag{14}$$

No reduction condition on the individually scaled basis is required.

Proof. A zero gcd would make the corresponding coefficient column zero, contradicting $\delta \neq 0$. Dividing each column by its gcd gives integer coefficients and determinant $\delta/(h_a h_b) \neq 0$. The preceding determinant identity proves nonvanishing of at least one form. The signed bound and Cauchy–Schwarz give $|L_a| \leq \sqrt{N}\, d_n^s S_{s,n} \sqrt{A}/h_a$, and similarly for b. The matrix $G^\sharp$ is the Gram matrix of $a/h_a, b/h_b$. Thus its diagonal entries are at most $\mu^\sharp$, and $\mu^\sharp \leq \operatorname{tr} G^\sharp$. Equivalently these inequalities follow directly from the quadratic root formula and $B^2 \leq AC$. Finally $\Gamma \mid h_a, h_b$ implies $h_a, h_b \geq \Gamma$, proving the last inequality. □

For every fixed $s \geq 7$, a sufficient condition for the nonzero selected primitive forms in (14) to tend to zero is

$$\limsup_{n\to\infty} \frac{1}{n} \log \tau^\sharp_{s,n} < 2(a_s - s). \tag{15}$$

Indeed the upper exponential rate in (14) is $s - a_s + \frac{1}{2} \limsup n^{-1} \log \tau^\sharp$. A rate at most $1/2$ would suffice at every such weight, since $a_s > s + 1/3$. This substitution permits additional divisibility in each selected vector; it does not require a separate lower bound for Γ. Neither (15) nor uniform augmented determinant nonvanishing has been proved for the actual higher-weight family. No Loewner comparison between $G^\sharp$ and G/Γ^2 is asserted: unequal diagonal scalings need not commute with G.

However, a signed combination that eliminates the target itself is a nonzero rational period and has an integral lower bound after clearing. In the notation of Section 6, put

$$\delta = Y(a)Z(b) - Y(b)Z(a) \neq 0.$$

Then $CU_Y = 0$, $Y(U_Y) = 0$, and $Z(U_Y) = \delta$. Since Γ divides all four displayed evaluations, $\Gamma^2 \mid \delta$, so $|\delta| \geq \Gamma^2$. Applying the cleared period identity and (13) gives

$$\boxed{\Gamma^2_{s,n} \leq |\delta| = d_n^s \left| \sum_j (U_Y)_j \mathcal{A}_{s,n,n-j} \right| \leq d_n^s \sqrt{r+1}\, \mathcal{M}_{s,n} S_{s,n}.} \tag{16}$$

Here $\|U_Y\|_1 \leq \sqrt{r+1}\|U_Y\|_2 \leq \sqrt{r+1}\mathcal{M}$. The determinant divisibility, signed bound, and integer-divisor lower bound are proved in `SignedCancellation.lean`; their application to these coefficient rows is the displayed paper derivation.

It follows from (8) and $\log d_n = n + o(n)$ that

$$\limsup \frac{1}{n} \log \frac{\Gamma^2_{s,n}}{\mathcal{M}_{s,n}} \leq s - a_s.$$

In particular, for $s \geq 7$ this upper rate is less than $-1/3$. At weight five the coarse base 140 gives the upper rate $5 - \log 140 < 0.058358$; the sharper saddle rate gives $5 + \lambda_5 < 0.029207$. Consequently the sufficient cofactor strengthenings $\Gamma^2_{5,n} \geq e^{0.06n}\mathcal{M}_{5,n}$, or $\Gamma^2_{5,n} \geq e^{0.03n}\mathcal{M}_{5,n}$ with the sharp saddle bound, cannot hold along an unbounded sequence with nonzero augmented determinant. This restriction does not invalidate the comparison in Section 6. It shows that using its cofactor upper bound in place of the actual second minimum is too strong for that proposed arithmetic target. The weaker bound for a selected primitive form remains a separate problem.

8 Row balance and positive integral cost

The relative-cost estimate needs arithmetic information about the specified family. In particular it does not follow from row balance and the existence of a positive integral kernel vector alone. This can be proved exactly, without extrapolating from numerical examples.

For an integer $t \geq 2$, take

$$C = (t, -(t-1), t), \qquad Y = (1, 0, 0), \qquad Z = (0, 0, 1).$$

The cancellation row is primitive, has full row rank, and its positive to negative row-sum ratio is $2t/(t-1) \leq 4$. The integral kernel contains $(1, 0, -1)$, on which Y evaluates to one, so its target-and-constant content is $\Gamma = 1$. If $w = (a, b, c) \geq 0$ is a nonzero integral kernel vector, then

$$t(a + c) = (t - 1)b, \qquad k = b - a - c \in \mathbb{Z}_{\geq 1}.$$

It follows that $a + c = (t-1)k$, $b = tk$, and

$$\|w\|_1 = (2t - 1)k \geq 2t - 1.$$

Equality is attained by $(t - 1, t, 0)$. Choosing $t = 2^n$ gives a bounded row-balance ratio but forces every nonzero nonnegative cancellation vector to have exponentially growing M/Γ. The lower bound, its attaining vector, the row-ratio bound, and the unit-content implication are checked in Lean. This family is not the odd-zeta family: it proves a non-implication between the general structural properties and the desired arithmetic estimate.

9 Formalization and reproducibility

The archive contains the pinned Lean project and every source module. The principal additions are:

- `OddZeta/HighWeightRelativeCost.lean`: the logarithm margin, convergence from eventual rate bounds, and the conditional irrationality theorem for `oddZetaValue k`, $k \geq 3$, meaning the real part of $\zeta(2k + 1)$.
- `OddZeta/KernelProjectorContent.lean`: the integral projection, idempotence, universal divisibility equivalence, and independence of the content from the section.
- `OddZeta/PositiveCancellationCost.lean`: the exact minimal positive mass calculation above, with the row-ratio and unit-content assertions.
- `OddZeta/GramSpectralBounds.lean`: the two-dimensional characteristic roots, reduced-basis eigenvalue bounds, condition-number bounds, and the uniform integral spectral gap.
- `OddZeta/PrimitivePairNormalization.lean`: the eigenvalue bounds for a general positive-semidefinite two-by-two Gram matrix, its individual scaling and trace bound, exact division of integer forms, and determinant-based primitive-pair nonvanishing.

The publication folder also contains `SignedCancellation.lean` and exact cancellation examples at weight seven and indices six and eight. Two independent rational coefficient algorithms agree at these indices, and Lean verifies the four resulting integer cancellation identities. They confirm actual cancellation, with no extrapolation to all indices. All parameters of the irrationality criterion remain explicit. In particular the cost bound and nonvanishing are hypotheses, and no axiom asserts their existence for the Padé family. The complete source declaration, rather than its theorem name alone, specifies the mathematical assertion checked by Lean.

A successful Lean compilation verifies these formal statements relative to the pinned library. It does not verify paper-only derivations such as (3), (7), or the Gauss-reduction comparison. The

accompanying verification report distinguishes these scopes. Rebuild with the pinned toolchain, using `lake exe cache get` and `lake build`. The provided lock file records the library revisions.

An unconditional application to all odd weights would require a proof of (10) for explicitly constructed nonnegative integral cancellation vectors at each fixed weight, or alternatively the signed pair rank and rate hypotheses in (15). The exact projector criterion and the Smith formula specify the arithmetic content entering the first route. Individual normalization weakens that route's shared-content requirement but does not establish the remaining construction-specific estimates.

A Global decay with the sharp saddle constant

Here is the full analytic proof of (7) and its sharper version. For $x > 1$, define

$$\Phi(x) = 6[x \log x - (x+1)\log(x+1)] + (x+2)\log(x+2) - (x-1)\log(x-1).$$

Its derivative has the sign of $x^6(x+2) - (x-1)(x+1)^6$. With $y = x(x+1)$, the negative of this expression is $p(y) = 3y^3 - 4y^2 - 5y - 1$. For $y \geq 2$, $p'(y) = 9y^2 - 8y - 5 > 0$, while $p(2) = -3$. Therefore Φ increases to a unique $x_* > 1$ and decreases thereafter. Subtracting $x_*\Phi'(x_*) = 0$ from its value gives

$$\lambda_5 = \Phi(x_*) = \log q(x_*), \qquad q(x) = \frac{(x+2)^2(x-1)}{(x+1)^6}.$$

Set $a = 1057/1000$. Exact integer arithmetic proves

$$1000^6 p(a(a+1)) = 54685013669880747 > 0,$$
$$2057^6 - 140 \cdot 3057^2 \cdot 57 \cdot 1000^3 = 1178944474661131249 > 0.$$

Hence $1 < x_* < a$. On this interval

$$(\log q)' = \frac{2}{x+2} + \frac{1}{x-1} - \frac{6}{x+1} > \frac{1000}{57} - 3 > 0.$$

The second integer inequality is exactly $q(a) < 1/140$, so $\lambda_5 < -\log 140$ is a global bound.

For the sharp certificate put

$$x_- = \frac{1056126503577}{10^{12}}, \qquad x_+ = \frac{1056126503578}{10^{12}}.$$

Exact rational evaluation gives $p(x_-(x_-+1)) < 0$ and $p(x_+(x_++1)) > 0$. Thus $x_- < x_* < x_+$, and monotonicity of q encloses λ_5 between the logarithms of these rational values. For $1 \leq u \leq 2$, let $z = (u-1)/(u+1)$. The convergent series gives

$$0 \leq \log u - 2\sum_{k=0}^{L-1} \frac{z^{2k+1}}{2k+1} \leq \frac{2z^{2L+1}}{(2L+1)(1-z^2)}.$$

For any positive rational argument, first factor out an integral power of two. Applying this formula with $L = 32$ and outward rational rounding proves

$$-4.970793860358 < \lambda_5 < -4.970793860341.$$

The signs and rational enclosures are reproduced by `check_high_weight_rates.py`. They certify a single global maximum, without extrapolation from indices.

To bound the infinite sum, write $T = T_{5,n,n}$. For $t \geq 2n$, logarithmic differentiation gives

$$\frac{d}{dt}\log(t^2T(t)) \leq -\frac{E}{t} + \frac{B}{t^2}, \qquad E = 3n+3, \quad B = 7n^2+5n.$$

Indeed apply $1/(t-v) \leq 1/t + 2v/t^2$, $1/(t+n+v) \leq 1/t$, and $1/(t+h) \geq 1/t - h/t^2$ to its finite logarithmic derivative. Set $K = \max(2n, n+1, \lceil B/E \rceil) \leq 3n$. Then $t^2T(t)$ is decreasing for $t \geq K$, so $T(k) \leq T(K)(K/k)^2$ there. The maximum summand P_n is attained at some $n+1 \leq k \leq K$. An integral comparison of the tail of $\sum k^{-2}$ gives

$$\mathcal{A}_{5,n,n} \leq (2K+1)P_n \leq (6n+1)P_n.$$

For $f(v) = v\log v - v$, $f(0) = 0$, integral comparison also gives

$$f(h) \leq \log(h!) \leq f(h) + 1 + \log(h+1) \qquad (h \in \mathbb{Z}_{\geq 0}).$$

At a maximizing index $n+1 \leq k \leq 3n$, the exact factorial identity is

$$T(k) = \frac{(n!)^3((k-1)!)^6(k+2n)!}{(k-n-1)!((k+n)!)^6}.$$

Apply the upper factorial bound to the ten numerator factors and the lower bound to the denominator. The total error is at most $10[1+\log(5n+1)]$. In the main terms, $f(k-1) - f(k) \leq 0$ and $f(k-n) - f(k-n-1) \leq \log(k-n) \leq \log(2n)$, including $k-n = 1$. Replacing these arguments accordingly leaves exactly $n\Phi(k/n) \leq n\lambda_5$, apart from error at most $\log(2n)$. Consequently, for every even $n \geq 4$,

$$\begin{aligned}\mathcal{A}_{5,n,n} &\leq 2e^{10}n(6n+1)(5n+1)^{10}e^{\lambda_5 n} \\ &\leq 12\cdot 5^{10}e^{10}(n+1)^{12}e^{\lambda_5 n} \leq C_0(n+1)^{12}140^{-n}.\end{aligned}$$

This proves the complete raw decay estimate. Section 3 extends it to every higher fixed weight. The primitive integer normalization cost remains in (5); no analytic step here discards that factor.

References

[1] *A Smith-content reduction for odd-zeta Padé forms*, user-supplied technical note, 28 October 2026, five pages.

[2] *Odd-zeta collected notes*, user-supplied compilation of six working notes, thirty pages.

[3] Companion Lean project and derivation notes, GitHub repository. The archive contains a source snapshot and a verification report.