MAT168 HW4

Andrew Jowe

November 28, 2022

(1)

Iteration 1

Draw out original problem

l		0			0			
	u			6		8		
		ζ		$-x_1$		_		
$-\infty$	5	w_1	=	$\begin{array}{c c} -x_1 \\ x_1 \end{array}$	+	x_2	=	0
$-\infty$	9	$ w_2 $	=	x_1	_	$2x_2$	=	0

Determine entering variable

 x_2 has positive coefficient and is at lower bound, x_2 enters.

Determine leaving variable

As $x_2 \rightarrow 8$:

$$w_1 \to -\infty \le x_2 \le 5 \to x_2 = 5$$

$$w_2 \to -\infty \le -2x_2 \le 9 \to x_2 = \infty$$

 w_1 leaves.

Iteration 2

Rewriting equation

l				0		$-\infty$		
4	$u \mid$			6		5		
		ζ	=	$0x_1$	+	w_1	=	5
0	8	x_2	=	x_1	+	w_1	=	5
$-\infty$	9	w_2	=	$-x_1$	_	$2w_2$	=	-10

Optimal

This is optimal since w_1 has positive coefficient and upper bound is chosen, and x_1 has coefficient of 0.

Solution

$$w_1 = 5$$
$$x_2 = 5$$

$$x_1 = 0$$

$$w_2 = -10$$

$$\zeta = 5$$

(2)

Primal

Given

$$c = \begin{bmatrix} -1\\1 \end{bmatrix}$$

$$A = \begin{bmatrix} -1 & 1 \\ 1 & -2 \end{bmatrix}$$

$$b = \begin{bmatrix} 5 \\ 9 \end{bmatrix}$$

$$a = \begin{bmatrix} -\infty \\ -\infty \end{bmatrix}$$

$$u = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$$

$$l = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Slack Variables

$$f = \begin{bmatrix} 0 \\ 19 \end{bmatrix}$$

$$p = \begin{bmatrix} \infty \\ \infty \end{bmatrix}$$

$$t = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$$

$$g = \begin{bmatrix} 0 \\ 5 \end{bmatrix}$$

Dual

Equations

minimize
$$\begin{bmatrix} 5 \\ 9 \end{bmatrix}^T v - \begin{bmatrix} -\infty \\ -\infty \end{bmatrix}^T q + \begin{bmatrix} 6 \\ 8 \end{bmatrix}^T s - \begin{bmatrix} 0 \\ 0 \end{bmatrix}^T h$$

subject to
$$\begin{bmatrix} -1 & 1 \\ 1 & -2 \end{bmatrix}^T (v - q) - (h - s) = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Complimentary Slackness

$$f_2 = 19 \rightarrow v_2 = 0$$

$$p_i \neq 0 \rightarrow q_i = 0 \quad \forall i \in [1, m]$$

$$t_j \neq 0 \rightarrow s_j = 0 \quad \forall j \in [1, n]$$

$$g_2 = 5 \rightarrow h_2 = 0$$

Rewrite Equations Iteration 1

minimize
$$5v_1^+$$
 0 subject to $\begin{bmatrix} -1 & 1 \\ 1 & -2 \end{bmatrix}v - h = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Rewrite Equations Iteration 2

minimize
$$5v_1^+$$

 0
subject to $-v_1 - h_1 = -1$
 $v_1 = 1$

Solve

$$v_1 = 1$$

 $v_2 = 0$
 $q_1, q_2 = 0$
 $s_1, s_2 = 0$
 $h_1, h_2 = 0$
 $\zeta = 5v_1 = 5(1) = 5$

Academic Integrity

On my personal integrity as a student and member of the UCD community, I have not given, nor received any unauthorized assistance on this assignment.

Andrew