

## 12. Definite Integration - II

**Ex. (1)** Evaluate  $\int_0^2 |4x-5| dx$

**Solution :**  $|4x-5| = \begin{cases} 4x-5, & \text{if } x \geq \frac{5}{4} \\ -(4x-5), & \text{if } x < \frac{5}{4} \end{cases}$

$$\begin{aligned} \int_0^2 |4x-5| dx &= \int_0^{\frac{5}{4}} |4x-5| dx + \int_{\frac{5}{4}}^2 |4x-5| dx \\ &= \int_0^{\frac{5}{4}} -(4x-5) dx + \int_{\frac{5}{4}}^2 (4x-5) dx \\ &= \int_0^{\frac{5}{4}} (-4x+5) dx + \int_{\frac{5}{4}}^2 (4x-5) dx \\ &= \left[ \frac{-4x^2}{2} + 5x \right]_0^{\frac{5}{4}} + \left[ \frac{4x^2}{2} - 5x \right]_{\frac{5}{4}}^2 \\ &= \left\{ \left[ -2\left(\frac{5}{4}\right)^2 + 5\left(\frac{5}{4}\right) \right] - [0] \right\} + \left\{ \left[ 2(2)^2 - 5(2) \right] - \left[ 2\left(\frac{5}{4}\right)^2 - 5\left(\frac{5}{4}\right) \right] \right\} \\ &= \frac{-50}{16} + \frac{25}{4} - 2 - \frac{50}{16} + \frac{25}{4} \\ &= \frac{-50+100-32-50+100}{16} = \frac{68}{16} = \frac{17}{4} \end{aligned}$$

$$\therefore \int_0^2 |4x-5| dx = \frac{17}{4}$$

**Ex. (2)** Evaluate  $\int_0^\pi \frac{x \tan x}{\sec x + \tan x} dx$

**Solution :**  $I = \int_0^\pi \frac{x \tan x}{\sec x + \tan x} dx = \int_0^\pi \frac{\frac{x \sin x}{\cos x}}{\frac{1}{\cos x} + \frac{\sin x}{\cos x}} dx$

$$= \int_0^\pi \frac{x \sin x}{1 + \sin x} dx = \int_0^\pi \frac{(\pi - x) \sin(\pi - x)}{1 + \sin(\pi - x)} dx$$

$$= \int_0^\pi \frac{(\pi - x) \sin x}{1 + \sin x} dx$$

$$I = \int_0^\pi \frac{\pi \sin x}{1 + \sin x} dx - \int_0^\pi \frac{x \sin x}{1 + \sin x} dx$$

$$I = \int_0^{\pi} \frac{\pi \sin x}{1 + \sin x} dx - I$$

$$2I = \int_0^{\pi} \frac{\pi \sin x}{1 + \sin x} dx$$

$$2I = \pi \int_0^{\pi} \frac{\sin x}{1 + \sin x} dx$$

$$2I = \pi \int_0^{\pi} \left( 1 - \frac{1}{1 + \sin x} \right) dx$$

$$2I = \pi \int_0^{\pi} \left( 1 - \frac{1 - \sin x}{1 - \sin^2 x} \right) dx$$

$$2I = \pi \int_0^{\pi} \left( 1 - \frac{1 - \sin x}{\cos^2 x} \right) dx$$

$$2I = \pi \int_0^{\pi} \left( 1 - \frac{1}{\cos^2 x} + \frac{\sin x}{\cos^2 x} \right) dx$$

$$2I = \pi \int_0^{\pi} (1 - \sec^2 x + \sec x \tan x) dx$$

$$2I = \pi [x - \tan x + \sec x]_0^{\pi}$$

$$2I = \pi [(\pi - \tan \pi + \sec \pi) - (0 - \tan 0 + \sec 0)]$$

$$2I = \pi [\pi - 0 - 1 - 0 - 1] = \pi (\pi - 2)$$

$$I = \frac{\pi}{2} (\pi - 2) = \left( \frac{\pi^2}{2} - \pi \right)$$

**Ex. (3)**  $\int_{-3}^3 \frac{x + x^2}{25 - x^2} dx$

**Solution :**  $I = \int_{-3}^3 \frac{x + x^2}{25 - x^2} dx$

$$I = \int_{-3}^3 \frac{x}{25 - x^2} dx + \int_{-3}^3 \frac{x^2}{25 - x^2} dx \dots (1)$$



$$\text{Let } f(x) = \frac{x}{25-x^2}$$

$$f(-x) = \frac{-x}{25-(-x)^2} = -\frac{x}{25-x^2} = -f(x)$$

$\therefore f(x)$  is an odd function.

$$\therefore \int_{-3}^3 \frac{x}{25-x^2} dx = 0 \quad \dots (2)$$

$$\text{Let } g(x) = \frac{x^2}{25-x^2}$$

$$g(-x) = \frac{(-x)^2}{25-(-x)^2} = \frac{x^2}{25-x^2} = g(x)$$

$\therefore g(x)$  is an even function.

$$\therefore \int_{-3}^3 \frac{x^2}{25-x^2} dx = 2 \int_0^3 \frac{x^2}{25-x^2} dx \quad \dots (3)$$

From (1), (2) and (3) we get,

$$\begin{aligned} I &= 2 \int_0^3 \frac{x^2}{25-x^2} dx \\ &= 2 \int_0^3 \frac{25-25+x^2}{25-x^2} dx \\ &= 2 \int_0^3 \frac{25-(25-x^2)}{25-x^2} dx \\ &= 2 \int_0^3 \frac{25}{25-x^2} - 1 dx \\ &= 2 \left[ \frac{25}{2(5)} \log \left( \frac{5+x}{5-x} \right) - x \right]_0^3 \\ &= 2 \left[ \left( \frac{5}{2} \log \left( \frac{5+3}{5-3} \right) - 3 \right) - \left( \frac{5}{2} \log \left( \frac{5+0}{5-0} \right) - 0 \right) \right] \\ &= 2 \left[ \frac{5}{2} \log \left( \frac{8}{2} \right) - 3 - \frac{5}{2} \log(1) \right] \\ &= 2 \left[ \frac{5}{2} \log(4) - 3 \right] \\ &= 5 \log 4 - 6 \end{aligned}$$

$$\therefore \int_{-3}^3 \frac{x+x^2}{25-x^2} dx = 5 \log 4 - 6$$



**Ex. (4)** Evaluate  $\int_0^2 ([x] + |x-1|) dx$

**Solution :**

$$I = \int_0^2 ([x] + |x-1|) dx$$

using  $\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx \quad (a < b < c)$

$$\therefore I = \int_0^1 ([x] + |x-1|) dx + \int_1^2 ([x] + |x-1|) dx$$

$$[x] \quad 0-1$$

$$|x| = x, \quad x > 0$$

$$[x] = 0$$

$$= -x, \quad x < 0$$

$$1-2$$

$$[x] = 1$$

$$|x-1| = x-1, \quad x-1 > 0$$

$$= -(x-1), \quad x-1 < 0$$

$$|x-1| = x-1, \quad x > 1$$

$$= 1-x, \quad x < 1$$

$$I = \int_0^1 (0 + 1 - x) dx + \int_1^2 (1 + x - 1) dx$$

$$= \int_0^1 (1 - x) dx + \int_1^2 x dx$$

$$= \left[ x - \frac{x^2}{2} \right]_0^1 + \left[ \frac{x^2}{2} \right]_1^2$$

$$= 1 - \frac{1}{2} - 2 \left( 0 - \frac{0}{2} \right) + \frac{4}{2} - \frac{1}{2} \Rightarrow \frac{1}{2} + 2 - \frac{1}{2}$$

$$I = 2$$

**Ex. (5)** Evaluate  $\int_e^{e^2} \frac{1}{\log x} - \frac{1}{(\log x)^2} dx$

**Solution :**

$$I = \int_e^{e^2} \left[ \frac{1}{\log x} - \frac{1}{(\log x)^2} \right] dx$$

$$t = \log x$$

$$\text{when } x = e, \quad t = \log e = 1$$

$$\text{when } x = e^2, \quad t = \log e^2 = 2$$

$$\text{put } \log x = t \Rightarrow x = e^t$$

$$\therefore dx = e^t dt$$



$$\therefore I = \int_1^2 \left[ \frac{1}{t} - \frac{1}{t^2} \right] dt$$

$$= \int_1^2 e^t \left[ \frac{1}{t} - \left( -\frac{1}{t^2} \right) \right] dt$$

$$= \left[ e^t \left( \frac{1}{t} \right) \right]_1^2 \quad \because \int e^x [f(x) + f'(x)] dx = e^x f(x) + c$$

$$= e^2 \left( \frac{1}{2} \right) - e^1 \left( \frac{1}{1} \right)$$

$$= \frac{e^2}{2} - e$$

$$I = \frac{e^2 - 2e}{2}$$

$$= \frac{e(e-2)}{2}$$

**Ex. (7)** Evaluate  $\int_0^{\pi} \frac{x \tan x}{\sec x \operatorname{cosec} x} dx$

**Solution :**

using  $\int_a^b f(x) dx = \int_a^b f(a-x) dx$

changing  $x \rightarrow \pi - x$

$$\begin{aligned} I &= \int_0^{\pi} \frac{(\pi-x) \tan(\pi-x)}{\sec(\pi-x) \operatorname{cosec}(\pi-x)} dx \\ &= \int_0^{\pi} \frac{(\pi-x)(-\tan x)}{(-\sec x) \operatorname{cosec} x} dx \\ &= \int_0^{\pi} \frac{(\pi-x) \tan x}{\sec x \cdot \operatorname{cosec} x} dx \\ &= \int_0^{\pi} \frac{\pi \tan x - x \tan x}{\sec x \cdot \operatorname{cosec} x} dx \\ &= \int_0^{\pi} \frac{\pi \tan x}{\sec x \cdot \operatorname{cosec} x} dx - \int_0^{\pi} \frac{x \tan x}{\sec x \cdot \operatorname{cosec} x} dx \end{aligned}$$

$$I = \int_0^{\pi} \frac{\pi \tan x}{\sec x \cdot \operatorname{cosec} x} dx - I \quad \text{by } I$$

$$2I = \pi \int_0^{\pi} \frac{\sin x}{\cos x} x \cos x \sin x dx$$

$$2I = \pi \int_0^{\pi} \sin^2 x dx$$

$$2I = \frac{\pi}{2} \int_0^{\pi} 2 \sin^2 x dx$$

$$\because \cos 2\theta = 1 - 2 \sin^2 \theta$$

$$I = \frac{\pi}{4} \int_0^{\pi} (1 - \cos 2x) dx$$

$$I = \frac{\pi}{4} \left[ x - \frac{\sin 2x}{2} \right]_0^{\pi}$$



$$I = \frac{\pi}{4} \left[ \pi - \frac{\sin 2\pi}{2} - \left( 0 - \frac{\sin 0}{2} \right) \right]$$

$$= \frac{\pi}{4} \times \pi$$

$$I = \frac{\pi^2}{4}$$

**Ex. (8)** Evaluate  $\int_0^3 x [x] dx$ , where  $[x]$  denote greatest integer function not greater than  $x$ .

**Solution :**

$$0-1 [x] = 0, \quad 1-2 [x] = 1, \quad 2-3 [x] = 2$$

$$I = \int_0^3 x [x] dx$$

if  $a < b < c$  then  $\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$

$$I = \int_0^1 x [x] dx + \int_1^2 x [x] dx + \int_2^3 x [x] dx \quad (0 < 1 < 2 < 3)$$

$$= \int_0^1 x(0) dx + \int_1^2 x(1) dx + \int_2^3 x(2) dx$$

$$= 0 + \int_1^2 x dx + \int_2^3 2x dx$$

$$= \left[ \frac{x^2}{2} \right]_1^2 + \left[ \frac{2x^2}{2} \right]_2^3$$

$$= \left[ \frac{x^2}{2} \right]_1^2 + [x^2]_2^3$$

$$= \frac{4}{2} - \frac{1}{2} + 9 - 4$$

$$= 7 - \frac{1}{2}$$

$$\therefore I = \frac{13}{2}$$

Sign of Teacher :

- Q. 26.** A solenoid of length  $\pi$  m and 5 cm in diameter has a winding of 1000 turns and carries a current of 5A. Calculate the magnetic field at its centre along the axis.

## SECTION – D

**Attempt any THREE questions of the following :**


**[12]**

- Q. 27.** What is Ferromagnetism? Explain it on the basis of domain theory.
- Q. 28.** Obtain an expression for average power dissipated in a series LCR circuit.
- Q. 29.** Distinguish between interference and diffraction of light.  
A double slit arrangement produces interference fringes for sodium light of wavelength 589 nm, that are 0.20 degree apart. What is the angular fringe separation if the entire arrangement is immersed in water?  
( R.I. of water = 1.33 )
- Q. 30.** State Einstein's photoelectric equation and mention physical significance of each term involved in it.  
The wavelength of incident light is  $4000\text{\AA}$ . Calculate the energy of incident photon.
- Q. 31.** State any four uses of Van de Graaff generator.  
In a parallel plate air capacitor, intensity of electric field is changing at the rate of  $2 \times 10^{11}$  V/ms. If area of each plate is  $20\text{ cm}^2$ , calculate the displacement current.





## Our Website



## Our Channel



 On Solution Buddy, You Will Get:

- ✓ Exercise solutions for Class 8–12
- ✓ Previous Year Question Papers (10th & 12th)
- ✓ Free Textbook Downloads
- ✓ Practical Solutions (Class 10, 11 & 12)
- ✓ Water Security Exercise & Activity Solution
- ✓ Defence Studies Exercise Solution
- 👉 Website: [solutionbuddy.netlify.app](https://solutionbuddy.netlify.app)
- 👉 YouTube: [youtube.com/@solutionbuddy](https://youtube.com/@solutionbuddy)



# Solution Buddy

