

2. Finding Inverse of a Matrix

A. Activities

1. If $\begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix} X = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$, find X by using the following activity

by $R_2 + R_1$

and $R_3 - R_1$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 3 & 5 \\ 0 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix}, \text{ By } \left(\frac{1}{3}\right)R_2$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 5/3 \\ 0 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \text{ By } R_1 - 2R_2$$

$$\begin{bmatrix} 1 & 0 & -1/3 \\ 0 & 1 & 5/3 \\ 0 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}, \text{ By } R_1 + \frac{1}{3}R_3, R_2 - \frac{5}{3}R_3$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} -7/3 \\ -1/3 \\ 2 \end{bmatrix}$$

$$\therefore X = \begin{bmatrix} -1/3 \\ 2 \\ 2 \end{bmatrix}$$

2. If $A = \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix}$, then show that $A^{-1} = \frac{1}{6} (A - 5I)$ by completing the following activity.

Consider $AA^{-1} = I$,

$$\therefore \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \frac{1}{4} R_1 \rightarrow \begin{bmatrix} 1 & 5/4 \\ 2 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1/4 & 0 \\ 0 & 1 \end{bmatrix}$$

$$R_2 - 2R_1 \rightarrow \begin{bmatrix} 1 & 5/4 \\ 0 & 3/2 \end{bmatrix} A^{-1} = \begin{bmatrix} 1/4 & 0 \\ -1/2 & 1 \end{bmatrix}$$

$$\frac{-2}{3} R_2 \rightarrow \begin{bmatrix} 1 & 5/4 \\ 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1/4 & 0 \\ 1/3 & 2/3 \end{bmatrix}$$

$$R_1 - \frac{5}{4}R_2, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} \frac{2}{3} & \frac{2}{3} \\ \frac{1}{3} & -\frac{2}{3} \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{6} \begin{bmatrix} -1 & 5 \\ 2 & -4 \end{bmatrix} \dots\dots\dots (i)$$

$$\begin{aligned} \text{Now } \frac{1}{6}(A - 5I) &= \frac{1}{6} \left(\begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix} - 5 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \\ &= \frac{1}{6} \begin{bmatrix} -1 & 5 \\ 2 & -4 \end{bmatrix} \dots\dots\dots (ii) \end{aligned}$$

$$\therefore \text{ From (i) and (ii) } A^{-1} = \frac{1}{6}(A - 5I)$$

3. Find the inverse of $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ by completing the following activity.

Consider $A.A^{-1} = I$

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad R_2 - R_1 \text{ and } R_3 - 2R_1$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & 2 \\ \square & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -2 & 0 & \square \end{bmatrix}, \text{ By } -R_2$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ \square & 0 & 0 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 0 \\ -2 & 0 & \square \end{bmatrix}, R_1 - 2R_2$$

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & +1 & -2 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} -1 & 2 & 0 \\ 1 & -1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

$R_1 - 7R_3$ and $R_2 + 2R_3$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 13 & 2 & -7 \\ 3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 13 & 2 & -7 \\ 3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

4. Carry out the following activity

$$\text{If } A = \begin{bmatrix} 1 & -1 & 2 \\ -2 & 3 & 5 \\ -2 & 0 & -1 \end{bmatrix}$$

$$M_{11} = \dots -3 \dots$$

$$M_{12} = \dots 12 \dots$$

$$M_{13} = \dots 6 \dots$$

$$M_{21} = \dots 1 \dots$$

$$M_{22} = \dots 3 \dots$$

$$M_{23} = \dots -2 \dots$$

$$M_{31} = \dots -11 \dots$$

$$M_{32} = \dots 9 \dots$$

$$M_{33} = \dots 1 \dots$$

$$A_{11} = (-1)^{1+1} M_{11} = \boxed{-3}$$

$$A_{12} = (-1)^{1+2} M_{12} = \boxed{-12}$$

$$A_{13} = (-1)^{1+3} \boxed{6} = 6$$

$$A_{21} = (-1)^{2+1} (1) = \boxed{-1}$$

$$A_{22} = (-1)^{2+2} \boxed{3} = 3$$

$$A_{23} = (-1)^{2+3} \boxed{-2} = \boxed{2}$$

$$A_{31} = (-1)^{3+1} \boxed{-11} = -11$$

$$A_{32} = (-1)^{3+2} \boxed{9} = -9$$

$$A_{33} = (-1)^{3+3} \boxed{1} = \boxed{1}$$

$$\text{Cofactor Matrix} = \begin{bmatrix} -3 & \boxed{-12} & \boxed{6} \\ \boxed{-1} & 3 & \boxed{2} \\ \boxed{-11} & \boxed{-9} & 1 \end{bmatrix}$$

$$\text{adjoint of } A = \begin{bmatrix} -3 & \boxed{-1} & -11 \\ \boxed{-12} & 3 & \boxed{-9} \\ 6 & \boxed{2} & 1 \end{bmatrix}$$

B. Solve the Following

Q.1. Find the inverse of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ by the elementary row transformations.

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$$

$$\therefore |A| = 1(7-20) - 2(7-10) + 3(4-2) = -13 + 6 + 6 = -1 \neq 0$$

$\therefore A^{-1}$ exists.

Consider $AA^{-1} = I$

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

By $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - 2R_1$, we get

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

By $R_2 \rightarrow (-1)R_2$, we get

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

By $R_1 \rightarrow R_1 - 2R_2$, we get

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} -1 & 2 & 0 \\ 1 & -1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

By $R_1 \rightarrow R_1 - 7R_3$ and $R_2 \rightarrow R_2 + 2R_3$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 13 & 2 & -7 \\ -3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 13 & 2 & -7 \\ -3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

Q.2. Find the inverse of the matrix $\begin{bmatrix} 1 & 2 & 1 \\ 3 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}$ using adjoint method.

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}$$

$$\therefore |A| = 1(0-2) - 2(3-0) + 1(6-0) = -2-6+6 = -2 \neq 0$$

$\therefore A^{-1}$ exists.

$$A_{11} = (-1)^{1+1} M_{11} = (1) \begin{vmatrix} 0 & 1 \\ 2 & 1 \end{vmatrix} = 0-2 = -2$$

$$A_{12} = (-1)^{1+2} M_{12} = (-1) \begin{vmatrix} 3 & 1 \\ 0 & 1 \end{vmatrix} = -(3-0) = -3$$

$$A_{13} = (-1)^{1+3} M_{13} = (1) \begin{vmatrix} 3 & 0 \\ 0 & 2 \end{vmatrix} = 6-0 = 6$$

$$A_{21} = (-1)^{2+1} M_{21} = (-1) \begin{vmatrix} 2 & 1 \\ 2 & 1 \end{vmatrix} = -(2-2) = 0$$

$$A_{22} = (-1)^{2+2} M_{22} = (1) \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = (1-0) = 1$$

$$A_{23} = (-1)^{2+3} M_{23} = (-1) \begin{vmatrix} 1 & 2 \\ 0 & 2 \end{vmatrix} = -(2-0) = -2$$

$$A_{31} = (-1)^{3+1} M_{31} = (1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix} = 2-0 = 2$$

$$A_{32} = (-1)^{3+2} M_{32} = (-1) \begin{vmatrix} 1 & 1 \\ 3 & 1 \end{vmatrix} = -(1-3) = 2$$

$$A_{33} = (-1)^{3+3} M_{33} = (1) \begin{vmatrix} 1 & 2 \\ 3 & 0 \end{vmatrix} = (0-6) = -6$$

$$\therefore \text{Cofactor matrix} = \begin{bmatrix} -2 & -3 & 6 \\ 0 & 1 & -2 \\ 2 & 2 & -6 \end{bmatrix}$$

$$\therefore \text{adj } A = [\text{Cofactor matrix}]^T = \begin{bmatrix} -2 & 0 & 2 \\ -3 & 1 & 2 \\ 6 & -2 & -6 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{-2} \begin{bmatrix} -2 & 0 & 2 \\ -3 & 1 & 2 \\ 6 & -2 & -6 \end{bmatrix}$$

Q.3. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, Show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .

$$\begin{aligned} \text{Let } A^2 - 5A + 7I &= \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} - 5 \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} + 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 9-1 & 3+2 \\ -3-2 & -1+4 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \\ &= \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \\ &= \begin{bmatrix} 8-15+7 & 5-5+0 \\ -5+5+0 & 3-10+7 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0 \end{aligned}$$

$$\therefore \boxed{A^2 - 5A + 7I = 0} \text{ --- (1)}$$

$$\text{Now } |A| = \begin{vmatrix} 3 & 1 \\ -1 & 2 \end{vmatrix} = 6 - (-1) = 7 \neq 0$$

$\therefore A^{-1}$ exists.

$$A_{11} = (-1)^{1+1} M_{11} = (-1)^2 \cdot 2 = 2$$

$$A_{12} = (-1)^{1+2} M_{12} = (-1)^3 \cdot (-1) = 1$$

$$A_{21} = (-1)^{2+1} M_{21} = (-1)^3 \cdot 1 = -1$$

$$A_{22} = (-1)^{2+2} M_{22} = 1 \cdot (3) = 3$$

$$\therefore \text{Cofactor matrix} = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix}$$

$$\therefore \text{Adj}(A) = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \cdot \text{Adj}(A) = \frac{1}{7} \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$$

Sign of Teacher :