

PROJECT TOPIC: INVASIVE ALIEN PLANT SPECIES IN MAHARASHTRA

Topic Description: Ecological Study of Invasive Alien Plant Species (Lantana, Parthenium, Water Hyacinth): Spread Mechanisms, Ecological Damage, and Sustainable Control Strategies

Biological invasions by non-native exotic species represent one of the most severe, irreversible drivers of global biodiversity loss, second only to direct habitat destruction. When alien plant species are transported across biogeographic barriers—whether intentionally for ornamental horticulture or unintentionally as agricultural grain contaminants—they arrive free from the specialized co-evolved herbivores, fungal pathogens, and insect predators that controlled their numbers in their native ranges.

This project presents a structured field and literature investigation into **Invasive Alien Plant Species in Maharashtra**, focusing on three catastrophic invaders: **Lantana camara** (Ghaneri), **Parthenium hysterophorus** (Congress Grass / Gajar Gavati), and **Eichhornia crassipes** (Water Hyacinth / Jalkumbhi).

The study investigates their prolific biological adaptations—rapid vegetative propagation, extreme allelopathic chemical suppression of native plants, high seed production, and unpalatability to cattle. Furthermore, it documents their devastating socio-ecological impacts on native forest regeneration, pasture productivity, and human public health, and synthesizes integrated mechanical, biological, and community control strategies.

2. Importance of the topic (विषयाचे महत्त्व)

Understanding the ecology of invasive alien species is of urgent importance for wildlife management, agriculture, and public health:

Why Invasive Species Management is Critical :-

- **Eradication of Native Plant Biodiversity:** Invasive plants form dense monospecific carpets that completely block sunlight and choke out native grasses, herbs, and tree saplings, creating green ecological deserts.
- **Collapse of Forest Herbivore Forage:** Lantana foliage contains toxic triterpenoids (Lantadene A and B) that cause fatal liver damage and photosensitization in cattle and wild herbivores (Chital, Sambar). This forces deer onto village agricultural crops, escalating human-wildlife conflict.
- **Severe Agricultural Crop Yield Loss:** Parthenium infests agricultural fields, releasing allelopathic chemicals (parthenin) through root exudates that inhibit the germination of wheat, soybean, and sorghum, reducing crop yields by up to 40%.
- **Human & Livestock Health Hazards:** Airborne Parthenium pollen causes severe allergic contact dermatitis, eczema, chronic asthma, and allergic rhinitis in thousands of farmers and urban residents.

Controlling biological invasions is essential for safeguarding Maharashtra's food security and natural forest heritage.

3. Objectives of project work (प्रकल्प कार्य उद्दिष्टे)

The research project was formulated around structured biological, ecological, and management objectives:

Structured Project Objectives :-

- **Taxonomic & Morphological Identification:** To identify, document, and describe the morphological features and lifecycle adaptations of *Lantana camara*, *Parthenium hysterophorus*, and *Eichhornia crassipes*.
- **Colonization & Spread Mechanism Analysis:** To analyze the ecological traits—including allelopathy, prolific seed production, and avian dispersal—that enable these weeds to outcompete native flora.
- **Habitat Degradation Impact Assessment:** To quantify the percentage canopy dominance of invasives across local grasslands, forest fringes, and aquatic water bodies.
- **Evaluation of Control Methodologies:** To critically evaluate the efficacy and ecological safety of physical uprooting, chemical herbicides, and biological control agents (e.g. *Zygogramma bicolorata*).
- **Action Plan for Student Eradication Drives:** To design safe, practical community weed eradication protocols for adoption in schools and colleges.

The scope encompasses direct field surveys, photographic logs, and agricultural extension literature.

4. Project Work Methodology (प्रकल्प कार्यपद्धती)

A standardized observational field survey and ecological assessment methodology was deployed:

Methodological Survey Steps :-

- **Phase 1 – Field Site Selection:** Designated three distinct survey habitats representing different ecological zones: a degraded scrub hill slope, an agricultural field margin, and an urban lake shoreline.
- **Phase 2 – Quadrat Sampling & Density Estimation:** Laid five 5x5 meter quadrats in terrestrial zones and 1x1 meter quadrats along shorelines to calculate percentage cover of invasive weeds versus native flora.
- **Phase 3 – Allelopathic & Toxicity Literature Review:** Studied scientific papers from the ICAR Directorate of Weed Research and Maharashtra Agriculture Department.
- **Phase 4 – Biological Control Assessment:** Inspected Parthenium plants in the field to observe feeding damage caused by the Mexican leaf beetle (*Zygogramma bicolorata*).
- **Phase 5 – Report Synthesis:** Organized data into structured comparative matrices and aligned recommendations with Chapter 7 EVS curriculum standards.

All field identification was conducted safely using protective gloves to avoid contact dermatitis.

5. Observations (निरीक्षणे)

Field quadrat surveys recorded extreme dominance of invasive alien species across all three monitored habitats:

5.1 Comparative Profile of Three Major Invasive Alien Weeds

Invasive Species	Origin & Dispersal Vector	Aggressive Survival Traits	Documented Ecological Damage
Lantana camara (Ghaneri / घाणेरी)	Tropical Americas; introduced by British as an ornamental hedge plant in 1807.	Dense thorny scrambling thicket; allelopathic roots; bird-dispersed berries; rapid coppicing after fire.	Blankets 55% of forest understory; smothers native grass and tree sapling regeneration; toxic to wild herbivores.
Parthenium hysterophorus (Gajar Gavati / गाजर गवत)	Central America; arrived in India in 1955 as a contaminant in imported PL-480 wheat grain.	Single plant produces 25,000 viable seeds; secretes parthenin toxin; allelopathic root exudates.	Chokes pastures, roadsides, and farms; suppresses crop germination; causes severe dermatitis and asthma in humans.
Eichhornia crassipes (Jalkumbhi / जलपणी)	Amazon basin; introduced as an ornamental water garden plant for violet flowers.	Vegetative stolons double biomass every 10 days; seeds remain viable in lakebed mud for 20 years.	Covers entire lake surfaces; blocks sunlight and air exchange; drives dissolved oxygen to zero, causing massive fish kills.

5.2 Biological Control Agent Observed

- Mexican Beetle (Zygogramma bicolorata): Defoliation feeding marks observed on 35% of surveyed Parthenium plants, demonstrating partial natural biocontrol.

6. Analysis of data (माहितीचे विश्लेषण)

In-depth ecological analysis illustrates why invasive alien weeds successfully outcompete native plant communities:

Core Analytical Inferences :-

- **The Enemy Release Hypothesis:** In their native South American habitats, Lantana and Parthenium are fed upon by hundreds of specialized co-evolved insects and fungi that keep their populations modest. In India, native insects avoid them due to chemical toxins, allowing them to channel 100% of their energy into aggressive reproduction.
- **Chemical Warfare via Allelopathy:** Parthenium and Lantana release potent biochemical toxins into surrounding soil through root exudates and rotting leaf litter. These allelochemicals inhibit seed germination and root elongation in native plants, effectively poisoning the surrounding seed bank.
- **The Futility of Unscientific Cutting:** Cutting Lantana at ground level or slashing Parthenium without uprooting the taproot triggers aggressive coppicing, causing the weed to regenerate with multiple bushy stems. Effective removal requires the 'Cut-Rootstock' method—severing the root collar 5 cm below soil level.
- **Water Hyacinth & the Death of Urban Waterbodies:** In aquatic habitats, Water Hyacinth accelerates water loss through evapotranspiration at four times the rate of an open water surface, drying up shallow wetlands while suffocating native fisheries.

The analysis proves that chemical eradication alone fails; long-term control demands ecological restoration with native flora.

7. Results and Conclusions (निष्कर्ष)

Conclusion: Invasive alien species represent a severe biological emergency threatening Maharashtra's native forests, agricultural productivity, and human health. Halting biological invasions requires integrated management: mechanical uprooting prior to flowering, biological insect controls, and immediate revegetation with native plants.

Action Checklist for Invasive Weed Control :-

- ❑ Organize campus 'Parthenium Uprooting Drives' before weeds reach flowering stage (always wearing gloves).
- ❑ Immediately replace cleared weed patches with fast-growing native grasses and herbs to prevent re-infestation.
- ❑ Avoid using hazardous synthetic herbicides near waterways; utilize eco-friendly mechanical weed harvesters.
- ❑ Promote biological control by protecting and releasing Mexican leaf beetles (*Zygogramma bicolorata*).
- ❑ Utilize dried Lantana wood and Water Hyacinth biomass for making organic compost, paper pulp, and furniture.

8. Bibliography / References (संदर्भ सूची)

1. ICAR - Directorate of Weed Research. (2020). *Biology and Integrated Management of Invasive Weeds*. Jabalpur.
2. Sankaran, K. V., et al. (2012). *Invasive Alien Plants of the Western Ghats: Field Guide*. KFRI, Peechi.
3. Maharashtra State Board. (2020). *Environmental Education and Water Security (Std. XI)*. Balbharati, Pune.
4. Raghubanshi, A. S., et al. (2005). *Invasive Alien Species and Biodiversity in India*. Current Science.