

The Gao Constant of Generalized Dihedral Groups with an Abelian p -Group Kernel, p Odd

Abstract

Let A be a nontrivial finite abelian p -group for an odd prime p , and let $\text{Dih}(A) = A \rtimes_{-1} C_2$ be the generalized dihedral group in which the nonidentity element of C_2 acts by inversion. We determine its Gao constant:

$$\mathsf{E}(\text{Dih}(A)) = 2|A| + \mathsf{D}(A) = |\text{Dih}(A)| + \mathsf{d}(\text{Dih}(A)).$$

Thus, if $A = \bigoplus_{i=1}^r C_{p^{\lambda_i}}$, then

$$\mathsf{E}(\text{Dih}(A)) = 2|A| + 1 + \sum_{i=1}^r (p^{\lambda_i} - 1).$$

The lower bound follows from the known small Davenport constant of $\text{Dih}(A)$. For the upper bound, we divide according to the number of reflections in the sequence. The two extreme ranges are handled by prescribed-length weighted zero-sum results and a plus-minus Davenport bound. In the remaining range, a weighted structural theorem either gives the required subsequence directly or forces many rotations into a proper subgroup. A simultaneous induction on the subgroup order completes the proof.

2020 Mathematics Subject Classification. 11B75; 20K01.

Keywords. Gao constant; product-one sequence; generalized dihedral group; Davenport constant; weighted zero-sum sequence.

1 Introduction

Let G be a finite group. Its *Gao constant* $\mathsf{E}(G)$ is the least integer L such that every sequence of length at least L over G contains a subsequence of exactly $|G|$ terms that can be ordered to have product one. The small Davenport constant $\mathsf{d}(G)$ is the largest length of a sequence over G having no nonempty product-one subsequence. The standard padding construction gives

$$\mathsf{E}(G) \geq |G| + \mathsf{d}(G).$$

For finite abelian groups, Gao proved equality [3]. Zhuang and Gao conjectured that the same equality holds for every finite group [10]. It is known for ordinary dihedral groups; see [1, 4, 10]. Exact calculations for other nearby nonabelian families include $D_{2n} \times C_2$ [2].

Let p be an odd prime, let

$$A = \bigoplus_{i=1}^r C_{p^{\lambda_i}}, \quad 1 \leq \lambda_1 \leq \cdots \leq \lambda_r,$$

and write

$$\text{Dih}(A) = A \rtimes_{-1} C_2,$$

where the nonidentity element of C_2 acts on A by inversion. Our main result is the following.

Theorem 1.1. *Let $Q = |A|$. Then*

$$E(\text{Dih}(A)) = 2Q + D(A) = 2Q + 1 + \sum_{i=1}^r (p^{\lambda_i} - 1).$$

Equivalently,

$$E(\text{Dih}(A)) = |\text{Dih}(A)| + d(\text{Dih}(A)).$$

The equivalence uses the identity $d(\text{Dih}(A)) = D(A)$ of Godara, Joshi, and Mazumdar [5]. Their work also gives the upper bound $E(\text{Dih}(A)) \leq 3|A|$. When A is cyclic, this upper bound meets the standard lower bound and recovers the known dihedral formula. For a noncyclic abelian p -group one has $D(A) < |A|$, so that bound does not determine the exact value. More recently, Zhao and Wang proved a uniform $3|G|/2$ bound for noncyclic groups whose order is not divisible by four, together with an equality characterization [9]. Applied here, their result gives a strict upper bound below $3|A|$ when A is noncyclic, but it still does not give the value in [Theorem 1.1](#).

The theorem extends the cyclic-kernel formula to all invariant-factor decompositions of finite abelian p -groups with p odd.

Example 1.2. Let $A = C_3^2$, with basis e_1, e_2 . Then $|A| = 9$ and $D(A) = 5$, so [Theorem 1.1](#) gives

$$E(\text{Dih}(C_3^2)) = 23.$$

The lower bound can be seen explicitly. Take two copies of the rotation $(e_1, 0)$, two copies of $(e_2, 0)$, and one reflection. This five-term sequence is product-one-free: a product-one subsequence has an even number of reflections, while the four rotations form a zero-sum-free sequence in C_3^2 . Appending seventeen identity rotations gives a sequence of length 22 with no product-one subsequence of length 18.

We outline the upper bound. Let S have length $2|A| + D(A)$, and let a be its number of reflections. If $a \leq 1$, there are enough rotations to apply an unweighted prescribed-length zero-sum theorem. If $a \geq D(A) + 1$, pairing terms within the two cosets reduces the problem to a plus-minus weighted zero-sum problem in A . In the intermediate range, the structural alternative in a theorem of Gryniewicz, Marchan, and Ordaz [6] either produces the desired subsequence or shows that many rotations lie in a proper subgroup $K < A$. The latter situation is resolved by simultaneous induction on $|K|$. The auxiliary induction statement is needed because one branch translates all rotation vectors; only an all-rotation conclusion can be pulled back through that translation without changing the product-one condition.

The proof uses Olson's formula for the Davenport constant of an abelian p -group and a plus-minus argument over a field of odd characteristic. We do not treat mixed-prime kernels or 2-groups in this paper.

2 Preliminaries

A sequence over a set X is an element of the free abelian monoid $\mathcal{F}(X)$. We write it multiplicatively, even when X is an additively written group. Thus

$$S = x_1 \cdot \dots \cdot x_\ell, \quad |S| = \ell.$$

Subsequences are selected by positions, so equal values occurring in different positions remain available separately. We write $T \mid S$ when T is a subsequence and ST^{-1} for the complementary

sequence. A sequence over a group is *product one* if its terms can be ordered so that their product is the identity.

For an abelian group B , the Davenport constant $D(B)$ is the least integer L such that every sequence of length L over B has a nonempty zero-sum subsequence. Thus $D(B) = d(B) + 1$. If $W \subseteq \mathbb{Z}$ is nonempty, the weighted Davenport constant $D_W(B)$ is defined analogously, with a coefficient from W assigned to each selected term. We write

$$D_{\pm}(B) = D_{\{-1,1\}}(B).$$

Throughout the paper, set

$$q_i = p^{\lambda_i}, \quad Q = |A| = \prod_{i=1}^r q_i, \quad D_A = D(A) = 1 + \sum_{i=1}^r (q_i - 1).$$

The formula for D_A is Olson's theorem [8]. In particular,

$$D_A \leq Q, \quad \exp(A) \mid Q, \quad D_A \text{ and } Q \text{ are odd.} \quad (1)$$

The first inequality follows by expanding $Q = \prod_i (1 + (q_i - 1))$.

We represent elements of $G = \text{Dih}(A)$ as pairs $(x, \varepsilon) \in A \times C_2$, with multiplication

$$(x, \varepsilon)(y, \delta) = (x + (-1)^{\varepsilon}y, \varepsilon + \delta). \quad (2)$$

Elements $(x, 0)$ are called *rotations*, and elements $(x, 1)$ are called *reflections*.

2.1 Realizing prescribed signs

The next elementary lemma is the bridge between weighted sums in A and product-one subsequences in G .

Lemma 2.1 (Realization of prescribed signs). *Suppose a selected sequence contains $2k > 0$ reflections and any number of rotations. Assign signs from $\{+1, -1\}$ to the terms so that exactly k reflections receive each sign; the signs on the rotations are arbitrary. Then the terms can be ordered so that their contributions to the A -coordinate in (2) have the assigned signs. Consequently, if the assigned signed vector sum is zero, the selected sequence is product one.*

If a selected sequence contains no reflections, it is product one if and only if the sum of its rotation vectors is zero.

Proof. Place the positively and negatively signed reflections alternately, starting with a positive one. In any ordered product, a term contributes its vector with sign $(-1)^t$, where t is the number of preceding reflections. Hence the chosen reflection ordering realizes the prescribed alternating signs.

The $2k$ reflections determine $2k + 1$ gaps: one before the first reflection, one after the last, and one between each consecutive pair. A rotation inserted in a gap preceded by an even number of reflections has sign $+1$; in a gap preceded by an odd number it has sign -1 . Since $k \geq 1$, gaps of both parities occur, and any number of rotations can be inserted in either type of gap. This realizes all prescribed rotation signs. The total number of reflections is even, so the C_2 -coordinate is zero; the A -coordinate is the prescribed signed sum. $\square$

2.2 Prescribed-length weighted sums

For a sequence T over a finite abelian group B , a nonempty weight set $W \subseteq \mathbb{Z}$, and an integer $n \leq |T|$, let

$$\Sigma_n^W(T) = \left\{ \sum_{j=1}^n w_j t_j : t_1, \dots, t_n \text{ occur in distinct positions of } T, w_j \in W \right\}.$$

For an integer n , write $nB = \{nx : x \in B\}$. We use the following two corollaries of Gryniewicz, Marchan, and Ordaz [6, Corollaries 1.2 and 1.3].

Theorem 2.2 (Gryniewicz–Marchan–Ordaz). *Let B be a finite abelian group, let $W \subseteq \mathbb{Z}$ be nonempty, let T be a sequence over B , and let $n \geq |B|$. If*

$$|T| \geq n + D_W(B) - 1,$$

then

$$\Sigma_n^W(T) \cap nB \neq \emptyset. \quad (3)$$

If, in addition, $\gcd(W) = 1$, then either

$$\Sigma_n^W(T) = B, \quad (4)$$

or there exist a proper subgroup $H < B$, elements $\alpha, \beta \in B$, and a subsequence $T' \mid T$ such that

$$|T'| \geq |T| - |B/H| + 2, \quad \text{supp}(T') \subseteq \alpha + H, \quad (5)$$

and

$$Wx \subseteq \beta + H \quad \text{for every } x \in \text{supp}(T'). \quad (6)$$

Two specializations will be used repeatedly. For $W = \{1\}$, one has $D_W(B) = D(B)$. For $W = \{-1, 1\}$, condition (6) says that x and $-x$ lie in the same H -coset for every term of T' . Thus $2x \in H$; if B/H has odd order, this implies $x \in H$.

The other external input is the following consequence of [5, Theorem 1.1]:

$$d(\text{Dih}(B)) = D(B) \quad \text{for every finite abelian group } B. \quad (7)$$

2.3 The lower bound and identity padding

Proposition 2.3. *For $G = \text{Dih}(A)$,*

$$E(G) \geq 2Q + D_A.$$

Proof. By (7), there is a product-one-free sequence U over G of length D_A . Append $2Q - 1$ copies of the identity. Every subsequence of length $2Q$ uses a nonempty part of U . If such a subsequence were product one, deleting the identity terms from a product-one ordering would give a nonempty product-one subsequence of U , a contradiction. $\square$

Lemma 2.4 (Identity padding). *Let Y be a sequence over G of length $2Q + D_A$. If at least D_A terms of Y are the identity rotation, then Y contains a product-one subsequence of length $2Q$.*

Proof. Let m_0 be the number of identity rotations. From the sequence of all nonidentity terms, successively remove nonempty product-one subsequences until the remainder R is product-one-free. Let P be the union of the removed subsequences. Concatenating one product-one ordering for each removed block makes P product one. By (7), $|R| \leq D_A$, and hence

$$|P| = (2Q + D_A - m_0) - |R| \geq 2Q - m_0.$$

Since $m_0 \geq D_A$, also $|P| \leq 2Q$. Therefore at most m_0 unused identity terms are needed to extend P to length $2Q$. $\square$

3 A plus-minus Davenport bound

The plus-minus Davenport constant has been studied systematically; see, for example, [7]. We need an estimate expressed in terms of the ordinary Davenport constant. The estimate below is known, but we include a short group-algebra proof because this precise form is used in each signed prescribed-length argument.

Proposition 3.1. *Let B be a nontrivial finite abelian p -group, where p is odd. Then*

$$D_{\pm}(B) \leq \frac{D(B) + 1}{2}.$$

Proof. Write

$$B = \bigoplus_{i=1}^s C_{p^{\mu_i}}, \quad r_i = p^{\mu_i}, \quad D_B = D(B) = 1 + \sum_{i=1}^s (r_i - 1).$$

Work in the group algebra $R = \mathbb{F}_p[B]$, write $[x]$ for the basis element indexed by $x \in B$, and let I be the augmentation ideal. If f_i is an invariant-factor generator and $y_i = [f_i] - [0]$, then

$$y_i^{r_i} = ([f_i] - [0])^{r_i} = [r_i f_i] - [0] = 0.$$

The resulting homomorphism

$$\mathbb{F}_p[y_1, \dots, y_s] / (y_1^{r_1}, \dots, y_s^{r_s}) \longrightarrow R$$

is surjective, and both sides have dimension $|B|$. It is therefore an isomorphism. Every monomial of total degree at least D_B vanishes, so

$$I^{D_B} = 0. \tag{8}$$

The number D_B is odd. Put $m = (D_B + 1)/2$, and take arbitrary terms $x_1, \dots, x_m \in B$. For every j ,

$$[x_j] + [-x_j] - 2[0] = [-x_j]([x_j] - [0])^2 \in I^2.$$

Hence the product of these m elements lies in $I^{2m} = I^{D_B+1} = 0$. The coefficient of $[0]$ in its expansion is

$$\sum_{\substack{\varepsilon \in \{-1, 0, 1\}^m \\ \sum_j \varepsilon_j x_j = 0}} (-2)^{|\{j: \varepsilon_j = 0\}|}.$$

If the sequence $x_1 \dots x_m$ had no nonempty plus-minus zero-sum subsequence, the only contributing vector would be $\varepsilon = 0$. The displayed coefficient would then be $(-2)^m \neq 0$ in $\mathbb{F}_p$, contradicting the vanishing of the product. Thus every sequence of length m has a nonempty plus-minus zero-sum subsequence. $\square$

4 Reduction by the number of reflections

Let S be an arbitrary sequence over $G = \text{Dih}(A)$ of length

$$|S| = 2Q + D_A. \quad (9)$$

Let a be the number of reflections in S , and put

$$b = 2Q + D_A - a$$

for the number of rotations. For a subgroup $K \leq A$ and a sequence Y over G , let $m_K(Y)$ denote the number of rotations $(x, 0)$ of Y with $x \in K$.

4.1 At most one reflection

Proposition 4.1. *If $a \leq 1$, then S contains a product-one subsequence of length $2Q$ consisting entirely of rotations.*

Proof. There are at least

$$b \geq 2Q + D_A - 1 = 2Q + D(A) - 1$$

rotations. Apply [Theorem 2.2](#), in its existence form (3), to the rotation sequence over A , with weight set $\{1\}$ and target $n = 2Q$. The target satisfies $2Q \geq |A|$, and the displayed inequality is the required length hypothesis. Since $\exp(A) \mid Q$, one has $2QA = \{0\}$. Thus there are exactly $2Q$ rotations with ordinary sum zero. $\square$

4.2 Many reflections

Proposition 4.2. *If $a \geq D_A + 1$, then S contains a product-one subsequence of length $2Q$.*

Proof. Pair reflections with reflections and rotations with rotations. Since $2Q + D_A$ is odd, exactly one term is left unpaired, and the number of pairs is

$$\left\lfloor \frac{a}{2} \right\rfloor + \left\lfloor \frac{b}{2} \right\rfloor = Q + \frac{D_A - 1}{2}.$$

Assign to a reflection pair $(x, 1) \cdot (y, 1)$ the label $x - y$, and to a rotation pair $(u, 0) \cdot (v, 0)$ the label $u + v$. By [Proposition 3.1](#), the sequence of pair labels has length at least

$$Q + D_{\pm}(A) - 1.$$

Apply (3) in A , with weight set $\{-1, 1\}$ and target $Q = |A|$. Since $QA = \{0\}$, we obtain exactly Q pairs and a sign for each selected pair such that the signed sum of their labels is zero.

At most $\lfloor b/2 \rfloor \leq Q - 1$ of all pairs are rotation pairs, because $a \geq D_A + 1$. Hence at least one selected pair is a reflection pair. For a reflection pair with selected weight ε , assign signs ε and $-\varepsilon$ to its two reflections. For a rotation pair, assign the selected weight to both rotations. The selected reflection signs are balanced, and the total signed vector sum is zero. By [Lemma 2.1](#), the $2Q$ selected terms have a product-one ordering. $\square$

4.3 The intermediate range

Assume now that

$$2 \leq a \leq D_A. \quad (10)$$

Let e be the largest even integer not exceeding a , and put

$$\ell = 2Q - e. \quad (11)$$

Then $e > 0$, $\ell \geq Q$, and $\ell \leq b$. Moreover,

$$b - \ell = \begin{cases} D_A, & a \text{ even,} \\ D_A - 1, & a \text{ odd,} \end{cases} \quad b - \ell \geq D_{\pm}(A) - 1. \quad (12)$$

Indeed, $e \leq D_A \leq Q$, and the final inequality follows from [Proposition 3.1](#).

Proposition 4.3 (Intermediate-range reduction). *Under (10), either S contains a product-one subsequence of length $2Q$, or there is a nonzero proper subgroup $K < A$ such that*

$$m_K(S) \geq b - [A : K] + 2. \quad (13)$$

Proof. Apply the structural part of [Theorem 2.2](#) to the sequence of all b rotation vectors, with ambient group A , weight set $\{-1, 1\}$, and target ℓ . The target condition $\ell \geq |A|$ and the length surplus are exactly (12).

If $\Sigma_{\ell}^{\{-1, 1\}}(S_{\text{rot}}) = A$, select any e reflections and assign half of them each sign. Let their signed vector sum be $t \in A$. Fullness supplies ℓ rotations with a signed sum of $-t$. The two selections are disjoint, and their total length is $e + \ell = 2Q$. The sign assignment on the reflections is balanced, so [Lemma 2.1](#) gives a product-one ordering.

Otherwise, [Theorem 2.2](#) gives a proper subgroup $K < A$ and a rotation subsequence of length at least $b - [A : K] + 2$ satisfying (6). For every one of its values x , the elements x and $-x$ lie in a common K -coset. Hence $2x \in K$. Since A/K has odd order, multiplication by two is injective on A/K , and therefore $x \in K$. This proves (13).

It remains to exclude $K = 0$ from the second alternative. In that case at least

$$b - Q + 2 = Q + D_A - a + 2 \geq D_A$$

rotations are the identity, so [Lemma 2.4](#) gives the required subsequence directly. $\square$

5 Subgroup descent in the intermediate range

Fix, for this section, a sequence S satisfying (9) and (10); retain its reflection and rotation counts a, b . For a sequence Y over G and a subgroup $K \leq A$, let $T_K(Y)$ be the sequence over $\text{Dih}(A/K)$ obtained by taking all reflections of Y , together with all rotations whose vectors lie outside K , and projecting their vectors modulo K . Rotations with vectors in K are omitted. Selections in $T_K(Y)$ always refer to the corresponding positions of Y .

We prove two statements simultaneously. For every proper subgroup $K < A$, let $\mathcal{P}_S(K)$ denote the assertion

$$m_K(S) \geq b - [A : K] + 2 \implies S \text{ has a product-one subsequence of length } 2Q.$$

Let $\mathcal{Q}(K)$ denote the following universal assertion: every sequence Y over G of length $2Q + D_A$, with the same counts a, b , has an all-rotation product-one subsequence of length $2Q$ provided that

1. $m_K(Y) \geq b - [A : K] + 2$, and
2. $T_K(Y)$ has no product-one subsequence containing a reflection.

The all-rotation conclusion is essential because the proof of $\mathcal{Q}(K)$ translates rotation vectors before descending to a smaller subgroup.

Theorem 5.1 (Subgroup descent). *For every proper subgroup $K < A$, both $\mathcal{P}_S(K)$ and $\mathcal{Q}(K)$ hold.*

5.1 Three subgroup estimates

Lemma 5.2 (Davenport inequality). *For every subgroup $K \leq A$,*

$$D_A \geq D(K) + D(A/K) - 1. \quad (14)$$

Proof. Take a zero-sum-free sequence in K of length $D(K) - 1$ and a zero-sum-free sequence in A/K of length $D(A/K) - 1$, and lift the latter arbitrarily to A . A zero-sum subsequence of the concatenation would project to a zero-sum subsequence of the quotient sequence. It therefore uses no lifted quotient term, after which it would be a zero-sum subsequence of the sequence in K . Thus the concatenation is zero-sum-free, giving (14). $\square$

Lemma 5.3 (Target-length estimate). *Let $0 < K < A$, put $I = [A : K]$ and $h = |K|$, so that $Q = Ih$. Then*

$$2Q - D_A - I + 2 \geq Q - I + 2 \geq h. \quad (15)$$

Proof. The first inequality is $D_A \leq Q$. For the second,

$$Q - I + 2 - h = Ih - I - h + 2 = (I - 1)(h - 1) + 1 > 0.$$

$\square$

Lemma 5.4 (Transitivity of concentration). *Let $H < K < A$, put $I = [A : K]$ and $J = [K : H]$, and let Y be a sequence with*

$$m_K(Y) \geq b - I + 2.$$

If at least $m_K(Y) - J + 2$ of the rotations counted by $m_K(Y)$ actually lie in H , then

$$m_H(Y) \geq b - [A : H] + 2.$$

Proof. Since $[A : H] = IJ$,

$$m_H(Y) \geq b - I - J + 4 \geq b - IJ + 2,$$

because $IJ - I - J + 2 = (I - 1)(J - 1) + 1 > 0$. $\square$

Lemma 5.5 (Correction of a quotient defect). *Let $K \leq A$. Suppose a selected sequence U has a positive even number of reflections and admits an ordering whose product has A -coordinate $z \in K$. Let C_0 be a disjoint rotation sequence, with prescribed signs, whose signed vector sum is $-z$. Then UC_0 is product one.*

Proof. Read from the fixed ordering of U the sign of each vector. The reflection signs are balanced, because they alternate and their number is positive and even. Together with the prescribed signs on C_0 , the total signed vector sum is zero. Apply Lemma 2.1 to the entire selected sequence. $\square$

5.2 A reflection-containing quotient block

Let $0 < K < A$, suppose the hypothesis of $\mathcal{P}_S(K)$ holds, and let C be the sequence of the $M = m_K(S)$ rotations of S whose vectors lie in K . Let B be the sequence of the remaining $c = b - M$ rotations, and put $T = T_K(S)$.

Proposition 5.6. *If T has a product-one subsequence containing a reflection, then either S has a product-one subsequence of length $2Q$, or there is a strict subgroup $H < K$ such that*

$$m_H(S) \geq b - [A : H] + 2.$$

Proof. Start with a reflection-containing product-one subsequence of T . Continue removing nonempty product-one subsequences from the current remainder until the final remainder R is product-one-free. Let U be the union of all removed blocks. Concatenating one product-one ordering for each block makes U product one in $\text{Dih}(A/K)$, and

$$|U| = a + c - |R|.$$

The sequence U contains a positive even number of reflections. By (7), applied to A/K ,

$$|R| \leq D(A/K).$$

Set

$$\tau = D_A - |R|.$$

By Lemma 5.2 and Proposition 3.1,

$$\tau \geq D(K) - 1 \geq D_{\pm}(K) - 1. \quad (16)$$

Lift the fixed quotient ordering of U to the corresponding terms of S . Its product has C_2 -coordinate zero and is trivial modulo K , so its A -coordinate is a defect $z \in K$. Put

$$m = M - \tau = M - D_A + |R|.$$

The concentration hypothesis and Lemma 5.3 give

$$\begin{aligned} m &\geq M - D_A \\ &\geq b - [A : K] + 2 - D_A \\ &= 2Q - a - [A : K] + 2 \\ &\geq 2Q - D_A - [A : K] + 2 \\ &\geq |K|. \end{aligned}$$

Equation (16) gives $M \geq m + D_{\pm}(K) - 1$. Thus the structural part of Theorem 2.2 applies to C in the ambient group K , with weight set $\{-1, 1\}$ and target m .

If $\Sigma_m^{\{-1, 1\}}(C) = K$, choose m rotations of C , with signs, whose signed sum is $-z$. Their positions are disjoint from those of U , because T contains no rotation from K . By Lemma 5.5, their union with the lift of U is product one. Its length is

$$\begin{aligned} |U| + m &= a + c - |R| + M - D_A + |R| \\ &= a + b - D_A = 2Q. \end{aligned}$$

If the spectrum is not all of K , the structural alternative gives a strict subgroup $H < K$ and at least $M - [K : H] + 2$ terms of C satisfying (6). For each such vector x , the elements x and $-x$ lie in a common H -coset, so $2x \in H$. The quotient K/H has odd order, and hence $x \in H$. Now Lemma 5.4 gives $m_H(S) \geq b - [A : H] + 2$. $\square$

5.3 The rotation-only quotient case

For $\alpha \in A$ and a sequence Y over G , define $\tau_\alpha(Y)$ by replacing every rotation $(v, 0)$ with $(v - \alpha, 0)$, leaving every reflection unchanged. This is a transformation of the values attached to the positions; it is not a group homomorphism.

Lemma 5.7 (Persistence of the quotient obstruction). *Let $H < K \leq A$ and $\alpha \in K$. If $T_K(Y)$ has no product-one subsequence containing a reflection, then $T_H(\tau_\alpha(Y))$ has no such subsequence.*

Proof. Suppose that a reflection-containing product-one subsequence existed in $T_H(\tau_\alpha(Y))$, and fix a product-one ordering. Project this ordered block further modulo K . A selected rotation whose original vector v lies in K has translated vector $v - \alpha \in K$, so it becomes the identity modulo K ; delete all such identity rotations from the ordering. A selected rotation whose original vector lies outside K has the same image modulo K before and after the translation. Reflections are unchanged. The remaining ordered block is a reflection-containing product-one subsequence of $T_K(Y)$, a contradiction. $\square$

Proposition 5.8. *Let $0 < K < A$. Suppose that Y has length $2Q + D_A$, has the same counts a, b as S , satisfies*

$$m_K(Y) \geq b - [A : K] + 2,$$

and $T_K(Y)$ has no product-one subsequence containing a reflection. Then either

1. *Y has an all-rotation product-one subsequence of length $2Q$, or*
2. *there are $H < K$ and $\alpha \in K$ such that $\tau_\alpha(Y)$ satisfies the hypotheses of $\mathcal{Q}(H)$.*

Proof. Let C be the sequence of the $M = m_K(Y)$ rotations in K , and let B be the sequence of the remaining $c = b - M$ rotations. In the quotient A/K , successively remove nonempty zero-sum subsequences from the image of B until a zero-sum-free remainder B_0 remains. Let $B' = BB_0^{-1}$, using the corresponding original positions, and put $r_0 = |B_0|$. The vector sum of B' is an element

$$z = \sigma(B') \in K. \tag{17}$$

The quotient sequence formed by all a reflections and the terms of B_0 is product-one-free. Indeed, a rotation-only product-one subsequence would give a nonempty zero-sum subsequence of B_0 , and a product-one subsequence containing a reflection is excluded by the hypothesis on $T_K(Y)$. Hence, by (7),

$$a + r_0 \leq D(A/K). \tag{18}$$

Set

$$d_0 = D_A - a - r_0.$$

Using Lemma 5.2 and (18),

$$d_0 \geq D_A - D(A/K) \geq D(K) - 1. \tag{19}$$

Put $m = M - d_0$. The concentration hypothesis gives

$$\begin{aligned} m &\geq b - [A : K] + 2 - (D_A - a - r_0) \\ &= 2Q - [A : K] + r_0 + 2 \\ &\geq 2Q - [A : K] + 2 \geq |K|, \end{aligned}$$

where the last inequality follows from Lemma 5.3. Equation (19) gives $M \geq m + D(K) - 1$. Apply the structural part of Theorem 2.2 to C in K , with weight set $\{1\}$ and target m .

Suppose first that $\Sigma_m^{\{1\}}(C) = K$. Positional complementation inside C gives

$$\Sigma_{d_0}^{\{1\}}(C) = \sigma(C) - \Sigma_m^{\{1\}}(C) = K.$$

Choose a d_0 -term subsequence $D_0 \mid C$ with $\sigma(D_0) = \sigma(C) + z$. The rotation sequences CD_0^{-1} and B' have disjoint positions and total vector sum

$$\sigma(C) - \sigma(D_0) + \sigma(B') = -z + z = 0.$$

Their total length is

$$\begin{aligned} |CD_0^{-1}| + |B'| &= M - d_0 + c - r_0 \\ &= b - (D_A - a - r_0) - r_0 = 2Q. \end{aligned}$$

This proves the first alternative.

If the spectrum is not all of K , the structural alternative gives a strict subgroup $H < K$, an element $\alpha \in K$, and at least $M - [K : H] + 2$ terms of C whose vectors lie in $\alpha + H$. After applying τ_α , these rotations lie in H , and [Lemma 5.4](#) gives

$$m_H(\tau_\alpha(Y)) \geq b - [A : H] + 2.$$

The translation preserves the length and the counts a, b , while [Lemma 5.7](#) preserves the absence of a reflection-containing quotient block. Thus $\tau_\alpha(Y)$ satisfies the hypotheses of $\mathcal{Q}(H)$. $\square$

Lemma 5.9 (Pullback through a rotation translation). *If the positions of an all-rotation subsequence of length $2Q$ form a product-one sequence in $\tau_\alpha(Y)$, then the same positions form a product-one sequence in Y .*

Proof. Pulling the selected rotations back adds α to each vector, so the total vector sum changes by $2Q\alpha = 0$, using $\exp(A) \mid Q$. There are no reflections, so no alternating signs occur. $\square$

5.4 Simultaneous induction

Proof of Theorem 5.1. We use strong induction on the order of the subgroup.

For $K = 0$, the hypothesis of $\mathcal{P}_S(0)$ says that the number m_0 of identity rotations satisfies

$$m_0 \geq b - Q + 2 = Q + D_A - a + 2 \geq D_A.$$

Thus $\mathcal{P}_S(0)$ follows from [Lemma 2.4](#).

To prove $\mathcal{Q}(0)$, let Y satisfy its hypotheses, and let m_0 again be the number of identity rotations. The sequence $T_0(Y)$ consists of all reflections and all nonidentity rotations. Successively remove nonempty product-one subsequences from $T_0(Y)$ until a product-one-free remainder remains, and let P be the union of the removed blocks. The quotient obstruction in the definition of $\mathcal{Q}(0)$ forces every removed block to consist only of rotations. Hence P is an all-rotation product-one sequence. The terminal remainder has length at most D_A , so

$$|P| \geq (2Q + D_A - m_0) - D_A = 2Q - m_0.$$

Moreover, the concentration hypothesis gives $m_0 \geq b - Q + 2$, and therefore

$$|P| \leq b - m_0 \leq Q - 2 < 2Q.$$

There are enough unused identity rotations to extend P to an all-rotation product-one subsequence of length $2Q$. This proves $\mathcal{Q}(0)$.

Now let $0 < K < A$, and assume both statements are known for every strict subgroup of K . To prove $\mathcal{P}_S(K)$, consider $T_K(S)$. If it has a product-one subsequence containing a reflection, [Proposition 5.6](#) either gives the required subsequence immediately or gives a strict subgroup $H < K$ satisfying the hypothesis of $\mathcal{P}_S(H)$. The induction hypothesis applies in the latter case. If $T_K(S)$ has no such subsequence, [Proposition 5.8](#) either gives an all-rotation subsequence of length $2Q$ directly or gives $H < K$ and $\alpha \in K$ such that $\tau_\alpha(S)$ satisfies the hypotheses of $\mathcal{Q}(H)$. The induction hypothesis gives the required all-rotation subsequence after translation, and [Lemma 5.9](#) transfers it back to S .

Finally, let Y satisfy the hypotheses of $\mathcal{Q}(K)$. Its quotient obstruction already rules out the reflection-containing case. Apply [Proposition 5.8](#). The first alternative is the desired conclusion. In the second, the translated sequence satisfies $\mathcal{Q}(H)$ for a strict subgroup $H < K$; apply the induction hypothesis and then [Lemma 5.9](#). $\square$

6 Completion of the proof

Proof of [Theorem 1.1](#). The lower bound is [Proposition 2.3](#). For the upper bound, let S be a sequence of length $2Q + D_A$, and let a be its number of reflections. If $a \leq 1$, apply [Proposition 4.1](#). If $a \geq D_A + 1$, apply [Proposition 4.2](#). In the remaining range $2 \leq a \leq D_A$, [Proposition 4.3](#) either gives the required subsequence directly or yields a nonzero proper subgroup $K < A$ satisfying the hypothesis of $\mathcal{P}_S(K)$. [Theorem 5.1](#) then gives a product-one subsequence of length $2Q = |G|$. Hence $E(G) \leq 2Q + D_A$, and equality follows. $\square$

Corollary 6.1. *If $A = C_{p^k}^r$, where p is odd, then*

$$E(\text{Dih}(A)) = 2p^{kr} + r(p^k - 1) + 1.$$

In particular,

$$E(\text{Dih}(C_p^r)) = 2p^r + r(p - 1) + 1.$$

Proof. Substitute $|A| = p^{kr}$ and $D(C_{p^k}^r) = 1 + r(p^k - 1)$ in [Theorem 1.1](#). $\square$

References

- [1] Jared Bass. Improving the Erdős–Ginzburg–Ziv theorem for some non-abelian groups. *Journal of Number Theory*, 126(2):217–236, 2007. doi: 10.1016/j.jnt.2006.10.020.
- [2] F. E. Brochero Martínez, A. Lemos, B. K. Moriya, and S. Ribas. The main zero-sum constants over $D_{2n} \times C_2$. *SIAM Journal on Discrete Mathematics*, 37(3):1496–1508, 2023. doi: 10.1137/22M1496013.
- [3] Weidong Gao. A combinatorial problem on finite abelian groups. *Journal of Number Theory*, 58(1):100–103, 1996. doi: 10.1006/jnth.1996.0067.
- [4] Weidong Gao and Zaiping Lu. The Erdős–Ginzburg–Ziv theorem for dihedral groups. *Journal of Pure and Applied Algebra*, 212(2):311–319, 2008. doi: 10.1016/j.jpaa.2007.04.012.

- [5] Naveen K. Godara, Renu Joshi, and Eshita Mazumdar. Combinatorial invariants for certain classes of non-abelian groups. *Journal of Number Theory*, 283:44–63, 2026. doi: 10.1016/j.jnt.2025.11.011.
- [6] David J. Grynkiewicz, Luz Elimar Marchan, and Oscar Ordaz. A weighted generalization of two theorems of Gao. *The Ramanujan Journal*, 28(3):323–340, 2012. doi: 10.1007/s11139-011-9350-x.
- [7] Luz E. Marchan, Oscar Ordaz, and Wolfgang A. Schmid. Remarks on the plus-minus weighted Davenport constant. *International Journal of Number Theory*, 10(5):1219–1239, 2014. doi: 10.1142/S1793042114500225.
- [8] John E. Olson. A combinatorial problem on finite abelian groups, I. *Journal of Number Theory*, 1(1):8–10, 1969. doi: 10.1016/0022-314X(69)90021-3.
- [9] Yang Zhao and Guoqing Wang. On the Gao constant of groups of even order. *Journal of Number Theory*, 292:1–11, 2027. doi: 10.1016/j.jnt.2026.05.015. Published online in 2026.
- [10] Jujuan Zhuang and Weidong Gao. Erdős–Ginzburg–Ziv theorem for dihedral groups of large prime index. *European Journal of Combinatorics*, 26(7):1053–1059, 2005. doi: 10.1016/j.ejc.2004.06.014.