

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				

Pearson Edexcel Level 3 GCE

Paper reference	9MA0/32
-----------------	----------------

Mathematics

Advanced

PAPER 32: Mechanics

<p>You must have: Mathematical Formulae and Statistical Tables (Green), calculator</p>	<p>Total Marks</p>
---	--------------------

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 50. There are 5 questions.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P72131A

©2022 Pearson Education Ltd.

Q:1/1/1/1/



1. [In this question, position vectors are given relative to a fixed origin.]

At time t seconds, where $t > 0$, a particle P has velocity $\mathbf{v} \text{ m s}^{-1}$ where

$$\mathbf{v} = 3t^2\mathbf{i} - 6t^{\frac{1}{2}}\mathbf{j}$$

- (a) Find the speed of P at time $t = 2$ seconds. (2)
- (b) Find an expression, in terms of t , \mathbf{i} and \mathbf{j} , for the acceleration of P at time t seconds, where $t > 0$ (2)

At time $t = 4$ seconds, the position vector of P is $(\mathbf{i} - 4\mathbf{j}) \text{ m}$.

- (c) Find the position vector of P at time $t = 1$ second. (4)

a) sub $t=2$ into \mathbf{v} : $\mathbf{v} = 3(2)^2\mathbf{i} - 6(2)^{\frac{1}{2}}\mathbf{j}$

$$\mathbf{v} = 12\mathbf{i} - 6\sqrt{2}\mathbf{j}$$

$$\text{speed} = \sqrt{12^2 + (-6\sqrt{2})^2} = 6\sqrt{6} = 15 \text{ ms}^{-1} \text{ (2sf)}$$

b) $\mathbf{a} = \frac{d\mathbf{v}}{dt} = 6t\mathbf{i} - 3t^{-\frac{1}{2}}\mathbf{j}$

c) $\mathbf{r} = \int \mathbf{v} dt = t^3\mathbf{i} - 4t^{\frac{3}{2}}\mathbf{j} + \mathbf{c}$

sub in $t=4$, $\mathbf{r} = \mathbf{i} - 4\mathbf{j}$

$$\mathbf{i} - 4\mathbf{j} = (4)^3\mathbf{i} - 4(4)^{\frac{3}{2}}\mathbf{j} + \mathbf{c}$$

$$\mathbf{i} - 4\mathbf{j} = 64\mathbf{i} - 32\mathbf{j} + \mathbf{c}$$

$$\mathbf{c} = -63\mathbf{i} + 28\mathbf{j}$$

$$\therefore \mathbf{r} = (t^3 - 63)\mathbf{i} + (-4t^{\frac{3}{2}} + 28)\mathbf{j}$$

sub in $t=1$

$$\mathbf{r} = -62\mathbf{i} + 24\mathbf{j}$$



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 8 marks)

2.




Figure 1

A rough plane is inclined to the horizontal at an angle α , where $\tan \alpha = \frac{3}{4}$

A small block B of mass 5 kg is held in equilibrium on the plane by a horizontal force of magnitude $X \text{ newtons}$, as shown in Figure 1.

The force acts in a vertical plane which contains a line of greatest slope of the inclined plane.

The block B is modelled as a particle.

The magnitude of the normal reaction of the plane on B is 68.6 N .

Using the model,

(a) (i) find the magnitude of the frictional force acting on B , (3)


(ii) state the direction of the frictional force acting on B . (1)

The horizontal force of magnitude $X \text{ newtons}$ is now removed and B moves down the plane.

Given that the coefficient of friction between B and the plane is 0.5


(b) find the acceleration of B down the plane. (6)

$$\tan \alpha = 3/4$$




$$\sin \alpha = 3/5$$

$$\cos \alpha = 4/5$$



Question 2 continued



"RC" means resolve

a)

(i) R(\nwarrow): $68.6 = X \sin \alpha + 5g \cos \alpha$ ①

$$\Rightarrow X = \frac{68.6 - 5g \cos \alpha}{\sin \alpha} = 49 \text{ N} \quad \text{①}$$

R(\nearrow): $X \cos \alpha = 5g \sin \alpha + F$


$$\Rightarrow F = X \cos \alpha - 5g \sin \alpha = 9.8 \text{ N} \quad \text{①}$$

(ii) Down the plane ①

Friction opposes motion; without friction the box would slide up the plane, so friction must act down to counteract this.

b) $\mu = 0.5$

$$F = \mu R \\ = 0.5R \quad \text{①}$$



- R changes as X is removed
- friction now acts up the plane

R(\nwarrow): $R = 5g \cos \alpha = 39.2$ ① $\therefore a = \frac{5g \sin \alpha - \mu R}{5}$

R(\searrow): $5g \sin \alpha - \mu R = 5a$ ①

$$a = 1.96 \text{ m s}^{-2} \text{ (3sf)} \quad \text{①}$$



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 2 is 10 marks)

3.

[In this question, \mathbf{i} and \mathbf{j} are horizontal unit vectors.]

A particle P of mass 4 kg is at rest at the point A on a smooth horizontal plane.

At time $t = 0$, two forces, $\mathbf{F}_1 = (4\mathbf{i} - \mathbf{j})\text{ N}$ and $\mathbf{F}_2 = (\lambda\mathbf{i} + \mu\mathbf{j})\text{ N}$, where λ and μ are constants, are applied to P

Given that P moves in the direction of the vector $(3\mathbf{i} + \mathbf{j})$

(a) show that

$$\lambda - 3\mu + 7 = 0 \quad (4)$$

At time $t = 4$ seconds, P passes through the point B .

Given that $\lambda = 2$

(b) find the length of AB .

(5)

$$a) \quad \mathbf{F}_1 + \mathbf{F}_2 = k \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} 4 \\ -1 \end{pmatrix} + \begin{pmatrix} \lambda \\ \mu \end{pmatrix} = \begin{pmatrix} 3k \\ k \end{pmatrix} \quad (1)$$

$$4 + \lambda = 3k \quad (1)$$

$$-1 + \mu = k \quad (2)$$

$$\begin{aligned} \text{sub (2) into (1): } 4 + \lambda &= 3(-1 + \mu) \quad (1) \\ 4 + \lambda &= -3 + 3\mu \quad (1) \\ \Rightarrow \lambda - 3\mu + 7 &= 0 \quad (1) \end{aligned}$$

b) given $\lambda = 2$, so find μ :

$$\begin{aligned} 2 - 3\mu + 7 &= 0 \\ \mu &= 3 \end{aligned}$$

\therefore resultant force is

$$\begin{pmatrix} 4 \\ -1 \end{pmatrix} + \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 6 \\ 2 \end{pmatrix} \quad (1)$$



Question 3 continued

$$\underline{F} = m \underline{a}$$

$$\begin{pmatrix} 6 \\ 2 \end{pmatrix} = 4 \underline{a} \quad (1)$$

$$\underline{a} = \begin{pmatrix} 1.5 \\ 0.5 \end{pmatrix}$$

motion AB:

$$\underline{s} = \underline{s}$$

$$\underline{u} = 0$$

$$\underline{v} =$$

$$\underline{a} = 1.5 \underline{i} + 0.5 \underline{j}$$

$$t = 4$$

$$\underline{s} = \underline{u}t + \frac{1}{2} \underline{a}t^2$$

$$= 0 + \frac{4^2}{2} \begin{pmatrix} 1.5 \\ 0.5 \end{pmatrix} \quad (1)$$

$$\underline{s} = \begin{pmatrix} 12 \\ 4 \end{pmatrix}$$

$u=0$ as starts from rest

$$\text{distance} = |\underline{s}| = \sqrt{12^2 + 4^2} = 4\sqrt{10} \quad (1)$$



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 is 9 marks)

4.




Figure 2

A uniform rod AB has mass M and length $2a$

A particle of mass $2M$ is attached to the rod at the point C , where $AC = 1.5a$

The rod rests with its end A on rough horizontal ground.

The rod is held in equilibrium at an angle θ to the ground by a light string that is attached to the end B of the rod.

The string is perpendicular to the rod, as shown in Figure 2.

- (a) Explain why the frictional force acting on the rod at A acts horizontally to the right on the diagram.

(1)

The tension in the string is T

- (b) Show that $T = 2Mg \cos \theta$

(3)

Given that $\cos \theta = \frac{3}{5}$

- (c) show that the magnitude of the vertical force exerted by the ground on the rod at A is $\frac{57Mg}{25}$

(3)

The coefficient of friction between the rod and the ground is μ

Given that the rod is in limiting equilibrium,

- (d) show that $\mu = \frac{8}{19}$

(4)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued



a) The only other force that has a horizontal component is T , which acts to the left. For the rod to be in equilibrium, friction must act to the right. ①

b) $M(A): Mg \times a \cos \theta + 2Mg \times \frac{3a}{2} \cos \theta = T \times 2a$ ①

①

$$4aMg \cos \theta = 2aT$$

$$T = \frac{4aMg \cos \theta}{2a} = 2Mg \cos \theta$$
 ①

c) $\cos \theta = \frac{3}{5} \quad \therefore T = 2Mg \left(\frac{3}{5} \right) = \frac{6Mg}{5}$

①

$$R(\uparrow): R + T \cos \theta = Mg + 2Mg$$
 ①


$$R = 3Mg - \frac{6}{5}Mg \left(\frac{3}{5} \right)$$

$$R = \frac{57Mg}{25}$$
 ①



Question 4 continued

d)



Limiting equilibrium
 $\therefore F = \mu R$ ①

$$R(\rightarrow): \mu R = T \sin \theta \quad \text{①}$$

$$\text{sub in } R = \frac{57Mg}{25}, \quad T = \frac{6Mg}{5}, \quad \sin \theta = \frac{4}{5}$$

$$\mu = \frac{T \sin \theta}{R} = \frac{\frac{6}{5} Mg \times \frac{4}{5}}{\frac{57Mg}{25}} = \frac{8}{19} \quad \text{①}$$



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 11 marks)

Diagram illustrating the projectile motion of a stone from point A to point B. The horizontal distance between A and B is 120 m. The stone is launched from A at an angle α to the horizontal with an initial velocity u .

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

a)



Motion AB: Method: eliminate t to form an equation in terms of only U and α .

Horizontal ($\rightarrow +$)

$$s = 120 \quad \textcircled{1} \quad s = ut + \frac{1}{2}at^2$$

$$u = U \cos \alpha$$

$$v = \quad \quad \quad 120 = Ut \cos \alpha \quad \textcircled{1}$$

$$a = 0$$

$$t = t \quad \quad \quad \Rightarrow t = \frac{120}{U \cos \alpha}$$

Vertical ($\uparrow +$)

$$s = 0 \quad \textcircled{1} \quad s = ut + \frac{1}{2}at^2$$

$$u = U \sin \alpha$$

$$v = \quad \quad \quad 0 = \frac{120 U \sin \alpha}{U \cos \alpha} - \frac{9}{2} \left(\frac{120}{U \cos \alpha} \right)^2 \quad \textcircled{1}$$

$$a = -g$$

$$t = \frac{120}{U \cos \alpha} \quad \textcircled{1} \quad \quad \quad 0 = \frac{120 \sin \alpha}{\cos \alpha} - 4.9 \left(\frac{14,400}{U^2 \cos^2 \alpha} \right)$$



$$0 = 120U^2 \sin \alpha \cos \alpha - 70560$$

$$120U^2 \sin \alpha \cos \alpha = 70560$$

$$U^2 \sin \alpha \cos \alpha = 588 \quad \textcircled{1}$$

Question 5 continued

b)



Motion AC:

Vertical: ($\uparrow +$) ①

$$s = 10$$

$$u = U \sin \alpha$$

$$v = 0$$

$$a = -g$$

$$t =$$

$$v^2 = u^2 + 2as$$

$$0 = U^2 \sin^2 \alpha - 20g \quad ①$$


$$U^2 \sin^2 \alpha = 196 \quad ①$$

$$\text{from (a), } U^2 \sin \alpha \cos \alpha = 588 \quad ②$$

$$① \div ②: \frac{U^2 \sin^2 \alpha}{U^2 \sin \alpha \cos \alpha} = \frac{196}{588} \quad ①$$

$$\tan \alpha = \frac{1}{3}$$

$$\sqrt{3^2 + 1^2} = \sqrt{10}$$



$$\therefore \sin \alpha = \frac{1}{\sqrt{10}} \Rightarrow \sin^2 \alpha = \frac{1}{10}$$

$$①: U^2 = \frac{196}{1/10} = 1960 \quad ①$$

we don't need to worry about the sign of $\sin \alpha$; α is acute so $\sin \alpha$, $\cos \alpha$, and $\tan \alpha$ are all positive.

c) V must be greater as air resistance has to be overcome ①

d) consider the dimensions of the ball. ①



DO NOT WRITE IN THIS AREA



Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 is 12 marks)**TOTAL FOR MECHANICS IS 50 MARKS**