An Effective Medium Field Theory Framework: Elastic Mode—$SU(3)$ Algebraic Correspondence, Soft-Wall Confinement, and Retarded Negative Running

QiDong Ran
Independent Researcher
ORCID: 0009-0004-1570-1915

Abstract

Background: While Quantum Chromodynamics (QCD) excels in the perturbative regime, non-perturbative color confinement lacks an intuitive continuous medium analogy. **Methods:** We construct an Effective Medium Field Theory (EMFT) framework. An explicit algebraic mapping from elastic modes to $SU(3)$ generators is established, and the one-loop β-function is recalculated. Thermodynamic linear confinement is constructed via the work done by an external soft-wall constraint. Dynamical cavitation is derived from the finite response retardation of the background. **Results:** The one-loop recalculation yields $b_0 = 9$ and infers $\Lambda_{\text{QCD}} \approx 246$ MeV. The soft-wall mechanism reproduces a linear confining potential $V = \sigma d$. Discrete Metropolis sampling and 1D PDE simulations verify the Haar limit and the dynamic cavitation effect, where the weak-response time fraction peaks at 86% near $\tau/T \sim 1$. **Conclusions:** This framework does not claim to be the ultimate microscopic theory of QCD, but rather provides a rigorously validated, fully open-source effective analogy framework for understanding strong interactions. **Keywords**: Effective Medium Field Theory, $SU(3)$ Algebraic Correspondence, One-Loop Recalculation, Color Confinement, Retarded Response, Dynamic Cavitation

1 Introduction

While Quantum Chromodynamics (QCD) has achieved tremendous success in the perturbative high-energy regime, the non-perturbative low-energy color confinement phenomenon still lacks an intuitive dynamical picture. The MIT Bag Model introduced vacuum pressure [3], and the flux tube model provided a linear potential [2]. However, an effective medium framework that simultaneously accommodates the algebraic structure, microscopic renormalization group running [4, 5], thermodynamic confinement, and dynamical cavitation is still missing.

In this paper, we propose an Effective Medium Field Theory (EMFT) framework. We make no overclaims of "first principles", but rather construct a quantifiable analogy model based on two macroscopic features of the medium: **volume pressure effects** and **finite response retardation**.

2 Methods

2.1 Algebraic Mapping to $SU(3)$

The traceless part of the 3D elastic medium deformation tensor M_{ij} contains 3 rotations and 5 shears. Mathematically, $\mathfrak{su}(3)$ is composed of anti-Hermitian traceless matrices iT^a; however, physics literature conventionally adopts Hermitian generators $T^a = \lambda^a/2$. We select 8 real basis

matrices e_k and map them as: $T^1 = \frac{1}{2}e_6$, $T^3 = \frac{1}{2}e_4$, $T^4 = \frac{1}{2}e_7$, $T^6 = \frac{1}{2}e_8$, $T^8 = \frac{1}{2}e_5$ and complex combinations: $T^2 = -\frac{i}{2}e_1$, $T^5 = -\frac{i}{2}e_2$, $T^7 = -\frac{i}{2}e_3$. These combinations strictly satisfy the commutation relations $[T^a, T^b] = if^{abc}T^c$. Furthermore, we embed three $SU(2)$ subgroups into 2D subspaces, yielding 9 generators that satisfy the redundancy relation $T^3_{(0,1)} - T^3_{(0,2)} + T^3_{(1,2)} = 0$, spanning a dimension of 8 to fully cover $\mathfrak{su}(3)$.

2.2 One-Loop β-Function Recalculation

Under the background field Feynman gauge ($\xi = 1$), the one-loop effective action is given by the functional determinant. Adopting $\beta(g) = -\frac{b_0}{(4\pi)^2}g^3 + \ldots$, the contributions are: Gluon (adjoint) $+12$, Ghost (adjoint) -1, Quark (fundamental, per flavor) $-2/3$. Summing these gives $b_0(n_f) = 12 - 1 - \frac{2}{3}n_f$. For $n_f = 3$, we obtain $b_0 = 9$. Substituting this into the RG running equation with $\alpha_s(M_Z) = 0.118$, we infer $\Lambda_{\text{QCD}} \approx 245.8$ MeV.

2.3 Soft-Wall Confinement

The confinement potential arises from the mechanical constraint of an external soft wall. The soft wall fixes the cross-sectional area to S. When the quark separation increases by dd, the constraint system displaces the background medium, overcoming the pressure P_{bg} to do work $dW = P_{\text{bg}}Sdd$. Thus, the constraint energy grows linearly as $V_{\text{confine}}(d) = P_{\text{bg}}Sd \equiv \sigma d$.

2.4 Retarded Response and Cavitation

Let the real part of the medium response function be $\text{Re}\,\chi(\omega) = \chi_0/(1 + \omega^2\tau^2)$. The inverse effective coupling correction is proportional to $1/\text{Re}\,\chi(\mu)$, yielding $\Pi(\mu) \simeq c\mu^2\tau^2$. Substituting into $1/g^2_{\text{eff}} = 1/g_0^2 + \Pi$, we obtain $g_{\text{eff}}(\mu) = g_0/\sqrt{1 + cg_0^2\mu^2\tau^2}$. Calculating $\beta(\mu) = \mu\partial_\mu g_{\text{eff}}$ gives $\beta(\mu) \approx -1/(\sqrt{c}\mu\tau) < 0$ in the formal limit $\mu\tau \gg 1$.

3 Results

3.1 Metropolis Monte Carlo Verification

We perform Metropolis sampling on a minimal two-link plaquette toy model. The analytic Haar limit is $P_0 = 8/9 \approx 0.8889$. Fixing $P_0 = 8/9$, we estimate the leading-order strong coupling correction using the data point at $\beta = 0.3$ as $c_1 \approx -0.22$. Parameters: thermalization $n_{\text{burn}} = 3000$, sampling $n_{\text{samp}} = 6000$, skip $n_{\text{skip}} = 2$, binning $n_{\text{bin}} = 30$. Results are shown in Table 1.

Table 1: Discrete Metropolis Monte Carlo Sampling Results

β	P (mean)	error	Accept Rate
0.3	0.822	$\pm$0.014	0.97
1.0	0.621	$\pm$0.011	0.89
4.0	0.233	$\pm$0.005	0.65
14.0	0.070	$\pm$0.002	0.45

3.2 1D PDE Confinement Simulation

Using grid size $N = 301$, spatial step $\Delta x = 0.02a$, and time step $\Delta t = 0.002t_0$, we simulate the defect interaction. Without external pressure, the force approaches zero for $d > 2.0$. With the external soft wall, the force stabilizes at a plateau $F = -6.001 \pm 0.002\, E_0/a$. This is consistent with the theoretical prediction $\sigma = P_{\text{bg}}S = 6.0\, E_0/a$ within 0.02%.

3.3 Retarded Cavitation Dynamics

We define $r(t) = B_{\text{quark}}(t)/B_0 < 0.5$ as the weak-response state. Scanning $\tau/T \in [0.01, 10]$, we find that at $\tau/T \sim 1$, the weak-response time fraction peaks at 86% (based on $N_{\text{samp}} = 10$ independent periods, statistical error $\pm 2\%$).

4 Discussion

Our results demonstrate that an effective medium with finite retardation can reproduce several qualitative features of QCD. The algebraic mapping provides a geometrical interpretation of the $SU(3)$ generators, while the soft-wall mechanism offers a thermodynamic analogy for linear confinement. The dynamic cavitation in the retarded response provides a negative power-law effective running, analogous to asymptotic freedom. It should be noted that the power-law running $\beta \sim -1/(\mu\tau)$ is a phenomenological feature of the medium EFT and does not supersede the standard QCD $-bg^3$ running at high energies. The framework is intended as a cross-disciplinary analogy rather than a fundamental replacement.

5 Conclusions

We have constructed an Effective Medium Field Theory framework. The one-loop recalculation yields $b_0 = 9$ and $\Lambda_{\text{QCD}} \approx 246$ MeV. The soft-wall mechanism constructs a linear confinement potential with constant string tension, and the retarded response induces dynamic cavitation. The model provides a rigorously validated, fully open-source effective analogy framework for understanding strong interactions.

A One-Loop Functional Determinant Integral Combination Table

Operator forms (background field Feynman gauge $\xi = 1$): $\Delta^{\mu\nu}_{\text{gluon}} = -D^2 g^{\mu\nu} + 2igF^{\mu\nu}$, $\Delta_{\text{ghost}} = -D^2$, $\Delta_{\text{quark}} = i\gamma^\mu D_\mu$. Extracting the $1/\epsilon$ poles in $d = 4 - \epsilon$ dimensions:

Table 2: One-loop integral combination table

Field	Rep.	Integrand	$\int_0^1 dx$	$\times$ Casimir/Stat.	Contribution
Gluon (trans.)	Adj.	$8x(1-x)$	$4/3$	$\times C_A = 3$	$+4$
Gluon (spin)	Adj.	$8/3$	$8/3$	$\times C_A = 3$	$+8$
Ghost	Adj.	$2x(1-x)$	$1/3$	$\times C_A = 3$, Fermi minus	-1
Quark	Fund.	$8T_F x(1-x)$	$4T_F/3$	$\times(-1)$, $T_F = 1/2$	$-2/3$

Declarations

Conflict of Interest

The author declares that there is no conflict of interest regarding the publication of this paper.

Ethics Statement

This research did not involve human participants, human data, human tissue, or animals. Therefore, no ethical approval was required.

Funding Statement

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. The author is an independent researcher.

Data Availability Statement

All code in this study is open-source and archived on Zenodo. `su3_full_pipeline.py` and `standalone_measurements.py` are independent recalculation scripts based solely on the Python Standard Library that can directly generate all numerical data and figures in the paper. The fixed version DOI is 10.5281/zenodo.24133353.

References

[1] M. Gell-Mann, "Symmetries of Baryons and Mesons," *Phys. Rev.* **125**, 1067 (1962).

[2] K. G. Wilson, "Confinement of Quarks," *Phys. Rev. D* **10**, 2445 (1974).

[3] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, and V. F. Weisskopf, "A New Extended Model of Hadrons," *Phys. Rev. D* **9**, 3471 (1974).

[4] D. J. Gross and F. Wilczek, "Ultraviolet Behavior of Non-Abelian Gauge Theories," *Phys. Rev. Lett.* **30**, 1343 (1973).

[5] H. D. Politzer, "Reliable Perturbative Results for Strong Interactions?," *Phys. Rev. Lett.* **30**, 1346 (1973).

[6] L. F. Abbott, "The Background Field Method Beyond One Loop," *Nucl. Phys. B* **185**, 189 (1981).

[7] J. J. Sakurai, *Currents and Mesons* (University of Chicago Press, 1969).

[8] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Westview Press, 1995).

[9] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller, "Equation of State Calculations by Fast Computing Machines," *J. Chem. Phys.* **21**, 1087 (1953).

[10] G. S. Bali, K. Schilling, and C. Schlichter, "Static quark-antiquark potential in QCD," *Phys. Rev. D* **51**, 5165 (1995).