

GATE-AII BRANCHES Engineering Mathematics



NUMERICAL METHODS

Lecture No.- 02



By- Rahul sir

Recap of previous lecture



Topic

Numerical methods

Topic

Newton Raphson method

Topic

Problems based on newton Raphson method and bisection rule

Topics to be covered



Topic

Numerical integration

Topic

Simpsons 1/3 rule

Topic

Simpsons 3/8 rule

Topic

Trapezoidal rule

Topic

Problems based on numerical integration

Numerical integration : $\int_a^b f(x) dx$


Area = sum of all rectangles
(Limit of sum)

$$= \underbrace{x_0 f(x_0)}_{\text{Rectangle Area}} + x_1 f(x_1) + \dots + x_n f(x_n)$$

x_0 = Initial guess

x_n = final value

$$\int_{x_0}^{x_n} f(x) dx = \text{Numerical integration}$$



$$\int_a^b f(x) dx = F(b) - F(a)$$

Newton's method

$$x_0 = x_0$$

$$x_1 = x_0 + h$$

$$x_2 = x_0 + 2h$$

$$x_3 = x_0 + 3h$$

⋮

$$\boxed{x_n = x_0 + nh}$$

final value - Initial
value

No. of Intervals.

$h = \text{step size}$

$$\int_{x_0}^{x_0 + nh} f(x) dx = \text{Numerical integral}$$

Calculation for $f(x_i)$ $i = 0, 1, 2, 3$ -
(more error)
 $y_i = f(x_i)$ → Trapezoidal method
→ Simpson's $\frac{1}{3}$ ✓
→ $\frac{3}{8}$ th Rule ✓

Trapezoidal Rule:

Total AREA

$$A_1 + A_2 + A_3 + A_4 + A_5$$

$$A_1 = \frac{h}{2} [f(x_0) + f(x_1)] \quad A_2 = \frac{h}{2} [f(x_1) + f(x_2)]$$


$$A_3 = \frac{h}{2} [f(x_2) + f(x_3)] \quad A_4 = \frac{h}{2} [f(x_3) + f(x_4)]$$

$$A_5 = \frac{h}{2} [f(x_4) + f(x_5)]$$

Total Area

$$\int_{x_0}^{x_5} f(x) dx = \frac{h}{2} [f(x_0) + f(x_5) + 2[f(x_1) + f(x_2) + \dots + f(x_4)]]$$

$$\begin{aligned} \text{AREA} &= \frac{1}{2} (b_1 + b_2) d \\ &= \frac{1}{2} (b_1 + b_2) h \end{aligned}$$



$$\int_{x_0}^{x_5} f(x) dx = \frac{h}{2} \left[(f(x_0) + f(x_5)) + 2(f(x_1) + f(x_2) + f(x_3) + f(x_4)) \right]$$

$$\int_{x_0}^{x_5} f(x) dx = \frac{h}{2} \left[(y_0 + y_5) + 2(y_1 + y_2 + y_3 + y_4) \right]$$

For n Trapezoidal Rule.

$$\int_{x_0}^{x_0+nh} f(x) dx = \frac{h}{2} \left[(y_0 + y_n) + 2(y_1 + y_2 + y_3 + \dots + y_{n-1}) \right] \quad \underline{\underline{\text{Rule.}}}$$

$x_0 = 0$ ✓ possible case

Trapezoidal Rule

#

Simpson $\frac{1}{3}$ rd Rule: any Polynomial ✓



$$\int_a^b f(x) dx = \frac{h}{3} \left[(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2}) \right]$$

(F+L) + 4(odd) + 2(even) ✓ F L O E ✓

Simpson $\frac{3}{8}$ th Rule:

$$h = \frac{\text{Last-final}}{n}$$

#

$$\int_a^b f(x) dx = \frac{3h}{8} \left[(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2}) \right]$$

(F+L) + 4(odd) + 2(even)



Topic : Numerical Methods



$$\int_0^1 e^x dx = e - e^0 = \underline{e - 1}$$

#Q. The value of $\int_0^1 e^x dx$ using the **trapezoidal rule** with four equal subintervals is

$\int_0^1 e^x dx \longrightarrow$ Numerically compute

For equal sub.

A

1.718

B

2.192

C

1.727

D

2.718

$$\int_0^1 e^x dx$$

c) corresponding value of y
 $y = f(x) = e^x$

$$f(x_0) = y_0 = e^{x_0} = e^0 = 1$$

$$f(x_1) = y_1 = e^{x_1} = e^{\frac{1}{4}} = e^{\frac{1}{4}}$$

$$f(x_2) = y_2 = e^{x_2} = e^{\frac{1}{2}} = e^{\frac{1}{2}}$$

$$f(x_3) = y_3 = e^{x_3} = e^{\frac{3}{4}} = e^{\frac{3}{4}}$$

$$f(x_4) = y_4 = e^{x_4} = e^1 = e$$

Step ①

$$n = 4$$

A) Calculate h

$$h = \frac{1-0}{4} = \frac{1}{4} = 0.25$$

(B) Calculate Iteration of x


$$x_0 = 0$$

$$x_1 = x_0 + h = 0 + \frac{1}{4} = \frac{1}{4}$$

$$x_2 = x_1 + h = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

$$x_3 = \frac{1}{2} + \frac{1}{4} = \frac{3}{4}$$

$$x_4 = \frac{3}{4} + \frac{1}{4} = 1$$



Using Trapezoidal Rule

$$\int_0^1 e^x dx = \frac{h}{2} \left[(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1}) \right]$$

$$= \frac{0.25}{2} \left[(e^0 + e^1) + 2(e^{0.25} + e^{0.50} + e^{0.75}) \right]$$

$$\Rightarrow \underline{1.727}$$

$$\text{Absolute Error} = |W - T|$$

①

$$= \underline{8.918 \times 10^{-3}}$$

$$\int_0^1 e^x dx$$

$$= e - 1$$

$$= \underline{\underline{1}}$$



Topic : Numerical Methods



#Q. Value of $\int_4^{5.2} \ln x \, dx$ using Simpson's one-third rule with interval size 0.3 is

using Simpson

$$= \frac{0.3}{3} \left[(\ln 4) + \ln(5.2) + 4(\ln 4.3 + \ln 4.9) + 2[\ln(4.6)] \right]$$

$$= 1.83$$

A 1.83

C 1.51

B 1.60

D 1.06

$x_0 + nh$ $h = 0.3$ $\xrightarrow{\text{final}}$
 x_0 Initial



$$x_0 = 4 \quad x_1 = 4.3 \quad x_2 = 4.6$$

$$x_3 = 4.9 \quad x_4 = 5.2$$

$$y_0 = \ln 4 \quad y_2 = \ln 4.6$$

$$y_1 = \ln 4.3 \quad y_3 = \ln 4.9$$

$$y_4 = \ln(5.2)$$



Topic : Numerical Methods



#Q. The evaluation of the definite integral $\int_{-1}^{1.4} x|x|dx$ by using Simpson's $1/3^{\text{rd}}$ (one-third) rule with step size $h = 0.6$ yields.

✓ N.W

A

0.914

B

1.248

C

0.581

D

0.592



Topic : Numerical Methods



$$h = \frac{(4 - 2)}{2} = 1$$

#Q. Evaluation of $\int_2^4 x^3 dx$ using a 2-equal- segment trapezoidal rule gives a value of _____.

$$\int_2^4 x^3 dx$$

$$y_0 = (2)^3 = 8$$

$$y_1 = (3)^3 = 27$$

$$y_2 = (4)^3 = 64$$

$$\begin{array}{c} x_0 = 2 \\ | \\ \hline | \\ \hline 3 = x_1 \\ | \\ \hline | \\ \hline x_2 = 4 \end{array}$$

$$\begin{aligned} \underline{\text{Error}} &= \frac{1}{2} [(8 + 64) + 2(27)] \\ &= \underline{63} \end{aligned}$$



Topic : Numerical Methods



#Q. Numerical integration using trapezoidal rule gives the best results for a single variable function, which is

A

Linear

B

Parabolic

C

Logarithmic

D

Hyperbolic



Topic : Numerical Methods



$$h = \frac{\pi - 0}{3} = \frac{\pi}{3}$$


#Q. The error in numerically computing the integral $\int_0^{\pi} (\sin x + \cos x) dx$ using the trapezoidal rule with three intervals of equal length between 0 and π is _____.

$$x_0 = 0 \quad y_0 = \sin 0 + \cos 0$$

$$x_1 = \frac{\pi}{3} \quad y_1 = \sin \frac{\pi}{3} + \cos \frac{\pi}{3}$$

$$x_2 = \frac{2\pi}{3} \quad y_2 = \sin \left(\frac{2\pi}{3} \right) + \cos \left(\frac{2\pi}{3} \right)$$

$$x_3 = \pi \quad y_3 = \sin \pi + \cos \pi$$



$$I = \int_0^{\pi} (\sin x + \cos x) dx = 2$$

$$\text{Trapezoidal} = \frac{h}{2} \left[(y_0 + y_3) + 2(y_1 + y_2) \right] = \underline{1.8137}$$

$$\text{Error} = |2 - 1.8137| = \underline{0.1863}$$



Topic : Numerical Methods



$$\checkmark \int_0^1$$

#Q. Simpson's $\frac{1}{3}$ rule is used to integrate the function $f(x) = \frac{3}{5}x^2 + \frac{9}{5}$ between $x = 0$ and $x = 1$ using the least number of equal sub-intervals. The values of the integral is _____.

\checkmark H.W



Topic : Numerical Methods



✓ h.w

#Q. The value of function $f(x)$ at 5 discrete points are given below :

X	0	0.1	0.2	0.3	0.4
f(x)	0	10	40	90	160

Using Trapezoidal rule with step size of 0.1, the value of $\int_0^{0.4} f(x)dx$ is ____.



Topic : Numerical Methods

#Q. Using a unit step size, the value of integral $\int_1^2 x \ln x dx$ by trapezoidal rule is_____

(H.W)



Topic : Numerical Methods



#Q. The definite integral $\int_1^3 \frac{1}{x} dx$ is evaluated using Trapezoidal rule with a step size of 1. The correct answer is _____.

H.W



Topic : Numerical Methods



#Q. The value of $\int_{2.5}^4 \ln x dx$ calculate using the Trapezoidal rule with five subintervals is _____.

✓ H.W



Topic : Numerical Methods



#Q. Using the trapezoidal rule, and dividing the interval of integration into three equal subintervals, the definite integral $\int_{-1}^{+1} |x| dx$ is___.

✓ H.W



Topic : Numerical Methods



#Q. The estimated of $\int_{0.5}^{1.5} \frac{dx}{x}$ Obtained using Simpson's rule with three-point function evaluation exceeds the exact value by

H.W

A

0.235

B

0.068

C

0.024

D

0.012



Topic : Numerical Methods



#Q. The integral $\int_1^3 \frac{1}{x} dx$, when evaluated by using Simpson's 1/3 rule on two equal subintervals each of length 1, equals

H.W

A

1.000

B

1.098

C

1.111

D

1.120



2 mins Summary



Topic

One

Trapezoidal

Topic

Two

Simpson $\frac{1}{3}$ rd

Topic

Three

Simpson $\frac{3}{8}$ th

Topic

Four

Topic

Five

THANK - YOU

Topics to be Covered