

GATE (ALL BRANCHES)

Engineering Mathematics

Differential Equation +
Partial differential

Lecture No. 03

By- Rahul Sir



TOPICS TO BE COVERED

o1

Problems based on Homogeneous
Differential Equation

o2

Non Homogeneous Differential Equation

o3

Questions based on Non Homogeneous
Differential Equation

o4

Linear Differential Equation

Imp ✓

Reducible to Homogenous form:


$$f(x, y) = \frac{ax + by + c}{a'x + b'y + c'}$$

$$\frac{dy}{dx} = \frac{ax + by + c}{a'x + b'y + c'}$$

Non Homogenous

Check whether this function Homogenous or Not

$$f(kx, ky) = \frac{a \cdot kx + b \cdot ky + c}{a'kx + b'ky + c'}$$



Remove The Non-Homogenous part
Convert to The Homogenous Eqⁿ

$$\frac{dy}{dx} = \frac{ax+by+c}{a'x+b'y+c'}$$

Put $\begin{cases} x = x+h \\ y = y+k \end{cases} \quad \frac{dy}{dx} = \frac{dy}{dx}$

$$\begin{aligned} x &= x+h \\ y &= y+k \\ dx &= dx \\ dy &= dy \end{aligned}$$

where h, k are constants
and x, y are functions
of x, y

$$\frac{dy}{dx} = \frac{a(x+h)+b(y+k)+c}{a'(x+h)+b'(y+k)+c'} = \frac{ax+by+(ah+bk+c)}{a'x+b'y+(a'h+b'k+c')}$$

$\begin{aligned} ah+bk+c &= 0 \\ a'h+b'k+c' &= 0 \end{aligned}$ both are
straight line eqⁿ

Solve the value of h, k



Homogenous

$$\frac{dy}{dx} = \frac{ax+by+0}{a'x+b'y+0} = \frac{ax+by}{a'x+b'y}$$

$$\frac{dy}{dx} = \frac{ax+by}{a'x+b'y}$$

If Eqnⁿ is Homogenous Then
Put $y = vx$

$$v + x \frac{dv}{dx} = \frac{ax+bvx}{a'x+b'vx} = \frac{(a+bv)}{(a'+b'v)}$$

Now Separate The variable and get The
Solution of D.E.

✓ Linear D.E

✓ Reduced to linear
D.E

✓ Variable separable.

✓ Reduced to V.S

✓ Homogenous

✓ Reduced to Homo

Linear Differential Equation:

Dep. variable $y = f(x)$ Independent variable

$$\frac{dy}{dx} \cdot y = \text{Non Linear}$$

$\frac{dy}{dx} \text{ (D)} = \checkmark$

$x \frac{dx}{dy} = \text{Non-Linear}$

Dependent derivative

$$x \frac{dy}{dx} = \text{Linear}$$

$$y \frac{dx}{dy} = \text{Linear}$$

$$\sin x \frac{dy}{dx} = \text{Linear}$$

$$\sin y \frac{dy}{dx} = \text{Non Linear}$$

$$t^2 \frac{dt}{dx} = \text{Non Linear}$$

(A) degree always $\frac{1}{\left(\frac{dy}{dx}\right)}$ Degree

(B) (Dependent var) \times (Dependent var derivative)

(C) No Non Linear function in dep. variable. is Not Present in the eqn.

$$t^2 \frac{dx}{dt} = \text{Linear}$$

$$\frac{dy}{dx} + y = e^x$$

D.E

Wrong Hit and trial method

Not Integrate

Wrong Hit and trial method

Not solved

D.E

- A) ✓ Var. separable form.
- B) ✓ Reduced var. sep.
- (C) ✓ Homogenous.
- (D) ✓ Reduced to Homogenous.

multiply both sides e^x

$$\frac{dy}{dx} e^x + y e^x = x e^x$$

$$\frac{d}{dx} [y e^x] = \frac{dy}{dx} e^x + y \cdot e^x = \frac{d}{dx} (y e^x)$$

I II

Important Key Point

multiply करें पर D.E solve

$$\Rightarrow \frac{d}{dx}(ye^x) = xe^x$$

both sides Integrate It

$$\Rightarrow \int \frac{d}{dx}(ye^x) = \int xe^x$$

$$\Rightarrow \boxed{ye^x = (x-1)e^x + c} \text{ solution of D.E}$$

D
↓
I
↓
D.E

Hit and trial method

→ factor multiply = ?
↙ D.E solve.

Linear D.E

$$\frac{dy}{dx} + Py = Q \quad \text{Where } P \text{ and } Q \text{ Are Function of } x \text{ only.}$$

Integrating factor = $e^{\int P dx}$ Linear form $ax + by + c = 0$

$$\Rightarrow e^{\int P dx} \cdot \frac{dy}{dx} + Py e^{\int P dx} = Q e^{\int P dx}$$

Solution of this D.E

$$\Rightarrow \frac{d}{dx} (y e^{\int P dx}) = Q e^{\int P dx}$$

$$y \cdot (I.F) = \int (RHS)(I.F) dx + C$$

Soln

$$\frac{dy}{dx} + Py = Q$$

both sides Integrate It

$$\int \frac{d}{dx} (y e^{\int P dx}) = \int Q e^{\int P dx} dx$$

$$\frac{dx}{dy} + Px \Rightarrow Q \quad \text{Where } P \text{ and } Q \text{ Are function of } y \text{ only}$$

Integrating Factor = $e^{\int P dy}$

→ solution

$$x(I.F) = \int (RHS)(I.F) dy + C \quad \text{solution of D.E}$$

Q.

Questions

#Q. The solution of the equation $\frac{dQ}{dt} + Q = 1$ with $Q = 0$ at $t = 0$ is

(a) $Q(t) = e^{-t} - 1$

(b) $Q(t) = 1 + e^{-t}$

(c) $Q(t) = 1 - e^t$

(d) ✓ $Q(t) = 1 - e^{-t}$

$$\frac{dQ}{dt} + Q = 1 \quad \text{with } Q=0 \text{ at } t=0$$

$$M=1 \quad N=1$$

$$I.F = e^{\int 1 dt} = e^t$$

$$Q \cdot e^t = \int e^t \cdot 1 dt$$

$$Qe^t = e^t + c$$

$$0 = e^0 + c \quad (c = -1)$$

$$Qe^t = e^t - 1$$

$$Q = 1 - e^{-t}$$

$\frac{dQ}{dt} + M Q = N$
where M and N
Are function
of t .

Q.

Questions

#Q. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function which satisfies

$$f(x) = \int_0^x f(t) dt, \text{ then the value of } f(\ln 5)$$

$$f'(x) = \frac{dy}{dx}$$

$$y = f(x)$$

Apply Newton Leibnitz Rule

$$f'(x) = f(x) \frac{d}{dx} \cdot 1$$

$$f'(x) = f(x)$$

$$\frac{dy}{dx} = y$$

$$f(x) = \int_0^x f(t) dt$$

$$x=0 \quad f(0) = \int_0^0 f(t) dt = 0$$

$$\int \frac{dy}{y} = \int dx$$

$$= \ln y = x + C$$

$$y = Ae^x$$

A = unknown

$$f(x) = y = Ae^x$$

$$A = 0$$

Solution $y = 0$


$$y(\ln 5) = 0$$

Ans

$$f(0) = 0$$

$$x = 0$$

$$y = 0$$



$$y(\ln 5) = 0 \quad y(\ln 2) = 0$$

$$y(81) = 0 \quad y(9) = 0$$

Q.

Questions

#Q. $\frac{dy}{dx} = \frac{2+(x-y)}{3+2x-2y}$

✓ Solve

✓ n.w

Do yourself

$$\frac{dy}{dx} = \frac{2+(x-y)}{3+2(x-y)}$$

Put

$$x-y = t$$

And solve

The D.E

#Q. If $x^2 \left(\frac{dy}{dx} \right) + 2xy = \frac{2 \ln x}{x}$ and $y(1) = 0$ then what is $y(e)$?

$$x^2 \frac{dy}{dx} + 2xy = \frac{2 \ln x}{x}$$

$$= \frac{dy}{dx} + \frac{2x}{x^2} y = \frac{2 \ln x}{x^3}$$

$$\Rightarrow \frac{dy}{dx} + \frac{2y}{x} = \frac{2 \ln x}{x^3} \quad \text{form}$$

$$P = \frac{2}{x} \quad Q = \frac{2 \ln x}{x^3}$$

$$\text{I.f} = x^2$$

$$\frac{dy}{dx} + Py = Q$$

P and Q function of x

$$\text{I.F} = e^{\int P dx}$$

$$= e^{\int \frac{2}{x} dx}$$

$$= e^{2 \ln x}$$

$$= e^{\ln x^2} = x^2$$

(a) e

(b) 1

(c) $\frac{1}{e}$

(d) $\frac{1}{e^2}$

$$y \cdot x^2 = \int \frac{2 \ln x \cdot x^2}{x^3 x} dx = 2 \int \frac{\ln x}{x} dx$$

Put $\ln x = t$

$$\frac{dx}{x} = dt$$

$$\int \frac{\ln x}{x} = \int t dt$$

$$= \frac{t^2}{2} + C$$

$$= \frac{(\ln x)^2}{2} + C$$

$$y x^2 = \frac{2 (\ln x)^2}{2} + C$$

Apply Initial condition $y(1) = 0$

$$0 = \frac{[\ln(1)]^2}{2} + C \quad \boxed{C = 0}$$

$$y x^2 = \frac{(\ln x)^2}{2} + 0$$

$$y = \frac{(\ln x)^2}{x^2}$$

$$y(e) = \frac{(\ln_e e)^2}{e^2} = \frac{1}{e^2}$$

Q.

Questions

#Q. The solution of the differential equation $\frac{dy}{dx} + 2xy = e^{-x^2}$ with $y(0) = 1$ is

$$P = 2x \quad Q = e^{-x^2} \quad I.F = e^{\int 2x dx} = e^{x^2}$$

$$y e^{x^2} = \int e^{x^2} \cdot e^{-x^2} dx$$

$$y e^{x^2} = \int dx$$

$$y e^{x^2} = x + C$$

$$1e^0 = 0 + C$$

$$C = 1$$

$$y e^{x^2} = x + 1$$

$$y = \frac{(x+1)}{e^{x^2}}$$

$$y = x e^{-x^2} + e^{-x^2} = e^{-x^2}(x+1)$$

(a) $(1+x)e^{x^2}$

(b) $(1+x)e^{-x^2}$

(c) $(1-x)e^{x^2}$

(d) $(1-x)e^{-x^2}$

Q.

Questions

#Q. The solution of $x \frac{dy}{dx} + y = x^4$ with condition $y(1) = \frac{6}{5}$

(a) $y = \frac{x^4}{5} + \frac{1}{x}$

(b) $y = \frac{4x^4}{5} + \frac{4}{5x}$

(c) $y = \frac{x^4}{5} + 1$

(d) $y = \frac{x^5}{5} + 1$

$$x \frac{dy}{dx} + y = x^4$$

$$\frac{dy}{dx} + \frac{y}{x} = \frac{x^4}{x}$$

$$\frac{dy}{dx} + \frac{y}{x} = x^3$$

$$P = \frac{1}{x} \quad Q = x^3$$

Do yourself

Q.

Questions

#Q2. Let $f: [1, \infty] \rightarrow (2, \infty)$ be a differentiable function such that $f(1) = 2$

$$6 \int_1^x f(t) dt = 3x f(x) - x^3$$

= 6

Then the value of $f(2)$, $\forall x \geq 1$

Ans

→ Using Leibnitz Rule

$$6 f(x) \cdot \frac{d}{dx}(x) = 3 [x \cdot f'(x) + f(x) \cdot 1] - 3x^2 \Rightarrow \frac{dy}{dx} - \frac{y}{x} = x$$

$$6f(x) = 3x f'(x) + 3f(x) - 3x^2$$

$$2f(x) = x f'(x) + f(x) - x^2$$

$$y = f(x)$$

$$y' = dy/dx = f'(x)$$

$$x f'(x) + f(x) - 2f(x) = x^2$$

$$x f'(x) - f(x) = x^2$$

$$f'(x) - \frac{f(x)}{x} = \frac{x^2}{x} = x$$

$$P = -\frac{1}{x} \quad Q = x$$

$$I.F = e^{\int -\frac{1}{x} dx} = e^{-\ln x} = \left(\frac{1}{x}\right)$$

#Q3. Let $f(x)$ be a differentiable function on the interval $(0, \infty)$ such that $f(1) = 1$

$$\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1 \text{ for } x > 0$$

$t \rightarrow \text{variable}$
 $x \rightarrow \text{const}$
 $\rightarrow 0$
 $\rightarrow 0$

$$\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1$$

Using L-Hospital Rule.

(a) $\frac{1}{3x} + \frac{2x^2}{3}$

(b) $-\frac{1}{3x} + \frac{4x^2}{3}$

(c) $-\frac{1}{x} + \frac{2}{x^2}$

(d) $\frac{1}{x}$

$$\Rightarrow \lim_{t \rightarrow x} \frac{2t f(x) - x^2 f'(t)}{1 - 0} = 1$$

$$= \lim_{t \rightarrow x} \frac{2t f(x) - x^2 f'(t)}{1} = 1$$

$$= \underline{2x f(x) - x^2 f'(x) = 1}$$

$$2xf(x) - x^2 f'(x) = 1$$

$$\Rightarrow 2xy - x^2 \frac{dy}{dx} = 1$$

$$\Rightarrow -x^2 \frac{dy}{dx} + 2xy = 1$$

$$\Rightarrow x^2 \frac{dy}{dx} - 2xy = -1$$

$$\Rightarrow \frac{dy}{dx} - \frac{2xy}{x^2} = -\frac{1}{x^2}$$

$$\Rightarrow \boxed{\frac{dy}{dx} - \frac{2y}{x} = -\frac{1}{x^2}}$$

Do yourself

$$y = f(x)$$

$$\frac{dy}{dx} = f'(x)$$

$$\frac{dy}{dx} - \frac{2y}{x} = -\frac{1}{x^2}$$

This is linear D.E form

$$P = -\frac{2}{x} \quad Q = -\frac{1}{x^2}$$

$$I.F = e^{\int P dx} = e^{\int -\frac{2}{x} dx}$$

$$= e^{-2 \ln x}$$

$$= e^{-\ln x^2} = \left(\frac{1}{x^2} \right)$$

#Q4. If $y(t)$ is a solution of $(1+t)\frac{dy}{dt} - ty = 1$ and $y(0) = -1$. Then $y(1)$ is equal to

(a) $\frac{1}{2}$

(b) $e + \frac{1}{2}$

(c) $e - \frac{1}{2}$

(d) $-\frac{1}{2}$

$$\frac{dy}{dt} - \frac{t}{(1+t)} y = \frac{1}{(1+t)} \quad P = -\frac{t}{(1+t)}$$

$$Q = \frac{1}{(1+t)}$$

$$\text{I.F} = e^{\int \frac{-t}{(1+t)} dt}$$

$$-\int \frac{t+1-1}{1+t} = -\left[\int \frac{t+1}{t+1} - \int \frac{1}{1+t} \right]$$

$$= -[t - \ln(1+t)] = \ln(1+t) - t$$

$$= e^{\ln(1+t) - t}$$

$$= e^{-t} e^{\ln(1+t)} = \underline{e^{-t}(1+t)}$$

$$I \cdot F = e^{-t}(1+t)$$

Solution of D.E

$$= y \cdot [e^{-t}(1+t)] = \int \cancel{e^{-t}(1+t)} \cdot \frac{1}{\cancel{(1+t)}} dt$$

$$= y e^{-t}(1+t) = \int e^{-t} dt$$

$$= y e^{-t}(1+t) = -e^{-t} + C$$

$$= -1 e^0(1+0) = -e^0 + C$$

$$= -1 = -1 + C$$

$$= -1 + 1 = C$$

$$(C=0)$$

$$\frac{y(0) = -1}{t=0}$$

$$y = -1$$

$$ye^{-t}(1+t) = -e^{-t} + 0$$

$$y = \frac{-e^{-t}}{e^{-t}(1+t)}$$

$$= \frac{-e^{-t} \cdot e^t}{(1+t)}$$

$$y = \frac{-1}{(1+t)}$$

$$y = \frac{-1}{1+1}$$

$$\boxed{y = -\frac{1}{2}} \quad \underline{\underline{Ans}}$$

#Q5. If $y=y(x)$, $\frac{2+\sin x}{(y+1)} \frac{dy}{dx} = -\cos x$
 $y(0)=1$, then $y(\pi/2)$ equals to

- (a) $1/3$
- (b) $2/3$
- (c) $-1/3$
- (d) 1

$$\frac{2+\sin x}{(y+1)} \frac{dy}{dx} = -\cos x$$

$$\frac{dy}{dx} = \frac{-\cos x (y+1)}{(2+\sin x)}$$

$$\frac{dy}{dx} + Py = Q$$

$$\frac{dy}{dx} = \frac{-\cos x \cdot y - \cos x}{2+\sin x}$$

$$\frac{dy}{dx} + y \frac{\cos x}{2+\sin x} = \frac{-\cos x}{2+\sin x}$$

$$P = \frac{\cos x}{2+\sin x} \quad Q = \frac{-\cos x}{(2+\sin x)}$$

$y(0)=1$ and get the solution

Do yourself

Q.

Questions

MSQ

#Q7. If $y(x)$ satisfies the D.E.

$$y' - y \tan x = 2 \sec x, \quad y(0) = 0 \text{ then}$$

(a) $y(\pi/4) = \frac{\pi^2}{8\sqrt{2}}$

(b) $y'(\pi/4) = \frac{\pi^2}{18}$

(c) $y(\pi/4) = \frac{\pi^2}{9}$

(d) $y'(\pi/3) = \frac{4\pi}{3} + \frac{2\pi^2}{3\sqrt{3}}$

Do yourself
✓

$$\frac{dy}{dx} - y \tan x = 2 \sec x$$

$$\begin{aligned} P &= -\tan x \\ Q &= 2 \sec x \end{aligned} \left. \begin{array}{l} \text{I.F} \\ \checkmark \text{Integral value} \\ \checkmark \text{Ans} \end{array} \right\}$$

Q.

Questions

#Q. $\frac{dy}{dx} + y = \frac{1 + \sin x}{1 + \cos x}$

$P = 1$
 $Q = \frac{1 + \sin x}{1 + \cos x}$

Solution
 do yourself

$\frac{dy}{dx} + Py = Q$

Q.

Questions

#Q. $(1+y^2)dx = (\tan^{-1}y - x)dy$

to yourself

Thank You!

PW Soldiers