

Data Science and Artificial Intelligence Probability and Statistics

Bivariate Random Variable

Lecture No.- **04**



By- Rahul Sir

Topics to be Covered



Topic

Problem Based on Bivariate Random Variable
02 (Univariate / Bivariate)



Topic : Problem Based on Bivariate Random Variable



Q1. A fair coin is tossed 3 times. Let the random variable X denote the number of heads in 3 tosses of the coin. Find the probability density function of X.

A. $\left(\frac{3}{x}\right) \left(\frac{1}{2}\right)^{2x} \left(\frac{1}{2}\right)^{2-x}$

B. $\left(\frac{3}{2x}\right) \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{1-x}$

C. $\left(\frac{3}{x}\right) \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{3-x}$

D. $\left(\frac{3}{x}\right) \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{4-x}$



Topic : Problem Based on Bivariate Random Variable

PW

$$\int_1^2 + \int_2^3 + \int_3^4 + \int_4^5 + \int_5^6 + \dots$$

Q2. If the probability of a random variable X is given by

$$\sum x = \frac{x(x+1)}{2}$$

1, - - 12

$$= \underline{12}(12+1)$$

$$= \frac{62}{x} \times 13 = 78$$

$$f(x) = k(2x - 1), \quad x = 1, 2, 3, \dots, 12. \text{ Find } k.$$

$$f(x) = K(2x - 1) \quad x = 1, 2, 3, \dots, 12$$

$$\sum \text{all Probs.} = 1$$

$$\sum_{k=1}^{12} 2kx - k = 1$$

$$2k \sqrt{\sum_{x=1}^{12} x} - k \sqrt{\sum_{k=1}^{12} 1} = 1$$

$$2kx \frac{12 \times 13}{2} - kx 12 = 1$$

$$K = \frac{1}{144}$$

$$\begin{array}{c} 1+1+1+1+1+1+ \\ 1+1+1+ \\ \hline = 12 \end{array}$$



Topic : Problem Based on Bivariate Random Variable



Q3. The density function for the continuous random variable X is

$$f_x(x) = \begin{cases} e^{-x} & \text{for } x > 0 \\ 0 & \text{for } x \leq 0 \end{cases}$$

Find the Probability $P[x \leq 2 | x > 1]$

$$P\left(\frac{A}{B}\right)$$

B is already
Happened
 $\frac{A}{B}$ Happening

$$f(x) = \begin{cases} e^{-x} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

$$P(x \leq 2 | x > 1) = \frac{P(x \leq 2 \wedge x > 1)}{P(x > 1)}$$
$$= \frac{\int_1^2 e^{-x} dx}{\int_1^{\infty} e^{-x} dx}$$

$$\checkmark \text{ Nr } \int_1^2 e^{-x} dx = \left[-e^{-x} \right]_1^2 = -e^{-2} + e^{-1} \\ = e^{-1} - e^{-2}$$

$$\checkmark \text{ Dr } \int_1^\infty e^{-x} dx = \left[-e^{-x} \right]_1^\infty = -e^{-\infty} + e^{-1} \\ = 0 + e^{-1}$$

$$P\left(\frac{X \leq 2 \wedge X > 1}{P(X > 1)}\right) = \frac{e^{-1} - e^{-2}}{e^{-1}}$$
$$\checkmark = \boxed{1 - e^{-1}} = \underline{0.632}$$



Topic : Problem Based on Bivariate Random Variable



Q4. Suppose the random variable X has a probability density function

Conti
Random
var

$$f(x) = \begin{cases} kx^3 e^{-x/2}, & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

$$f(x) = \begin{cases} kx^3 e^{-\frac{x}{2}} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

The value of k is

- A. 1/96
- B. 96
- C. 8/3
- D. 1/4

$$\int_{-\infty}^{\infty} f(x) dx = 1$$
$$= \left[\int_{-\infty}^{\infty} kx^3 e^{-x/2} dx \right] = 1$$

$$= \int_0^\infty K e^{-x^2} 2x^3 dx = 1$$

$$\Rightarrow K \int_0^\infty e^{-\frac{x^2}{2}} x^3 dx = 1$$

$$\begin{aligned} t &= \frac{D}{2} = D & \frac{x}{2} = t & \cdot x = 2t \\ t &= \frac{\infty}{2} = \infty & dx = 2dt \end{aligned}$$

$$\Rightarrow K \int_0^\infty e^{-t} \cdot (2t)^3 \cdot 2 dt = 1$$

$$\Rightarrow 16K \int_0^\infty e^{-t} t^3 dt = 1$$

Convert The Gamma function

$$K = \frac{1}{96} \quad \checkmark$$

$$\boxed{n-1=3} \rightarrow \boxed{n=4}$$

D to ∞ DR - $\omega_0 \rightarrow \infty$

$$\Gamma n = \int_0^\infty e^{-x} x^{n-1} dx$$

~~exponential + algebraic~~

$$16K \left[\int_0^\infty e^{-t} t^3 dt \right] = 1$$

$$\begin{aligned} 16K \times \sqrt{4} &= 1 \\ 16K \times 3! &= 1 \end{aligned}$$



Topic : Problem Based on Bivariate Random Variable



Q5. Let X be a continuous random variable with pdf

$$f_x(x) = \begin{cases} cx^2, & \text{for } 0 < x \leq 1, \\ 0 & \text{otherwise} \end{cases}$$

$$\int_0^1 cx^2 dx = 1$$

$$c = 3$$

$$f(x) = cx^2$$
$$0 < x \leq 1$$
$$= 0 \quad \text{otherwise}$$

$$\text{For some positive constant } c. \text{ The value of } P\left(X \leq \frac{2}{3} \mid X > \frac{1}{3}\right)$$

$$P\left(X \leq \frac{2}{3} \mid X > \frac{1}{3}\right)$$

A. $3/26$

B. $5/26$

C. $7/26$

D. $11/26$

$$P\left(X \leq \frac{2}{3} \mid X > \frac{1}{3}\right) = \frac{P\left(X \leq \frac{2}{3} \wedge X > \frac{1}{3}\right)}{P\left(X > \frac{1}{3}\right)}$$
$$= \frac{\int_{1/3}^{2/3} 3x^2 dx}{\int_{1/3}^1 3x^2 dx}$$
$$= \frac{\left[x^3\right]_{1/3}^{2/3}}{\left[x^3\right]_{1/3}^1}$$
$$= \frac{\left(\frac{8}{27} - \frac{1}{27}\right)}{\left(1 - \frac{1}{27}\right)}$$
$$= \frac{7/27}{26/27}$$
$$= \frac{7}{26}$$



Topic : Problem Based on Bivariate Random Variable



Q6. Suppose the random variable X has the probability density function

$$f(x) = \begin{cases} ce^{x/3}, & x \leq 0, \\ ce^{-x/3}, & x > 0, \end{cases}$$



Do yourself

For some positive constant c. The value of $P [x > 6/x > 0]$ is

- A. e^{-2}
- B. ce^{-2}
- C. 0
- D. $1-e^{-2}$



Topic : Problem Based on Bivariate Random Variable



Q7. Let X be a discrete random variable with probability function $P(X = x) = \frac{2}{3^x}$ for

$x = 1, 2, 3, \dots$. What is the probability that X is even?

- A. $\frac{1}{4}$
- B. $\frac{2}{7}$
- C. $\frac{1}{3}$
- D. $\frac{2}{3}$

$P(X=x)$	2	4	6	8	10
$\frac{2}{3^2}$	$\frac{2}{3^4}$	$\frac{2}{3^6}$	$\frac{2}{3^8}$	$\frac{2}{3^{10}}$	

$$P(X=x) = \frac{2}{3^x} \quad x=1, 2, 3$$

What is The
Probability X is
even

$$= P(X=2) + P(X=4) + P(X=6) + P(X=8) + \dots$$

Discrete Random [countable SET)

$$= \underbrace{\frac{2}{3^2} + \frac{2}{3^4} + \frac{2}{3^6} + \frac{2}{3^8} + \frac{2}{3^{10}} + \dots}_{\text{SERIES}}$$

$$= \frac{2}{9} \left[1 + \underbrace{\frac{1}{9} + \left(\frac{1}{9}\right)^2 + \left(\frac{1}{9}\right)^3 + \left(\frac{1}{9}\right)^4 + \dots}_{\text{G.P.}} \right]$$

$$\Rightarrow \frac{2}{9} \left[1 + \frac{\frac{1}{9}}{1 - \frac{1}{9}} \right]$$

$$= \frac{2}{9} \left[1 + \frac{\frac{1}{9}}{\frac{8}{9}} \right] = \left(\frac{1}{4} \right)$$

$$\begin{aligned} S_{\infty} &= \frac{a}{1-r} = \frac{\text{Ist term}}{1-\text{common ratio}} \\ r &= \frac{T_2}{T_1} = \end{aligned}$$



Topic : Problem Based on Bivariate Random Variable




Q8. Let $f(x) = \frac{k|x|}{(1+|x|)^4}, -\infty < x < \infty$

$$f(x) = \frac{K|x|}{(1+|x|)^4} \quad -\infty < x < \infty$$

Then the value of k for which $f(x)$ is a probability density function is

- A. $1/6$
- B. $1/2$
- C. 3
- D. 6

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{K|x|}{1+|x|^4} dx &= 1 \\ &= 2 \int_0^{\infty} \frac{Kx}{1+x^4} dx = 1 \\ &= 2 \int_0^{\infty} \frac{Kx}{1+x^4} dx \end{aligned}$$



$$= 2 \int_0^\infty \frac{kx}{(1+x)^4} dx = 2 \int_0^\infty \frac{k \cdot x^{l-1}}{(1+x)^{l+m}}$$

bracket

$$= 2 \int_0^\infty \frac{k \cdot x}{(1+x)^4} dx = 2 \int_0^\infty \frac{k \cdot x^{2-1}}{(1+x)^{2+2}}$$

$$= 2 \times k \frac{\Gamma 2 \Gamma 2}{\Gamma 4}$$

$$= \frac{2k \cdot 1}{3!} = \frac{1}{\Gamma 2 + 2}$$

$|x|$ even
Function

$$I = \int_{-\infty}^{\infty} \frac{k|x|}{(1+|x|)^4}$$

(odd).

Beta Function

$$\int_0^\infty \frac{k \cdot x}{(1+x)^4} dx$$

$$\frac{x^{l-1}}{(1+x)^{l+m}}$$

$$= \frac{\Gamma l \Gamma m}{\Gamma l+m}$$

$$k=3$$



Topic : Problem Based on Bivariate Random Variable



Q10. A continuous random variable X has density function

$$f(x) = \begin{cases} 2x & 0 < x < \frac{1}{2} \\ \frac{4-2x}{3} & \frac{1}{2} \leq x < 2 \\ 0, & \text{elsewhere} \end{cases}$$

✓ Do yourself

Find $P[0.25 < x \leq 1.25]$



Topic : Problem Based on Bivariate Random Variable

P
W

(H2B)

Q11. Let X be a continuous random variable with probability density function


$$f(x) = \frac{1}{2} e^{-|x-1|}, -\infty < x < \infty$$

10th $|X| < a, |X| > a$

$$P[1 < |X| < 2] = |x| > 1, |x| < 2$$

OR

Find the value of $P(1 < |X| < 2)$



$$\begin{aligned} & \left\{ \begin{array}{l} (-\infty, -1) \cup (1, \infty) \\ -2 < x < 2 \end{array} \right. \\ & \Rightarrow \int_{-1}^1 \frac{1}{2} e^{-(x-1)} dx + \int_1^2 \frac{1}{2} e^{-(x-1)} dx \\ & \Rightarrow \frac{1}{2} (e^{-1} + e)(e^{-1} + e^{-2}) \end{aligned}$$

$$\int_0^{\infty} \frac{x^{l-1}}{(1+x)^{m+l}} = \frac{\Gamma(l)\Gamma(m)}{\Gamma(l+m)} = \frac{\Gamma(l)l!}{\Gamma(m)m!} = \frac{l!(l-1)!}{(m-1)!}$$



Topic : Problem Based on Bivariate Random Variable



Q12. The following table represents the joint probability distribution of the discrete random variable (X, Y):

		Y	1	2
		X		
X	1	0.1	0.2	
	2	0.1	0.3	
		3	0.2	0.1

Find :

- (i) $F(2, 2)$, $F(3, 2)$
- (ii) $F_x(3)$
- (iii) $F_Y(1)$



Topic : Problem Based on Bivariate Random Variable



Q13. Consider an experiment of drawing randomly three balls from an urn containing two red, three white, and four blue balls. Let (X, Y) be a bivariate r.v. where X and Y denote, respectively, the number of red and white balls chosen.




$X = \text{No. of red ball}$
 $Y = \text{No. of white ball}$

- A. Find the range of (X, Y) .
- B. Find the joint pmf's of (X, Y) .
- C. Find the marginal pmf's of X and Y .
- D. Are X and Y independent?

$$P[X=x_i \cap Y=y_j] = P[X=x_i] P[Y=y_j]$$

marginal Prob.



$X = \text{No. of red ball}$
 $Y = \text{No. of white ball}$

$X = \text{No. of red ball } 0, 1, 2$
 $Y = \text{No. of white ball } 0, 1, 2, 3$

Joint Prob. Mass Function
 $P[X=x_i, Y=y_j]$ where

$$\begin{array}{l} i=0,1,2 \\ j=0,1,2,3 \end{array}$$

3 balls



$$\frac{4C_3}{9C_3}$$

$$P[X=0, Y=0] = P[X=\text{No red ball} \wedge Y=\text{No white ball}] \Rightarrow \frac{4C_3}{9C_3} = \frac{4}{84}$$

$$P[X=0, Y=1] \Rightarrow \frac{18}{84}$$

$$P[X=0, Y=2] = \frac{12}{84}$$

$$P[X=1, Y=0] = \frac{12}{84}$$

$$P[X=1, Y=1] = \frac{24}{84}$$

$$P[X=1, Y=2] = \frac{6}{84}$$

$$P[X=2, Y=0] = \frac{4}{84}$$

$X \setminus Y$	0	1	2	3
0	(0,0)	(0,1)	(0,2)	(0,3)
1	(1,0)	(1,1)	(1,2)	(1,3)
2	(2,0)	(2,1)	(2,2)	(2,3)

$$\text{Range} = \{(x, y)\}$$

$$P[X=2, Y=1] = \frac{3}{84}$$

$$P[X=2, Y=2] =$$



Topic : Problem Based on Bivariate Random Variable

Q16. Consider the binary communication channel shown in Fig. 3-4. Let (X, Y) be a bivariate r.v., where X is the input to the channel and Y is the output of the channel. Let $P(X = 0) = 0.5$, $P(Y = 1|X = 0) = 0.1$ and $P(Y = 0|X = 1) = 0.2$.

- (a) ✓ Find the joint pmf's of (X, Y)
- (b) ✓ Find the marginal pmf's of X and Y .
- (c) ✓ Are X and Y independent?

$$P\left(\frac{Y=1}{X=0}\right) = 0.1$$

$$P\left(\frac{Y=0}{X=0}\right) = 1 - 0.1 = 0.9$$

$X = \text{input channel}$

$Y = \text{output channel}$


$$P[X=1] = 1 - P[X=0] = 1 - 0.5 = 0.5$$

$$P\left(\frac{Y=0}{X=1}\right) = 0.2$$

$$P\left(\frac{Y=1}{X=1}\right) = 1 - 0.2 = 0.8$$



Topic : Problem Based on Bivariate Random Variable



$$\begin{aligned} P(X=0) &= 0.5 \\ P(X=1) &= 1 - P(X=0) \\ &= 1 - 0.5 = 0.5 \\ P\left(\frac{Y=1}{X=0}\right) &= 0.1 \\ P\left(\frac{Y=0}{X=1}\right) &= 0.2 \\ P\left(\frac{Y=0}{X=0}\right) &= 1 - 0.1 = 0.9 \\ P\left(\frac{Y=1}{X=1}\right) &= 1 - 0.2 = 0.8 \end{aligned}$$

$$P[X=0, Y=0] = P[X=0 \wedge Y=0]$$

$$\checkmark P\left[\frac{Y=0}{X=0}\right] = \frac{P[Y=0 \wedge X=0]}{P[X=0]} = \frac{0.9 \times 0.5}{0.45} = 0.45$$

$$\checkmark P\left[\frac{Y=0}{X=1}\right] = \frac{P[Y=0 \wedge X=1]}{P[X=1]} = 0.2 \times 0.5 = 0.1$$

		0	1
X	0	(0,0)	(0,1)
1	(1,0)	(1,1)	

0.45, 0.05, 0, 0.4

$$\checkmark P\left[\frac{Y=1}{X=1}\right] = \frac{P[Y=1 \wedge X=1]}{P[X=1]} = 0.8 \times 0.5$$

$$P\left[\frac{Y=1}{X=0}\right] = \frac{P[Y=1 \wedge X=0]}{P[X=0]} = \frac{0.1 \times 0.5}{0.45} = \underline{\underline{0.05}}$$

✓ X and Y Are Not Independent

THANK - YOU