

Spatio-temporal variability of zooplankton standing stock in  
eastern Arabian Sea inferred from ADCP backscatter  
measurements

February 20, 2024

**Abstract**

The study focuses on the zooplankton standing stock in the eastern Arabian sea (EAS) and aims to understand its spatio-temporal variation using ADCP(acoustic Doppler current profiler) backscatter measurements. The ADCP moorings were deployed at multiple locations on the continental slope of the west coast of India; of which we have used data from October 2017 to January 2023. The ADCP (operating frequency 153.3 kHz) uses backscatter from or sediments or organisms such as copepods, ctenophores, salps and amphipods greater than 1 cm to calculate current profile. The conversion from backscatter to biomass is based on volumetric zooplankton sampling at the respective locations. Analysis of the data over 25-140 m shows that the backscatter and zooplankton biomass decrease from the upper ocean ( $215 \text{ mg m}^{-3}$  biomass contour) to the lower depths. Seasonal variation is noticed in the monthly climatology zooplankton standing stock (integral of

<sup>12</sup> the biomass over the 20-140m water column) along with change as we move to northward slope  
<sup>13</sup> moorings in EAS. Complementary parameters (mixed layer depth, net primary production, Chl-a,  
<sup>14</sup> sea surface temperature) is used to explain the processes leading to growth or decay in zooplankton  
<sup>15</sup> biomass and on their migratory behaviour. Additionally, we have studied the effect of wind induced  
<sup>16</sup> vertical mixing events. The findings of this research will contribute to a better understanding of  
<sup>17</sup> the zooplankton dynamics in the EAS and provide valuable insights into the seasonal and annual  
<sup>18</sup> cycles of zooplankton standing stock.

<sup>19</sup> **1 Introduction**

<sup>20</sup> **1.1 Background**

<sup>21</sup> Zooplankton plays a vital role in food web of pelagic ecosystem by enabling the hierarchical trans-  
<sup>22</sup> port of organic matter from primary producers to higher trophic levels impacting the fish population  
<sup>23</sup> and the carbon pump of the deep ocean [Ohman and Hirche, 2001, Le Qu et al., 2016]. They are  
<sup>24</sup> presumably the largest migrating organisms in terms of biomass [Hays, 2003] which occurs in Diel  
<sup>25</sup> Vertical Migration (DVM). Zooplanktons depend not only on phytoplankton but other environ-  
<sup>26</sup> mental parameters (e.g. Mixed layer depth, insolation, Oxygen, thermocline, nutrient availability,  
<sup>27</sup> chlorophyll concentration and daily primary production). The biological productivity of the ocean  
<sup>28</sup> is essentially connected with physics and chemistry [Subrahmanyam, 1959, Ryther et al., 1966,  
<sup>29</sup> Qasim, 1977, Nair, 1970, Banse, 1995, McCreary et al., 2009, Vijith et al., 2016]. The dynamic  
<sup>30</sup> ocean results in varying physico-chemical properties, leading to bloom and growth of planktons  
<sup>31</sup> in favourable conditions. The changes are strongly influenced by the seasonal cycle in the North  
<sup>32</sup> Indian Ocean (NIO; north of 5 °N of Indian Ocean). The eastern boundary of Arabian Sea contains  
<sup>33</sup> the West India Coastal Current (WICC; [Patil et al., 1964, Ramamirtham and Murty, 1965, Banse,  
<sup>34</sup> 1968, Shetye et al., 1991, McCreary Jr et al., 1993, Shankar and Shetye, 1997, Shetye and Gouveia,  
<sup>35</sup> 1998, Maheswaran et al., 2000, Amol et al., 2014, Chaudhuri et al., 2020, 2021]) which reverses  
<sup>36</sup> seasonally, flowing poleward (equatorward) during November to February (June to September).  
<sup>37</sup> The direct consequence of this reversal is the seasonal cycle of thermocline, oxycline and thick-  
<sup>38</sup> ness of mixed Layer Depth (MLD) induced by upwelling favourable conditions in summer and down-  
<sup>39</sup> welling favourable conditions in winter in eastern Arabian Sea (EAS). Further, the phytoplankton  
<sup>40</sup> biomass and chlorophyll concentration changes with the season [Subrahmanyam and Sarma, 1960,

<sup>41</sup> Banse, 1968, Lévy et al., 2007, Vijith et al., 2016]. Upwelling in summer monsoon leads to maxi-  
<sup>42</sup> mum chlorophyll growth in the entire EAS [Banse, 1968, Banse and English, 2000, McCreary et al.,  
<sup>43</sup> 2009, Hood et al., 2017]. During winter monsoon, the convective mixing induced winter mixed layer  
<sup>44</sup> [Shetye et al., 1992, Madhupratap et al., 1996b, Lévy et al., 2007, Vijith et al., 2016, Shankar et al.,  
<sup>45</sup> 2016, Keerthi et al., 2017] results in winter chlorophyll peak in northern EAS (NEAS) while the  
<sup>46</sup> downwelling Rossby waves modulate chlorophyll along the southern EAS (SEAS) albeit limited to  
<sup>47</sup> coast and islands [Amol et al., 2020]. (For a detailed description on EAS division, please refer figure  
<sup>48</sup> 1 of [Shankar et al., 2019].

<sup>49</sup> The zooplankton grazing peak is instantaneous with no time delay from peak phytoplankton  
<sup>50</sup> production [Li et al., 2000], but its population growth lags [Rehim and Imran, 2012, Almén and  
<sup>51</sup> Tamelander, 2020] depending on its gestation period and other limiting aspects. While some studies  
<sup>52</sup> suggest that the peak timing of zooplankton may not change in parallel with phytoplankton blooms  
<sup>53</sup> [Winder and Schindler, 2004], others indicate that lag exists between primary production and the  
<sup>54</sup> transfer of energy to higher trophic levels [Brock and McClain, 1992, Brock et al., 1991]. A recent  
<sup>55</sup> work [Aparna et al., 2022] had shown that peak zooplankton population may never occur even  
<sup>56</sup> with a bloom in phytoplankton such as in SEAS, leading to the collapse of ecological models and  
<sup>57</sup> succeeding food webs of higher trophic levels.

<sup>58</sup> The conventional zooplankton measurements, where only few snapshot/s of the event is cap-  
<sup>59</sup> tured gives an incoherent or incomplete understanding in terms of spatio-temporal variation of  
<sup>60</sup> zooplankton [Ramamurthy, 1965, Piontkovski et al., 1995, Madhupratap et al., 1992, 1996a] as  
<sup>61</sup> much information is revealed by later studies [Jyothibabu et al., 2010, Vijith et al., 2016, Shankar  
<sup>62</sup> et al., 2019, Aparna et al., 2022] using high resolution data. Calibrated acoustic instruments such as  
<sup>63</sup> Acoustic Doppler Current Profiler (ADCP) along with relevant data can be utilised to understand

64 small scale variability [Nair et al., 1999, Edvardsen et al., 2003, Smith and Madhupratap, 2005,  
65 Smeti et al., 2015, Kang et al., 2024], the complex interplay between the physico-chemical parame-  
66 ters and ecosystem [Jiang et al., 2007, Potiris et al., 2018, Shankar et al., 2019, Aparna et al., 2022,  
67 Nie et al., 2023], the zooplankton migration [Ursella et al., 2018, 2021] and their seasonal to annual  
68 variation [Jiang et al., 2007, Hobbs et al., 2021, Liu et al., 2022, Aparna et al., 2022].

## 69 **1.2 ADCP backscatter and zooplankton biomass**

70 At present, there are two types of acoustic samplers: non-calibrated single frequency acoustic pro-  
71 filer such as ADCP or calibrated multi and mono frequency acoustic profilers such as zooplankton  
72 acoustic profiler (ZAP) and Tracor acoustic profiling system(TAPS). The use of acoustics as a proxy  
73 for zooplankton biomass estimation can be traced to [Pieper, 1971, Sameoto and Paulowich, 1977]  
74 and earlier studies which used echograms to approximate the large-scale horizontal extents [Barr-  
75 aclough et al., 1969], and small scale vertical extent [McNaught, 1968]. The relationship between  
76 backscatter and the abundance and size of zooplankton was described by [Greenlaw] wherein it was  
77 pointed out that single frequency backscatter can be used to estimate abundance if mean zooplank-  
78 ter size is known. This paved the way for use of single frequency acoustic profiler. A drastic increase  
79 in study temporal and spatial variation of zooplankton biomass using backscatter-proxy came in  
80 1990s by introduction of high frequency echosounders, with studies [Flagg and Smith, 1989, Wiebe  
81 et al., 1990, Batchelder et al., 1995, Greene et al., 1998, Rippeth and Simpson, 1998] methodically  
82 showing acoustic backscatter estimated zooplankton biomass in various shelf and slope locations  
83 around North Atlantic, North pacific location. The foundation for further research that investigated  
84 the potential of acoustic backscatter from ADCPs and multi frequency echosounders in assessing  
85 zooplankton biomass and comprehending zooplankton dynamics in diverse maritime habitats was

<sup>86</sup> established by these initial explorative experiments.

<sup>87</sup> Acoustic backscatter and zooplankton biomass have been better understood as a result of tech-  
<sup>88</sup> nological and methodological developments over time. Net sampling augmented ADCP backscatter  
<sup>89</sup> have been used to study DVM and the spatial and temporal variability of zooplankton biomass by  
<sup>90</sup> [Cisewski et al., 2010, Smeti et al., 2015, Guerra et al., 2019] in various marine regions, such as the  
<sup>91</sup> Southwestern Pacific, the Lazarev Sea in Antarctica and the Corsica Channel in the north-western  
<sup>92</sup> Mediterranean Sea. The zooplankton biomass variation in the Arabian sea has been studied during  
<sup>93</sup> JGOFS programme in 1990s [Herring et al., 1998, Nair et al., 1999, Fielding et al., 2004, Smith and  
<sup>94</sup> Madhupratap, 2005]. However, their studies were limited to the cruise duration as vessel mounted  
<sup>95</sup> ADCPs were predominantly used; hence longterm data was sparsely produced. The first such study  
<sup>96</sup> to fully exploit the immense potential of ADCPs in EAS was carried out by [Aparna et al., 2022]  
<sup>97</sup> using ADCP moorings deployed on continental slopes off the Indian west coasts [Amol et al., 2014,  
<sup>98</sup> Chaudhuri et al., 2020].

### <sup>99</sup> **1.3 Objective and scope of the manuscript**

<sup>100</sup> A network of ADCPs has been installed off the continental slope and shelf on the west coast of  
<sup>101</sup> India. This ADCPs have enabled a rigorous view of intraseasonal to seasonal scale variability [Amol  
<sup>102</sup> et al., 2014, Chaudhuri et al., 2020]. Initially a network of 4 ADCPs (off Mumbai, Goa, Kollam and  
<sup>103</sup> Kanyakumari) on continental slope, it has been extended to include 3 more moorings (off Okha from  
<sup>104</sup> 2018, Jaigarh and Udupi from 2017). In the recent study [Aparna et al., 2022] have used ADCP  
<sup>105</sup> moorings off Mumbai, Goa and Kollam to explain the temporal variability of zooplankton biomass.  
<sup>106</sup> The study showed that the zooplankton peaks (and troughs) is not only non-uniform in latitude but  
<sup>107</sup> also heavily influenced by the oxygen minimum zone, MLD and the seasonal upwelling/downwelling

108 conditions. Stark contrast in the phytoplankton bloom and subsequence growth of zooplankton or  
109 the lack thereof was observed in the EAS regimes.

110 We build upon the existing work by extending to include the newly incorporated ADCPs so as to  
111 have a better understanding in the latitudinal variation of zooplankton biomass in EAS. The paper  
112 is organized as follows, datasets and methods employed are described in detail in Secion 2. Section  
113 3 describes the observed seasonal cycle of zooplankton biomass and standing stock. The role of  
114 mixed layer depth, net primary production, sea surface temperature, wind forcing and circulation  
115 in determining the biomass is discussed in results section 4, with conclusion in section 5.

116 **2 Data and methods**

## <sup>117</sup> References

- <sup>118</sup> Anna-Karin Almén and Tobias Tamelander. Temperature-related timing of the spring bloom and match between  
<sup>119</sup> phytoplankton and zooplankton. *Marine Biology Research*, 16(8-9):674–682, 2020.
- <sup>120</sup> P Amol, D Shankar, V Fernando, A Mukherjee, SG Aparna, R Fernandes, GS Michael, ST Khalap, NP Satelkar,  
<sup>121</sup> Y Agarvadekar, et al. Observed intraseasonal and seasonal variability of the west india coastal current on the  
<sup>122</sup> continental slope. *Journal of Earth System Science*, 123(5):1045–1074, 2014.
- <sup>123</sup> P Amol, Suchandan Bemal, D Shankar, V Jain, V Thushara, V Vijith, and PN Vinayachandran. Modulation of  
<sup>124</sup> chlorophyll concentration by downwelling rossby waves during the winter monsoon in the southeastern arabian  
<sup>125</sup> sea. *Progress in Oceanography*, 186:102365, 2020.
- <sup>126</sup> SG Aparna, DV Desai, D Shankar, AC Anil, Shrikant Dora, and R Khedekar. Seasonal cycle of zooplankton standing  
<sup>127</sup> stock inferred from adcp backscatter measurements in the eastern arabian sea. *Progress in Oceanography*, 203:  
<sup>128</sup> 102766, 2022.
- <sup>129</sup> K Banse and DC English. Geographical differences in seasonality of czcs-derived phytoplankton pigment in the  
<sup>130</sup> arabian sea for 1978–1986. *Deep Sea Research Part II: Topical Studies in Oceanography*, 47(7-8):1623–1677, 2000.
- <sup>131</sup> Karl Banse. Hydrography of the arabian sea shelf of india and pakistan and effects on demersal fishes. In *Deep sea*  
<sup>132</sup> *research and oceanographic Abstracts*, volume 15, pages 45–79. Elsevier, 1968.
- <sup>133</sup> Karl Banse. Zooplankton: pivotal role in the control of ocean production: I. biomass and production. *ICES Journal*  
<sup>134</sup> *of marine Science*, 52(3-4):265–277, 1995.
- <sup>135</sup> WE Barraclough, RJ LeBrasseur, and OD Kennedy. Shallow scattering layer in the subarctic pacific ocean: detection  
<sup>136</sup> by high-frequency echo sounder. *Science*, 166(3905):611–613, 1969.
- <sup>137</sup> H. P. Batchelder, J. R. VanKeuren, R. D. Vaillancourt, and E. Swift. Spatial and temporal distributions of acoustically  
<sup>138</sup> estimated zooplankton biomass near the marine light-mixed layers station (59°30'n, 21°00'w) in the north atlantic  
<sup>139</sup> in may 1991. *Journal of Geophysical Research: Oceans*, 100:6549–6563, 1995. doi: 10.1029/94jc00981.
- <sup>140</sup> John C Brock and Charles R McClain. Interannual variability in phytoplankton blooms observed in the northwestern  
<sup>141</sup> arabian sea during the southwest monsoon. *Journal of Geophysical Research: Oceans*, 97(C1):733–750, 1992.
- <sup>142</sup> John C Brock, Charles R McClain, Mark E Luther, and William W Hay. The phytoplankton bloom in the north-  
<sup>143</sup> western arabian sea during the southwest monsoon of 1979. *Journal of Geophysical Research: Oceans*, 96(C11):  
<sup>144</sup> 20623–20642, 1991.
- <sup>145</sup> Anya Chaudhuri, D Shankar, SG Aparna, P Amol, V Fernando, A Kankonkar, GS Michael, NP Satelkar, ST Khalap,  
<sup>146</sup> AP Tari, et al. Observed variability of the west india coastal current on the continental slope from 2009–2018.  
<sup>147</sup> *Journal of Earth System Science*, 129(1):57, 2020.

- 148 Anya Chaudhuri, P Amol, D Shankar, S Mukhopadhyay, SG Aparna, V Fernando, and A Kankonkar. Observed  
149 variability of the west india coastal current on the continental shelf from 2010–2017. *Journal of Earth System*  
150 *Science*, 130:1–21, 2021.
- 151 Boris Cisewski, Volker H Strass, Monika Rhein, and Sören Krägesky. Seasonal variation of diel vertical migration  
152 of zooplankton from adcp backscatter time series data in the lazarev sea, antarctica. *Deep Sea Research Part I:*  
153 *Oceanographic Research Papers*, 57(1):78–94, 2010.
- 154 A Edvardsen, D Slagstad, KS Tande, and P Jaccard. Assessing zooplankton advection in the barents sea using  
155 underway measurements and modelling. *Fisheries Oceanography*, 12(2):61–74, 2003.
- 156 S Fielding, G Griffiths, and HSJ Roe. The biological validation of adcp acoustic backscatter through direct comparison  
157 with net samples and model predictions based on acoustic-scattering models. *ICES Journal of Marine Science*,  
158 61(2):184–200, 2004.
- 159 Charles N Flagg and Sharon L Smith. On the use of the acoustic doppler current profiler to measure zooplankton  
160 abundance. *Deep Sea Research Part A. Oceanographic Research Papers*, 36(3):455–474, 1989.
- 161 Charles H Greene, Peter H Wiebe, Chris Pelkie, Mark C Benfield, and Jacqueline M Popp. Three-dimensional  
162 acoustic visualization of zooplankton patchiness. *Deep Sea Research Part II: Topical Studies in Oceanography*, 45  
163 (7):1201–1217, 1998.
- 164 Charles F Greenlaw. Acoustical estimation of zooplankton populations 1.
- 165 Davide Guerra, Katrin Schroeder, Mireno Borghini, Elisa Camatti, Marco Pansera, Anna Schroeder, Stefania  
166 Sparnocchia, and Jacopo Chiggiato. Zooplankton diel vertical migration in the corsica channel (north-western  
167 mediterranean sea) detected by a moored acoustic doppler current profiler. *Ocean Science*, 15(3):631–649, 2019.
- 168 Graeme C Hays. A review of the adaptive significance and ecosystem consequences of zooplankton diel vertical  
169 migrations. In *Migrations and Dispersal of Marine Organisms: Proceedings of the 37 th European Marine Biology*  
170 *Symposium held in Reykjavík, Iceland, 5–9 August 2002*, pages 163–170. Springer, 2003.
- 171 PJ Herring, MJR Fasham, AR Weeks, JCP Hemmings, HSJ Roe, PR Pugh, S Holley, NA Crisp, and MV Angel.  
172 Across-slope relations between the biological populations, the euphotic zone and the oxygen minimum layer off  
173 the coast of oman during the southwest monsoon (august, 1994). *Progress in Oceanography*, 41(1):69–109, 1998.
- 174 Laura Hobbs, Neil S Banas, Jonathan H Cohen, Finlo R Cottier, Jørgen Berge, and Øystein Varpe. A marine  
175 zooplankton community vertically structured by light across diel to interannual timescales. *Biology Letters*, 17(2):  
176 20200810, 2021.
- 177 Raleigh R Hood, Lynnath E Beckley, and Jerry D Wiggert. Biogeochemical and ecological impacts of boundary  
178 currents in the indian ocean. *Progress in Oceanography*, 156:290–325, 2017.

- 179 Songnian Jiang, Tommy D Dickey, Deborah K Steinberg, and Laurence P Madin. Temporal variability of zooplankton  
180 biomass from adcp backscatter time series data at the bermuda testbed mooring site. *Deep Sea Research Part I: Oceanographic Research Papers*, 54(4):608–636, 2007.
- 182 R Jyothibabu, NV Madhu, H Habeebrehman, KV Jayalakshmy, KKC Nair, and CT Achuthankutty. Re-evaluation  
183 of ‘paradox of mesozooplankton’ in the eastern arabian sea based on ship and satellite observations. *Journal of Marine Systems*, 81(3):235–251, 2010.
- 185 Myounghee Kang, Sunyoung Oh, Wooseok Oh, Dong-Jin Kang, SungHyun Nam, and Kyounghoon Lee. Acoustic  
186 characterization of fish and macroplankton communities in the seychelles-chagos thermocline ridge of the southwest  
187 indian ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 213:105356, 2024.
- 188 Madhavan Girijakumari Keerthi, Matthieu Lengaigne, Marina Levy, Jerome Vialard, Vallivattathillam Parvathi,  
189 Clément de Boyer Montégut, Christian Ethé, Olivier Aumont, Iyyappan Suresh, Valiya Parambil Akhil, et al.  
190 Physical control of interannual variations of the winter chlorophyll bloom in the northern arabian sea. *Biogeosciences*, 14(15):3615–3632, 2017.
- 192 Corinne Le Qu, Robbie M Andrew, Josep G Canadell, Stephen Sitch, Jan Ivar Korsbakken, Glen P Peters, Andrew C  
193 Manning, Thomas A Boden, Pieter P Tans, Richard A Houghton, et al. Global carbon budget 2016. *Earth System  
194 Science Data*, 8(2):605–649, 2016.
- 195 Marina Lévy, D Shankar, J-M André, SSC Shenoi, Fabien Durand, and Clément de Boyer Montégut. Basin-wide  
196 seasonal evolution of the indian ocean’s phytoplankton blooms. *Journal of Geophysical Research: Oceans*, 112  
197 (C12), 2007.
- 198 M Li, A Gargett, and K Denman. What determines seasonal and interannual variability of phytoplankton and  
199 zooplankton in strongly estuarine systems? *Estuarine, Coastal and Shelf Science*, 50(4):467–488, 2000.
- 200 Yanliang Liu, Jingsong Guo, Yuhuan Xue, Chalermrat Sangmanee, Huiwu Wang, Chang Zhao, Somkiat Khokiat-  
201 tiwong, and Weidong Yu. Seasonal variation in diel vertical migration of zooplankton and micronekton in the  
202 andaman sea observed by a moored adcp. *Deep Sea Research Part I: Oceanographic Research Papers*, 179:103663,  
203 2022.
- 204 M Madhupratap, P Haridas, Neelam Ramaiah, and CT Achuthankutty. Zooplankton of the southwest coast of india:  
205 abundance, composition, temporal and spatial variability in 1987. 1992.
- 206 M Madhupratap, TC Gopalakrishnan, P Haridas, KKC Nair, PN Aravindakshan, G Padmavati, and Shiney Paul.  
207 Lack of seasonal and geographic variation in mesozooplankton biomass in the arabian sea and its structure in the  
208 mixed layer. *Current science. Bangalore*, 71(11):863–868, 1996a.
- 209 M Madhupratap, S Prasanna Kumar, PMA Bhattathiri, M Dileep Kumar, S Raghukumar, KKC Nair, and N Rama-  
210 iah. Mechanism of the biological response to winter cooling in the northeastern arabian sea. *Nature*, 384(6609):  
211 549–552, 1996b.

- 212 PA Maheswaran, G Rajesh, C Revichandran, and KKC Nair. Upwelling and associated hydrography along the west  
213 coast of india during southwest monsoon, 1999. 2000.
- 214 JP McCreary, Raghu Murtugudde, Jerome Vialard, PN Vinayachandran, Jerry D Wiggert, Raleigh R Hood,  
215 D Shankar, and S Shetye. Biophysical processes in the indian ocean. *Indian Ocean biogeochemical processes*  
216 and ecological variability, 185:9–32, 2009.
- 217 Julian P McCreary Jr, Pijush K Kundu, and Robert L Molinari. A numerical investigation of dynamics, thermody-  
218 namics and mixed-layer processes in the indian ocean. *Progress in Oceanography*, 31(3):181–244, 1993.
- 219 Donald C McNaught. Acoustical determination of zooplankton distribution. In *Proc. 11th Conf. Great lakes Res.*,  
220 pages 76–84, 1968.
- 221 KKC Nair, M Madhupratap, TC Gopalakrishnan, P Haridas, and Mangesh Gauns. The arabian sea: physical  
222 environment, zooplankton and myctophid abundance. 1999.
- 223 PV Nair. Primary productivity in the indian seas. *CMFRI Bulletin*, 22:1–63, 1970.
- 224 Lingyun Nie, Jianchao Li, Hao Wu, Wenchoao Zhang, Yongjun Tian, Yang Liu, Peng Sun, Zhenjiang Ye, Shuyang  
225 Ma, and Qinfeng Gao. The influence of ocean processes on fine-scale changes in the yellow sea cold water mass  
226 boundary area structure based on acoustic observations. *Remote Sensing*, 15(17):4272, 2023.
- 227 MD Ohman and H-J Hirche. Density-dependent mortality in an oceanic copepod population. *Nature*, 412(6847):  
228 638–641, 2001.
- 229 MR Patil, CP Ramamirtham, P Udaya Varma, and CP Nair. Hydrography of the west coast of india during the  
230 pre-monsoon period of the year 1962. *Journal of marine biological association of India*, 6(1):151–164, 1964.
- 231 Richard Edward Pieper. *A study of the relationship between zooplankton and high-frequency scattering of underwater*  
232 *sound*. PhD thesis, University of British Columbia, 1971.
- 233 SA Piontkovski, R Williams, and TA Melnik. Spatial heterogeneity, biomass and size structure of plankton of the  
234 indian ocean: some general trends. *Marine ecology progress series*, pages 219–227, 1995.
- 235 Emmanuel Potiris, Constantin Frangoulis, Alkiviadis Kalampokis, Manolis Ntoumas, Manos Pettas, George Peti-  
236 hakis, and Vassilis Zervakis. Acoustic doppler current profiler observations of migration patterns of zooplankton  
237 in the cretan sea. *Ocean Science*, 14(4):783–800, 2018.
- 238 SZ Qasim. Biological productivity of the indian ocean. 1977.
- 239 CP Ramamirtham and AVS Murty. Hydrography of the west coast of india during the pre-monsoon period of the  
240 year 1962—part 2: in and offshore waters of the konkan and malabar coasts. *Journal of the Marine Biological*  
241 *Association of India*, 7(1):150–168, 1965.
- 242 S Ramamurthy. Studies on the plankton of the north kanara coast in relation to the pelagic fishery. *Journal of*  
243 *Marine Biological Association of India*, 7(1):127–149, 1965.

- 244 Mehbuba Rehim and Mudassar Imran. Dynamical analysis of a delay model of phytoplankton–zooplankton interac-  
245 tion. *Applied Mathematical Modelling*, 36(2):638–647, 2012.
- 246 T. P. Rippeth and J. H. Simpson. Diurnal signals in vertical motions on the hebridean shelf. *Limnology and*  
247 *Oceanography*, 43:1690–1696, 1998. doi: 10.4319/lo.1998.43.7.1690.
- 248 John H Ryther, John R Hall, Allan K Pease, Andrew Bakun, and Mark M Jones. Primary organic production in  
249 relation to the chemistry and hydrography of the western indian ocean 1. *Limnology and Oceanography*, 11(3):  
250 371–380, 1966.
- 251 D Sameoto and S Paulowich. The use of 120 khz sonar in zooplankton studies. In *OCEANS'77 Conference Record*,  
252 pages 523–528. IEEE, 1977.
- 253 D Shankar and SR Shetye. On the dynamics of the lakshadweep high and low in the southeastern arabian sea.  
254 *Journal of Geophysical Research: Oceans*, 102(C6):12551–12562, 1997.
- 255 D Shankar, R Remya, PN Vinayachandran, Abhisek Chatterjee, and Ambica Behera. Inhibition of mixed-layer  
256 deepening during winter in the northeastern arabian sea by the west india coastal current. *Climate Dynamics*,  
257 47:1049–1072, 2016.
- 258 D Shankar, R Remya, AC Anil, and V Vijith. Role of physical processes in determining the nature of fisheries in the  
259 eastern arabian sea. *Progress in Oceanography*, 172:124–158, 2019.
- 260 SR Shetye and AD Gouveia. Coastal circulation in the north indian ocean: Coastal segment (14, sw). John Wiley  
261 and Sons, New York, USA, 1998.
- 262 SR Shetye, AD Gouveia, SSC Shenoi, GS Michael, D Sundar, AM Almeida, and K Santanam. The coastal current  
263 off western india during the northeast monsoon. *Deep Sea Research Part A. Oceanographic Research Papers*, 38  
264 (12):1517–1529, 1991.
- 265 SR Shetye, AD Gouveia, and SSC Shenoi. Does winter cooling lead to the subsurface salinity minimum off saurashtra,  
266 india? *Oceanography of the Indian Ocean*, pages 617–625, 1992.
- 267 Houssem Smeti, Marc Pagano, Christophe Menkes, Anne Lebourges-Dhaussy, Brian PV Hunt, Valerie Allain, Martine  
268 Rodier, Florian De Boissieu, Elodie Kestenare, and Cherif Sammari. Spatial and temporal variability of zooplank-  
269 ton off n ew c alifornia (s outhwestern p acific) from acoustics and net measurements. *Journal of Geophysical*  
270 *Research: Oceans*, 120(4):2676–2700, 2015.
- 271 SL Smith and M Madhupratap. Mesozooplankton of the arabian sea: patterns influenced by seasons, upwelling, and  
272 oxygen concentrations. *Progress in Oceanography*, 65(2-4):214–239, 2005.
- 273 R Subrahmanyam. Studies on the phytoplankton of the west coast of india: Part ii. physical and chemical factors  
274 influencing the production of phytoplankton, with remarks on the cycle of nutrients and on the relationship of  
275 the phosphate-content to fish landings. In *Proceedings/Indian Academy of Sciences*, volume 50, pages 189–252.  
276 Springer, 1959.

- 277 R Subrahmanyam and AH Sarma. Studies on the phytoplankton of the west coast of india. part iii. seasonal variation  
278 of the phytoplankters and environmental factors. *Indian Journal of Fisheries*, 7(2):307–336, 1960.
- 279 Laura Ursella, Vanessa Cardin, Mirna Batistić, Rade Garić, and Miroslav Gačić. Evidence of zooplankton vertical  
280 migration from continuous southern adriatic buoy current-meter records. *Progress in oceanography*, 167:78–96,  
281 2018.
- 282 Laura Ursella, Sara Pensieri, Enric Pallàs-Sanz, Sharon Z Herzka, Roberto Bozzano, Miguel Tenreiro, Vanessa Cardin,  
283 Julio Candela, and Julio Sheinbaum. Diel, lunar and seasonal vertical migration in the deep western gulf of mexico  
284 evidenced from a long-term data series of acoustic backscatter. *Progress in Oceanography*, 195:102562, 2021.
- 285 V Vijith, PN Vinayachandran, V Thushara, P Amol, D Shankar, and AC Anil. Consequences of inhibition of mixed-  
286 layer deepening by the west india coastal current for winter phytoplankton bloom in the northeastern arabian sea.  
287 *Journal of Geophysical Research: Oceans*, 121(9):6583–6603, 2016.
- 288 Peter H Wiebe, Charles H Greene, Timothy K Stanton, and Janusz Burczynski. Sound scattering by live zooplankton  
289 and micronekton: empirical studies with a dual-beam acoustical system. *The Journal of the Acoustical Society of  
290 America*, 88(5):2346–2360, 1990.
- 291 Monika Winder and Daniel E Schindler. Climatic effects on the phenology of lake processes. *Global change biology*,  
292 10(11):1844–1856, 2004.

Table 1: ADCP deployment details at the locations. The temporal resolution is 1 hour, bin size(vertical resolution) 4 m. All ADCPs are operated at 153.3 kHz. The moorings are at a water column depth of 950 - 1200 m on the continental slope and are serviced on yearly basis according to ship availability. The 6th column consists of Reference echo intensity (Er) for each beam, while the 7th column contains the corresponding RSSI conversion factor [?].

Station (Position; °E, °N)	Date		Depth			
	Deployment	Recovery	Ocean	ADCP	Er	Kc
Okha (67.47, 22.26)	01/10/2018	01/12/2019	996	118	37 , 37 , 37 , 36	0.42 , 0.44 , 0.42 , 0.43
	01/12/2019	04/12/2020	1166	312	39 , 36 , 38 , 36	0.42 , 0.44 , 0.42 , 0.43
	04/12/2020	08/03/2022	1021	144	41 , 37 , 38 , 37	0.42 , 0.44 , 0.42 , 0.43
	08/03/2022	01/01/2023	1019	142	37 , 38 , 39 , 36	0.42 , 0.44 , 0.42 , 0.43
Mumbai (69.24, 20.01)	09/11/2017	29/09/2018	1025	150	36 , 34 , 39 , 42	0.40 , 0.40 , 0.40 , 0.40
	29/09/2018	29/11/2019	1122	125	35 , 36 , 39 , 42	0.40 , 0.40 , 0.40 , 0.40
	29/11/2019	02/12/2020	1143	164	37 , 34 , 39 , 43	0.40 , 0.40 , 0.40 , 0.40
	02/12/2020	06/03/2022	1125	142	36 , 34 , 39 , 42	0.40 , 0.40 , 0.40 , 0.40
	07/03/2022	02/01/2023	1103	158	37 , 34 , 40 , 43	0.40 , 0.40 , 0.40 , 0.40
Jaigarh (71.12, 17.53)	27/10/2017	27/09/2018	1039	198	32 , 35 , 33 , 32	0.45 , 0.45 , 0.45 , 0.45
	27/09/2018	30/10/2019	1032	164	32 , 35 , 33 , 31	0.45 , 0.45 , 0.45 , 0.45
	03/11/2019	30/11/2020	1142	264	32 , 36 , 33 , 32	0.45 , 0.45 , 0.45 , 0.45
	30/11/2020	05/03/2022	1099	119	33 , 36 , 34 , 32	0.45 , 0.45 , 0.45 , 0.45
	03/04/2022	26/06/2022	1120	136	68 , 71 , 69 , 66	0.45 , 0.45 , 0.45 , 0.45
Goa (72.74, 15.17)	03/10/2017	25/09/2018	1000	174	35 , 37 , 34 , 35	0.44 , 0.44 , 0.40 , 0.41
	25/09/2018	16/10/2019	969	145	38 , 36 , 36 , 34	0.44 , 0.44 , 0.40 , 0.41
	16/10/2019	29/11/2020	966	143	44 , 38 , 36 , 43	0.44 , 0.44 , 0.40 , 0.41
	29/11/2020	03/03/2022	985	157	35 , 40 , 35 , 38	0.44 , 0.44 , 0.40 , 0.41
	03/03/2022	05/01/2023	984	159	35 , 38 , 35 , 34	0.44 , 0.44 , 0.40 , 0.41
Udupi (74.04, 12.5)	05/10/2017	06/10/2018	1028	176	44 , 46 , 29 , 35	0.45 , 0.45 , 0.45 , 0.45
	06/10/2018	18/10/2019	1027	179	32 , 38 , 30 , 36	0.45 , 0.45 , 0.45 , 0.45
	18/10/2019	11/12/2020	1018	168	33 , 37 , 31 , 38	0.45 , 0.45 , 0.45 , 0.45
	11/03/2022	06/01/2023	1036	155	31 , 32 , 32 , 33	0.45 , 0.45 , 0.45 , 0.45
Kollam (75.44, 9.05)	07/10/2017	08/10/2018	1174	200	43 , 55 , 45 , 43	0.49 , 0.50 , 0.49 , 0.50
	08/10/2018	20/10/2019	1160	123	49 , 62 , 46 , 46	0.49 , 0.50 , 0.49 , 0.50
	20/10/2019	13/12/2020	1209	176	52 , 61 , 54 , 55	0.49 , 0.50 , 0.49 , 0.50
	13/12/2020	13/03/2022	1129	91	49 , 51 , 46 , 47	0.49 , 0.50 , 0.49 , 0.50
	13/03/2022	08/01/2023	1149	164	41 , 48 , 43 , 41	0.49 , 0.50 , 0.49 , 0.50
Kanyakumari (77.39, 6.96)	16/11/2016	08/10/2017	1096	252	37 , 36 , 37 , 37	0.42 , 0.44 , 0.42 , 0.43
	08/10/2017	10/10/2018	1055	181	32 , 34 , 38 , 35	0.45 , 0.45 , 0.45 , 0.45
	10/10/2018	22/10/2019	1075	180	36 , 34 , 39 , 36	0.45 , 0.45 , 0.45 , 0.45
	22/10/2019	14/12/2020	1060	167	33 , 35 , 36 , 35	0.45 , 0.45 , 0.45 , 0.45
	14/12/2020	14/03/2022	1184	287	34 , 36 , 36 , 35	0.45 , 0.45 , 0.45 , 0.45
	14/03/2022	10/01/2023	1069	172	33 , 36 , 42 , 36	0.45 , 0.45 , 0.45 , 0.45

Table 2:  
Volumetric samples of zooplankton of various stations. The tags corresponds to cruise and particular station. The sampling depth range is standardised for later years for bin range of 0-25m, 25-50m, 50-75m, 75-100m, 100-150m

Sample number	Tag	Lat( $^{\circ}$ N)	Lon( $^{\circ}$ E)	Date	Time (IST)	Sampling depth range (m)
1-3	G1	15.18	72.79	25 Sep 18	452	50–25, 100–50, 150–100
4-6	G2	15.16	72.71	25 Sep 18	2108	50–25, 100–50, 150–100
7-10	G2	15.16	72.71	25 Sep 18	2137	40–20, 60–40, 80–60, 100–80
11-14	J1			26 Sep 18	2000	40–20, 60–40, 80–60, 100–80
15-17	J2			27 Sep 18	2000	50–25, 100–50, 150–100
18-21	J2			27 Sep 18	2100	40–20, 60–40, 80–60, 100–80
22-25	M1	20	69.19	28 Sep 18	2135	40–20, 60–40, 80–60, 100–80
26-27	M1	20	69.19	28 Sep 18	2205	50–25, 100–50
28-29	M2	20.01	69.2	29 Sep 18	2035	50–25, 100–50
30-33	M2	20.01	69.2	29 Sep 18	2057	40–20, 60–40, 80–60, 100–80
34-37	U1			5 Oct 18	2000	40–20, 60–40, 80–60, 100–80
38-40	U1			5 Oct 18	2100	50–25, 100–50, 150–100
41-43	U2			6 Oct 18	2000	50–25, 100–50, 150–100
44-47	U2			6 Oct 18	2100	40–20, 60–40, 80–60, 100–80
48-51	K1	9.06	75.42	8 Oct 18	421	40–20, 60–40, 80–60, 100–80
52-54	K1	9.06	75.42	8 Oct 18	449	50–25, 100–50, 150–100
55-56	K2	9.04	75.4	8 Oct 18	2027	50–25, 100–50
57-60	K2	9.04	75.4	8 Oct 18	2045	40–20, 60–40, 80–60, 100–80
61-64	G2	15.16	72.74	16 Oct 19	829	50–25, 75–50, 100–75, 150–100
65-67	G3	15.16	72.74	16 Oct 19	1812	50–25, 75–50, 100–75
68-70	K2	9.02	75.42	20 Oct 19	840	50–25, 75–50, 100–75
71-74	K3	9.04	75.43	20 Oct 19	1934	50–25, 75–50, 100–75, 150–100
75-78	KK1			22 Oct 19	742	50–25, 75–50, 100–75, 150–100
79-82	KK2			22 Oct 19	1925	50–25, 75–50, 100–75, 150–100
83-86	J1			30 Oct 19	324	50–25, 75–50, 100–75, 150–100
87-89	J2			4 Nov 19	946	75–50, 100–75, 150–100
90-92	M2	19.98	69.22	29 Nov 19	1434	50–25, 75–50, 100–75
93-96	M3	20.01	69.23	30 Nov 19	958	50–25, 75–50, 100–75, 150–100
97-100	O1	22.24	67.49	1 Dec 19	937	50–25, 75–50, 100–75, 150–100
101	O2	22.25	67.46	1 Dec 19	1957	150-100
102-105	G3	15.68	73.22	28 Nov 20	930	50–25, 75–50, 100–75, 150–100
105-108	G4	15.32	73.22	29 Nov 20	1558	50–25, 75–50, 100–75, 150–100
108-110	J2	17.85	71.21	30 Nov 20	1458	75–50, 100–75, 150–100
111-114	J3	17.91	71.21	1 Dec 20	1052	50–25, 75–50, 100–75, 150–100
115-118	M4	20.03	69.38	2 Dec 20	2016	50–25, 75–50, 100–75, 150–100
119-00	O2	22.41	67.8	4 Dec 20	953	150-100
120-123	O3	22.41	67.79	4 Dec 20	2011	50–25, 75–50, 100–75, 150–100
124-127	K3	9.11	75.72	12 Dec 20	2335	50–25, 75–50, 100–75, 150–100
128-131	K4	9.06	75.74	13 Dec 20	1507	50–25, 75–50, 100–75, 150–100
132-134	KK1	7.62	77.63	14 Dec 20	1226	50–25, 75–50
135-138	KK2	7.62	77.63	14 Dec 20	2047	50–25, 75–50, 100–75, 150–100
139-142	G4	15.32	73.21	3 Mar 22	823	50–25, 75–50, 100–75, 150–100
143-146	G5	15.68	73.21	4 Mar 22	1030	50–25, 75–50, 100–75, 150–100
147-150	M5	19.99	69.23	7 Mar 22	957	50–25, 75–50, 100–75, 150–100
151-154	O3	22.24	67.5	8 Mar 22	806	50–25, 75–50, 100–75, 150–100
155-158	U3	12.5	74.04	12 Mar 22	1156	50–25, 75–50, 100–75, 150–100
159-160	K4	9.04	75.42	13 Mar 22	1027	50–25, 75–50, 100–75
161-164	KK3	6.97	77.4	15 Mar 22	1220	50–25, 75–50, 100–75, 150–100




Figure 1: Map showing region of interest. The slope moorings are deployed at 1000 m depth.




Figure 2: The linear fit line of Biomass (taken in log of biomass) and Backscatter. The linear fit line is within the error range of previous result of [Aparna et al., 2022] onto which latest zooplankton volumetric sample data is added.




Figure 3: The Daily and monthly averaged biomass for EAS moorings, north (top) to south (bottom). The dark contours are marking  $215 \text{ mg m}^{-3}$ .




Figure 4: The daily biomass at depth of 40 m and 104 m for all locations shown by grey and cyan curves. The black and blue lines shows the 30 day rolling averaged biomass.




Figure 5