

Topics in Set Theory

February 16, 2019

Contents

0	Introduction	3
1	Model theory of set theory	5
1.1	Absoluteness	5
1.2	Long List of Δ_0^T formulas	8
1.3	Absoluteness of well-foundedness	10
1.4	Concrete transitive models of ZFC	11
1.5	Inaccessible cardinals	13
1.6	The second construction: Models of hereditarily small sets	15
2	Example Class 1	20
2.1	Question 1	20
2.2	Question 2	20
2.3	Question 3	20
2.4	Question 6	20
2.5	Question 8	20
2.6	Question 9	21
2.7	Question 10	21
2.8	Question 11	21
2.9	Question 12	21

0 Introduction

—Lecture 1—

Example classes: 4th Feb, 18th Feb, 4th Mar 330-5pm MR5; fourth class undecided (probably on 15th).

Although the name of this course is *Topics* in Set theory, for all of its history only one topic is discussed. So maybe this course should be called *One Topic in Set Theory*, or probably just *the Continuum Hypothesis*: in this course we'll just solve one problem: the continuum problem, which we've known in the end that the problem is independent from ZFC.

Let's have some background stories first. In the second ICM congress (1900, Paris), Hilbert posed the famous 23 Hilbert questions, with the first one being the Continuum Hypothesis (a hypothesis at that time). The original formulation of CH was:

Any infinite subset of real numbers is either equinumerous to the set of natural numbers, or to the set of real numbers.

We could definitely formulate it better, but that is less important. More modern version of CH would be a short equation

$$2^{\aleph_0} = \aleph_1$$

which seemingly has nothing to do with the previous problem. However, in ZFC these two statements are equivalent:

- if $2^{\aleph_0} > \aleph_1$, in particular, $2^{\aleph_0} \geq \aleph_2$. Since $2^{\aleph_0} \sim \mathbb{R}$, we get an injection $i : \aleph_2 \rightarrow \mathbb{R}$. Consider $X := i[\aleph_1] \subseteq \mathbb{R}$. Clearly, $i \upharpoonright \aleph_1$ (i restricted to \aleph_1) is a bijection between \aleph_1 and X , so $X \sim \aleph_1$; but \aleph_1 , being uncountable, is not in bijection with natural numbers, and is not in bijection with real numbers. Thus X refutes CH.

If $2^{\aleph_0} = \aleph_1$, let $X \subseteq \mathbb{R}$. Consider $b : 2^{\aleph_0} \rightarrow \mathbb{R}$ a bijection. If X is infinite, then $b^{-1}[X] \subseteq 2^{\aleph_0}$. Thus the cardinality of X is either \aleph_0 or \aleph_1 (which $\sim \mathbb{N}$ and \mathbb{R} respectively). So $2^{\aleph_0} = \aleph_1 \implies \text{CH}$.

In 1938, Gödel proved that ZFC does not prove $\neg\text{CH}$, and in 1961 Cohen proved that ZFC does not prove CH, by methods of *inner models* and *forcing* (sometimes also called *outer models*, which is not incorrect) respectively. The latter has become the most important method in Set theory since then.

From logic (see Part II Logic and Set Theory) we have Gödel's Completeness Theorem: a theory T is consistent iff it has a model. So from the above two statements, it seems that we're going to prove that there are models for ZFC+CH and ZFC+ $\neg\text{CH}$; but this is obviously not possible because of the incompleteness phenomenon: we know we can't prove the consistency of ZFC (as a result, we can't even prove there is a model of ZFC)! So instead we could only prove the following:

$$\text{Cons}(\text{ZFC}) \rightarrow \text{Cons}(\text{ZFC} + \text{CH})$$

or equivalently, if $M \models ZFC$, then there is $N \models ZFC + CH$ (and similar for the other half).

1 Model theory of set theory

1.1 Absoluteness

For a moment, we will assume that we have a model $(M, \in) \models ZFC$. Unfortunately this first assumption doesn't make much sense, because model theory is based on set theory and we don't have anything if ZFC is inconsistent. We refer to the canonical objects in M by the usual symbols, e.g. $0, 1, 2, 3, 4, \dots, \omega, \omega+1, \dots$

What would an *inner model* be? Take $A \subseteq M$, and consider (A, \in) . It is a substructure of (M, \in) , because there are no function symbols or constant symbols in the language of set theory. This might be counterintuitive, because we're using symbols like ϕ and $\{\cdot\}$ all the time! However, these are technically not part of language of set theory as they can all be defined without any use of function symbols, i.e. they are just abbreviations. For example, $X = \phi$ abbreviates $\forall w(\neg w \in X)$; $X = \{Y\}$ abbreviates $\forall w(w \in X \iff w = Y)$, and similarly for \cup and \mathcal{P} ; and also for relation symbols such as \subseteq , which abbreviates $\forall w(w \in X \rightarrow w \in Y)$. Note that $X = \phi$ is NOT the formula that it looks like; in particular, it is not quantifier free (because it abbreviates $\forall w(\neg w \in X)$)! So we need to take extra care when we do things in this course.

Definition. If φ is a formula in n free variables, we say φ is *upwards absolute* between A and M if for all $a_1, \dots, a_n \in A$,

$$(A, \in) \models \varphi(a_1, \dots, a_n) \implies (M, \in) \models \varphi(a_1, \dots, a_n)$$

and we say φ is *downwards absolute* between A and M if for all $a_1, \dots, a_n \in A$,

$$(M, \in) \models \varphi(a_1, \dots, a_n) \implies (A, \in) \models \varphi(a_1, \dots, a_n)$$

and φ is *absolute* between A and M if it is both upwards and downwards absolute.

Observation:

(a) If φ is *quantifier-free*, then φ is absolute between A and M . But this doesn't really help much, because almost nothing is quantifier-free: without quantifiers we can only say things like $A \in B$ and $A = B$, and conjunctions of those; that's pretty much all.

(b) We say that a formula is Σ_1 if it is of the form

$$\exists x_1 \dots \exists x_n \varphi(x_1, \dots, x_n)$$

where φ is q.f.;

we say a formula is Π_1 if it is of the form

$$\forall x_1 \dots \forall x_n \varphi(x_1, \dots, x_n)$$

where φ is q.f..

(c) If φ is Π_1 , it is downward absolute; if it's Σ_1 then it is upwards absolute. So in particular, note that $X = \phi$ is downward absolute.

As an example, write $0, 1, 2, 3, \dots$ for the ordinals in M , and let $A := M \setminus \{1\}$. In A , we have $0, 2$, but no 1 ; we also have $\{1\}$. If we use $\Phi_0(x)$ to denote the formula $\forall w(\neg w \in x) \iff x = \phi$. Clearly $(M, \in) \models \Phi_0(0)$, so by π_1 -downwards absoluteness, $(A, \in) \models \Phi_0(0)$.

Now, how many elements does $2 = \{0, 1\}$ have? In M we obviously know 2 has 2 elements; but in A , 2 only has one element 0 , and $\{1\}$ has no element: $(A, \in) \models \Phi_0(\{1\})$! Clearly $(M, \in) \not\models \Phi_0(\{1\})$, so Φ_0 is not absolute between A and M . As a corollary, we get $(A, \in) \not\models$ extensionality (we can uniquely specify sets by specifying their elements).

Remark. We could go on, defining formulas $\Phi_1(x), \Phi_2(x)$, etc to analyse which of the elements correspond to the natural numbers in A .

Reminder (from Part II Logic and Set Theory): we say A is *transitive in M* if for all $a \in A$ and $x \in M$ s.t. $(M, \in) \models x \in a$, we have $x \in A$. The problem for the above A is that it is not transitive. As long as that is fixed, we have the following:

Proposition. If A is transitive, then Φ_0 is absolute between A and M .

Proof. Since Φ_0 is Π_1 , we only need to show upwards absoluteness. Suppose $a \in A$ s.t. $(A, \in) \models \Phi_0(a)$, and suppose for contradiction that $a \neq 0$. Then there is some $x \in a$. By transitivity, $x \in A$. But then $\Phi_0(a) : \forall w(w \notin a)$ is not true in (A, \in) . \square

Similarly, if Φ_n is the formula describing the natural number n , and there is $a \in A$ s.t. $(A, \in) \models \Phi_n(a)$, and A is transitive, then $a = n$.

Proposition. If A is transitive in M , then $(A, \in) \models$ extensionality.

Proof. Take $a, b \in A$ with $a \neq b$. So by extensionality in (M, \in) , find, WLOG some $c \in a \setminus b$. Since $c \in a \in A$, by transitivity $c \in A$. Note that all of these quantifier-free formulas are absolute, so (A, \in) also models them; in particular, $(A, \in) \models c \in a, c \notin b$. So a, b do not satisfy the assumptions of extensionality. \square

Consider now $A = \omega + 2 = \{0, 1, 2, \dots, \omega, \omega + 1\} \subseteq M$. This is clearly transitive subset of M because it's an ordinal. So $(A, \in) \models$ extensionality, but clearly it isn't anything like a model of set theory as it is too thin. Consider the formula $x = \mathcal{P}(y)$. Unfortunately, this is not a formula, as \mathcal{P} is undefined. We have to expand it properly:

$$\begin{aligned} x &= \mathcal{P}(y) \\ \iff x &= \{z; z \subseteq y\} \\ \iff \forall w(w \in x &\leftrightarrow w \subseteq y) \\ \iff \forall w(w \in x &\leftrightarrow (\forall v(v \in w \rightarrow v \in y))) \end{aligned}$$

In A , what is $\mathcal{P}(\omega)$? We have $(A, \in) \models \omega + 1 = \mathcal{P}(\omega)$, which is obviously not what we want for $\mathcal{P}(\omega)$ to be.

Definition. (Bounded quantification)

We first define the notations $\exists v \in w \varphi$ to be $\exists v(v \in w \wedge \varphi)$, and $\forall v \in w \varphi$ to be $\forall v(v \in w \rightarrow \varphi)$, and we call these quantifiers *bounded*.

Now we say a formula φ is Δ_0 if it is in the smallest set S of formulas with the following properties:

1. All q-f formulas are in S ;
2. If $\varphi, \psi \in S$, then so are:

- 2a. $\varphi \wedge \psi, \varphi \vee \psi, \varphi \rightarrow \psi, \varphi \leftrightarrow \psi$;
- 2b. $\neg\varphi$;
- 2c. $\exists x \in w \varphi, \forall v \in w \varphi$.

Theorem. If φ is Δ_0 and A is transitive, then φ is absolute between A and M .

Proof. We already know that quantifier free formulas are absolute, and absoluteness is obviously preserved under propositional connectives. The only case left is (2c).

Let's just do $\varphi \rightarrow \exists v \in w \varphi = \exists v(v \in w \wedge \varphi)$. So suppose φ is absolute. We need to deal with downwards absoluteness: we have $(M, \in) \models \exists v \in a \varphi(v, a)$ for some $a \in A$, i.e. $(M, \in) \models \exists v(v \in a \wedge (\varphi(v, a)))$.

Let's find $m \in M$ s.t. $(M, \in) \models m \in a \wedge \varphi(m, a)$.

Now $m \in a \in A$, so $m \in A$. By absoluteness of φ , we get $(A, \in) \models m \in a \wedge \varphi(m, a) \implies (A, \in) \models \exists v \in a \varphi(v, a)$. \square

Let T be any *set theory*. Then we say that φ is Δ_0^T if there is a Δ_0 formula ψ s.t. $T \vdash \varphi \leftrightarrow \psi$. So we get, as a corollary:

Corollary. If A is transitive in M , and both (M, \in) and (A, \in) are models of T , then Δ_0^T formulas are absolute between A and M .

We may also define Σ_1^T formulas to be the formulas that are T -equivalent to $\exists v_1 \dots \exists v_n \psi$ where ψ is Δ_0 , and similarly for Π_1^T formulas. So $\Sigma_1^T(\Pi_1^T)$ formulas are upwards(downwards) absolute between A and M respectively.

On Wednesday we will look at what formulas are actually in these classes.

—Lecture 3—

Last time we fixed some *set theory* T , and defined formula classes Δ_0^T , Σ_0^T and Π_1^T . We showed that Δ_0^T formulas are absolute between A, M if A is transitive and $A, M \models T$, and also Σ_1^T and Π_1^T upwards and downwards respectively.

Even if you haven't paid attention you would have realize that we have some 0 and some 1 as subscripts here. So what is Δ_1^T ?

Definition. A formula is Δ_1^T if it is both Σ_1^T and Π_1^T .

Note that this definition is only possible upon taking equivalence classes on T , else no formula could be both Σ_1 and Π_1 .

Corollary. If A is transitive, $A, M \models T$, and ϕ is Δ_1^T , then ϕ is absolute between A and M .

Now we have to think of what a *set theory* is. We have to think of which axioms we're using. Preferably we would have extensionality, and then let's have pairing, union, power set, separation.

We denote this by FST_0 (finite set theory), with the 0 denoting that we don't have foundation yet. We use FST to denote FST_0 +foundation(regularity).

Now if we add infinity in, we reach the original version of Zermelo set theory Z_0 . However, nowadays we often call $Z = Z_0$ +foundation the Zermelo set theory.

For ordinary people these are enough (or far more than enough). But set-theorists realized later that they need replacement; we call this ZF_0 (of course ZF for the version with foundation). And lastly if we add choice in we get ZFC_0 (with foundation we get ZFC).

1.2 Long List of Δ_0^T formulas

Now we find a long list of Δ_0^T formulas. We start with the more trivial ones:

1. $x \in y$;
2. $x = y$;

These two are Δ_0 without T needed.

3. $x \subseteq y$. Apparently this is not a formula: we think it means $\forall w(w \in x \rightarrow w \in y)$, which we might abbreviate it as $\forall w \in x(w \in y)$, which is exactly the (2c) in definition of Δ_0 . So this is Δ_0 without T as well.

4. $\Phi_0(x) : \forall w(w \notin x) : \iff \forall w(\neg w \in x)$. If you took part II Logic and Set theory, you'll disagree that this is a formula, because \neg is not a thing; but let's not be so parsimonious on the syntax, but write it as $\forall w(w \in x \rightarrow \neg x = x)$, so this is also Δ_0 in predicate logic.

We say that an operation $x_1, \dots, x_n \rightarrow F(x_1, \dots, x_n)$ is defined by a formula in class Γ (where Γ is any class of formulas) in the theory T if there is a formula $\Psi \in \Gamma$ s.t.

- (1) $T \vdash \forall x_1 \dots \forall x_n \exists y \Psi(x_1, \dots, x_n, y)$;
- (2) $T \vdash \forall x_1 \dots \forall x_n \forall y, z \Psi(x_1, \dots, x_n, y) \wedge \Psi(x_1, \dots, x_n, z) \rightarrow y = z$;
- (3) $\Psi(x_1, \dots, x_n, y)$ iff $y = F(x_1, \dots, x_n)$.

Note that the first two are formal requirements, but the last one is an informal requirement as we haven't defined what F is.

Examples: $x \rightarrow \{x\}$, $x, y \rightarrow \{x, y\}$ (these are operations in FST_0). Note that these are informal because notations like $\{\cdot\}$ are undefined.

Let's now continue our lists:

5. $x \rightarrow \{x\}$. We need a formula $\Psi(x, z) \leftrightarrow 'z = \{x\}' \leftrightarrow \forall w(w \in z \leftrightarrow w = x)$.

This is not Δ_0 yet because we have a \leftrightarrow here. We rewrite it as $\forall w((w \in z \rightarrow w = x) \wedge (w = x \rightarrow w \in z))$, but the second part is not Δ_0 . So we again rewrite it as

$\exists w \in z(w = x) \wedge \forall w \in z(w \in z \rightarrow w = x)$. So this is Δ_0 , with some very weak set theory being sufficient.

Similar to 5, we also have

6. $x, y \rightarrow \{x, y\}$;
7. $x, y \rightarrow x \cup y$;
8. $x, y \rightarrow x \cap y$;

9. $x, y \rightarrow x \setminus y$;

10. $x, y \rightarrow (x, y)$, the ordered pair, where we define it as $\{\{x\}, \{x, y\}\}$. Note that we could apply 5 and 6 (twice) to get this one.

The last one gives us the motivation that if two operations f, g_1, \dots, g_k are defined by Δ_0^T -formulas, then so is the operation

$$x_1, \dots, x_n \rightarrow f(g_1(x_1, \dots, x_n), \dots, g_k(x_1, \dots, x_n))$$

We then naturally have

11. $x \rightarrow x \cup \{x\} := S(x)$ (by the previous fact from 5 and 7).

12. $x \rightarrow \cup x$; (obvious if we write this fully out)

13. the formula φ describing " x is transitive".

14. the formula describing x is an ordered pair. At first look it looks like this is unbounded, but that's not the case: the quantifiers for the two components of x are bounded by $\cup x$.

15. $a, b \rightarrow a \times b$;

16. the formula " x is a binary relation";

17. $x \rightarrow \text{dom}(x) := \{y : \exists p \in x (p \text{ is an ordered pair, } p = (v, w), y = v)\}$;

18. $x \rightarrow \text{range}(x) := \{y : \exists p \in x (p \text{ is an ordered pair and } p = (v, y))\}$;

19. the formula ' x is a function';

20. the formula ' x is injective';

21. the formula ' x is a function from A to B ';

22. the formula ' x is a surjection from A to B ';

23. the formula ' x is a bijection from A to B '.

Note that we've only used some very few axioms: union, pairing and some finite version of separation, and nothing more.

Let's also agree on the definition of an ordinal: α is an ordinal if α is a transitive set well-ordered by \in . Of course we have to also agree on what being well-ordered means: it's totally ordered + well-founded, i.e. $\forall X (X \subseteq \alpha \rightarrow X \text{ has a } \in\text{-least element})$.

Being totally ordered is Δ_0 formula (check); however, the sentence (X, R) being well-founded is not obviously absolute, since the bound for the $\forall Z (Z \subseteq X \dots)$ quantifier is the power set. We'll talk about absoluteness of well-foundedness on Friday.

However, we don't actually need the general well-foundedness; we only need well-foundedness by \in , but that is given by axiom of foundation! So in models with the axiom of foundation, α is an ordinal iff α is transitive and totally ordered by \in .

—Lecture 4—

We're still on our list of things that are absolute for transitive models. We ended with ordinals last time, where we defined that x is an ordinal iff x is transitive and (x, \in) is a well-order. We went into an issue there, because that consists of (x, \in) is a total order, which is fine; but then it also needs \in is a well-founded relation on x , which is only Π_1 . The good side is that if T contains the axiom

of foundation, then $\Phi_{ord}(x)$ is equivalent to x is transitive and (x, \in) is a total order (as the last part is guaranteed), which is Δ_0^T . Therefore we can expand our list:

- 24. ' x is an ordinal' is Δ_0^T (for the right choice of T);
This is not as harmless as before, because we actually need T to include the axiom of foundation.
- 25. ' x is a successor ordinal', which is equivalent to ' x is an ordinal' and $\exists y \in x (y$ is the \in -largest element of x);
- 26. ' x is a limit ordinal';
- 27. $x = \omega$ (the smallest limit ordinal; similarly, $x = \omega + \omega$, $x = \omega + 1$, $x = \omega + \omega + 1$, $x = \omega^2$, $x = \omega^3$, $x = \omega^\omega$, ...)

1.3 Absoluteness of well-foundedness

If (X, R) is well-founded, we can define a rank function $rk : X \rightarrow \alpha$, where α is some ordinal, s.t. rk is order-preserving between (X, R) and (α, \in) . This theorem is proved using the right instances of Replacement. In particular, ZF proves:

(X, R) is well-founded $\iff \exists \alpha \exists f$ α is an ordinal, and f is an order-preserving function from (X, R) to (α, \in) . RHS is Σ_1^{ZF} , and LHS is Π_1^{ZF} . Thus for sufficiently strong T , (X, R) is well-founded is Δ_1^T and hence absolute for transitive models of T .

We generalize this to concepts defined by transfinite recursion. But first let's recall what it is: let (X, R) be well-founded, let F be 'functional', so for every x there is a unique y s.t. $x = F(y)$. Then there is a unique f with domain X and for all $x \in X$, $f(x) = F(f|IS_R(x))$, where $IS_R(x) = \{z \in x : zRx\}$.

Proposition. Let T be a set theory that is strong enough to prove the transfinite recursion theorem for F . Let F be absolute for transfinite models of T . Let (X, R) be in A . Then f defined by transfinite recursion is absolute between A and M .

Example. Let L be any first-order language whose symbols are all in A . Then the set of L -formulas and the set of L -sentences are in A .

The relation $S \models \varphi$ (note that this is not q-f, although it is bounded by S) is defined by recursion, and thus is absolute between A and M .

So: if S is an L -structure, $S \in A$,

$(A, \in) \models 'S \models \varphi' \iff (M, \in) \models 'S \models \varphi'$.

Gödel's incompleteness theorem roughly says that, if T is a theory whose set of axioms are recursive enumerable, and its axioms are strong enough to do some arithmetics, then $T \not\models Cons(T)$ (which is a sentence in L). Examples for T : PA, Z, ZF, ZFC, ZFC+ φ any one additional formula.

In particular, $ZFC^* := ZFC + Cons(ZFC) \not\models Cons(ZFC + Cons(ZFC))$.

By completeness theorem, $Cons(T) \iff \exists M (M \models T)$. Note that LHS is Π_1 and RHS is Σ_1 , so completeness theorem tells us that this is a Δ_1 concept. Note

also that LHS is a bounded concept, since all quantifiers are bounded (LHS is Δ_0^Z).

Now write β for 'there is a transitive set A s.t. $(A, \in) \models ZFC$ '. Note that this is stronger than the consistency of ZFC as it also specifies that there is a particular model for it. In particular, we can't prove it even in ZFC^* :

Theorem. If ZFC^* is consistent, then $ZFC^* \not\vdash \beta$.

Proof. Let $(M, \in) \models ZFC^*$. Suppose $ZFC^* \vdash \beta$. So $(M, \in) \models \beta$. So we found a transitive set A in M s.t. (A, \in) is a model of ZFC. By assumption, $(M, \in) \models \text{Cons}(ZFC)$, so $(A, \in) \models \text{Cons}(ZFC)$ since $\text{Cons}(ZFC)$ is absolute between transitive models. So $(A, \in) \models ZFC^*$. So we proved $\text{Cons}(ZFC^*)$, contradicting Gödel's incompleteness theorem. \square

That means that assuming β is not an obvious assumption, so we need to study under which (natural) assumptions β is true.

1.4 Concrete transitive models of ZFC

So let's investigate transitive models A inside M . The two most basic constructions:

- (1) von Neumann hierarchy (cumulative hierarchy);
- (2) hereditarily small sets.

(1) is defined as follows by transfinite recursion: $V_0 := \phi$, $V_{\alpha+1} := \mathcal{P}(V_\alpha)$, $V_\lambda := \bigcup_{\alpha < \lambda} V_\alpha$ for λ being limit ordinals.

Proposition. $\forall \alpha V_\alpha$ is transitive (see Part II Logic and Set Theory). [Induction, with key lemma: if X is transitive, then $\mathcal{P}(X)$ is transitive]

If λ is a limit ordinal, then $V_\lambda \models FST$.

If $\lambda > \omega$ and a limit ordinal, then $V_\lambda \models Z$.

For (2): let κ be a cardinal. We say X is hereditarily smaller than κ if $|tcl(X)| < \kappa$ (transitive closure).

Let $H_\kappa := \{X : X \text{ is hereditarily smaller than } \kappa\}$. This is obviously transitive.

We'll continue next Monday.

—Lecture 5—

We'll start with V_α today.

We know (if you don't know then I invite you to learn it): if λ is a limit then $V_\lambda \models FST$; if further $\lambda > \omega$, then $V_\lambda \models Z$. (On example sheet 1. I haven't printed them today, so if you want to try them before Wednesday, find an envelope next to my office at C0.10 from this afternoon).

The critical axiom here is *Replacement*. The test case is $\lambda = \omega + \omega$, which we expect replacement fails. Remember replacement says that if we have a function $F : V_{\omega+\omega} \rightarrow V_{\omega+\omega}$ that is definable in $V_{\omega+\omega}$, and $x \in V_{\omega+\omega}$, then $\{F(y) : y \in x\} \in V_{\omega+\omega}$, i.e. image of elements of a set under a function is a set.

The idea is to take $x = \omega$, and define F to be the function that takes $n \rightarrow \omega + n$, and $y \rightarrow 0$ if $y \notin \omega$. Remember that definable just means that we need a formula that uniques specify the image (see previous notes). Let $Y = \{F(n) : n \in \omega\}$, which needs to be a set if replacement holds. Then Y is a subset of $V_{\omega+\omega}$, so $Y \in V_{\omega+\omega+1}$; but it is not bounded, so it is not in $V_{\omega+\omega}$. This example shows concretely that $V_{\omega+\omega} \models \neg \text{Replacement}$.

What we needed in this particular example is a function that takes a bounded sequence to an unbounded one. Similarly, if α is any ordinal s.t. there is a definable function $f : \omega \rightarrow \alpha$, s.t. the range of f is unbounded in α , then $V_\alpha \models \neg \text{Replacement}$.

Even more general, if $\beta < \alpha$, and a definable function $f : \beta \rightarrow \alpha$ with unbounded range, then $V_\alpha \models \neg \text{Replacement}$.

Reminder: we call a cardinal κ *regular* if there is no partition $\kappa = \bigcup_{i \in I} A_i$ s.t. $|I|, |A_i| < \kappa$ for all $i \in I$.

Equivalently, for every $\alpha < \kappa$, there is no unbounded function $f : \alpha \rightarrow \kappa$.

We know, for example, that \aleph_1 is regular. Moreover, for any α , $\aleph_{\alpha+1}$ is regular. So this gives us the next candidate, $\alpha = \aleph_1$, which replacement cannot fail in the above way. So how does it fail? Note that $\mathcal{P}(\omega) \in V_{\omega+2} \subseteq V_{\omega_1}$. Clearly, there is a surjection $s : \mathcal{P}(\omega) \rightarrow \omega_1$. But that means its range is unbounded in ω_1 . Thus $V_{\omega_1} \models \neg \text{Replacement}$ either.

In general, if κ is regular and there is $\lambda < \kappa$ with $|\mathcal{P}(\lambda)| \geq \kappa$, then the same argument shows $V_\kappa \models \neg \text{Replacement}$.

Definition. A cardinal κ is called *inaccessible* if

- (a) κ is regular;
- (b) $\forall \lambda < \kappa, |\mathcal{P}(\lambda)| < \kappa$.

Side note: Related to the question, 'are there regular limit cardinals'? Under GCH: $\forall \kappa, 2^\kappa = \kappa^+$, we have that κ is inaccessible iff κ is regular limit. This suggests that the above question cannot be answered that easily (we usually call regular limit cardinals *weakly inaccessible*).

Let's assume that $\kappa > \omega$ is inaccessible (because ω is actually inaccessible, which kind of make sense, because *infinity* is the ultimate thing that is not accessible from everything smaller than it).

Lemma. $\forall \lambda < \kappa, |V_\lambda| < \kappa$.

Proof. Clearly $|V_\omega| = \aleph_0$, so $|V_\omega| < \kappa$.

By induction, suppose $|V_\lambda| < \kappa$. Then $V_{\lambda+1} = \mathcal{P}(V_\lambda)$. Therefore $|V_{\lambda+1}| = |\mathcal{P}(V_\lambda)| < \kappa$ by (b).

Now let $\lambda < \kappa$ be a limit ordinal. Then $V_\lambda = \bigcup_{\alpha < \lambda} V_\alpha$. So suppose by contradiction that $|V_\lambda| = \kappa$. But $|V_\alpha| < \kappa$ for all $\alpha < \kappa$, so we can write κ as a union of λ many things of smaller cardinals, contradicting regularity. \square

Theorem. If κ is inaccessible, then $V_\kappa \models \text{Replacement}$.

Proof. We're actually going to something slightly stronger: take any function $F : V_\kappa \rightarrow V_\kappa$ (without caring of its definability), and any $x \in V_\kappa$. Note that $V_\kappa = \bigcup_{\alpha < \kappa} V_\alpha$. So we can find $\alpha \in \kappa$ s.t. $x \in V_\alpha$. Since V_α is transitive, $x \subseteq V_\alpha$. But that means $|x| \leq |V_\alpha| < \kappa$ (by lemma). Note that this is exactly what went wrong in V_{ω_1} .

Now consider $X := \{F(y) : y \in x\}$. For each $y \in x$, consider $\rho(F(y)) :=$ the least α s.t. $F(y) \in V_{\alpha+1} \setminus V_\alpha$. By assumption, $\rho(F(y)) < \kappa$. Consider $\{\rho(F(y)) : y \in x\} := R$, then $|R| \leq |x| < \kappa$. By regularity, $\alpha := \bigcup R < \kappa$. But now $\forall y \in x F(y) \in V_{\alpha+1}$. So $X \subseteq V_{\alpha+1}$, so $X \in V_{\alpha+2}$, i.e. $V_\kappa \models \text{Replacement}$. \square

Note that this proves that the existence of inaccessible cardinals cannot be proved from ZFC.

1.5 Inaccessible cardinals

—Lecture 6—

Course webpage:

https://www.math.uni-hamburg.de/home/loewe/Lent2019/TST_L19.html.

Hand in work at the start of example class.

On the example sheet we've seen(or will see) that we write IC for the axiom 'there is an inaccessible cardinal.

We've seen that if κ is inaccessible, then $V_\kappa \models \text{ZFC}$ (which is a transitive model). So $\text{ZFC} + \text{IC} \vdash$ 'there is a transitive set that is a model of ZFC', which we have called this β at some point; we've also proved that β is stronger than $\text{Cons}(\text{ZFC})$, in the sense that $\text{ZFC} + \text{Cons}(\text{ZFC}) \not\models \beta$. Therefore $\text{ZFC} + \text{Cons}(\text{ZFC}) \not\models \text{IC}$.

Two model-theoretic reminders (see Part II Logic and Set Theory for both):

(1) Löwenheim-Skolem theorem. We want to formalize this: if \mathcal{S} is any structure in some countable first-order language L , and $X \subseteq S$ is any subset, then there is a *Skolem hull* of X in \mathcal{S} , usually written as $\mathcal{H}^{\mathcal{S}}(X)$, with $X \subseteq \mathcal{H}^{\mathcal{S}}(X) \subseteq \mathcal{S}$, s.t.

(a) $\mathcal{H}^{\mathcal{S}}(X) \prec \mathcal{S}$ (elementary substructure of \mathcal{S});

(b) $|\mathcal{H}^{\mathcal{S}}(X)| \leq \max(\aleph_0, |X|)$.

You probably have seen the proof of this theorem, but let's have a sketch again because it's important to see how theorems of this kind are proved.

Proof. (sketch)

Key ingredient – Tarski-Vaught criterion: $Z \subseteq S$ then $Z \prec S$ iff for every φ and all z_1, \dots, z_n , if $S \models \exists x \varphi(x, z_1, \dots, z_n)$, then $Z \models \exists x \varphi(x, z_1, \dots, z_n)$.

In order to construct the Skolem hull, we define $Z_1 = Z_0 \cup$ witnesses for all

tuples $\varphi_1(z_1, \dots, z_n)$ where $z_1, \dots, z_n \in Z_0 = X$, and then $Z_{n+1} := Z_n \cup \dots$. By induction, each Z_i has the same size. In the end we define $Z := \bigcup_{n \in \mathbb{N}} Z_n$, which will satisfy the Tarski-Vaught criterion. \square

Consequence: work in $ZFC + IC$, $(M, \in) \models ZFC + IC$; inside $V_\kappa \subseteq M \models ZFC$. Apply L-S theorem to V_κ with $X := \phi$, we find a structure $H := \mathcal{H}^{V_\kappa}(\phi) \prec V_\kappa$, which is countable (since it cannot be finite). So we've found a countable model of ZFC – isn't this a contradiction? Not really, because this H is, in general, not transitive – and it probably shouldn't be – consider the sentence ' $\exists x$ s.t. x is the least uncountable cardinal. This is certainly true in V_κ , so it needs to be also true in H ; but its only witness in V_κ is \aleph_1 , so $\aleph_1 \in Z_1 \subseteq H$; so H cannot be transitive since \aleph_1 has uncountably many elements.

(2) Mostowski Collapse Theorem. If X is any set, and $R \subseteq X \times X$ s.t. R is well-founded and extensional, then there is a transitive set T s.t. $(T, \in) \cong (X, R)$. Consider $(H, \in) \models ZFC$: \in is extensional on H by the axioms of ZFC; \in is well-founded on M , so \in is well-founded on H . So, let T be the Mostowski collapse of H : T is transitive, $(T, \in) \cong (H, \in)$. So $(T, \in) \models ZFC$, and now T is transitive, and since this is a bijection, $|T| = |H| \leq \aleph_0$, so we get a countable transitive model of ZFC!

(some diagrams, where we used T to denote the countable transitive model of ZFC)

Consider $\varphi(x) := x$ is countable: $\exists f(\underbrace{f : x \rightarrow \mathbb{N}}_{\Delta_0^{ZFC}}, \underbrace{f \text{ is injective}}_{\Delta_0^{ZFC}})$, which is Σ_1^{ZFC} ,

so is upwards absolute. But this formula is *not* downwards absolute: if α is an ordinal, $\alpha \in T$, then $V_\kappa \models \alpha$ is countable. But since $(T, \in) \models ZFC$, there is some $\alpha \in T$ s.t. $(T, \in) \models \alpha$ is uncountable, so V_κ and T disagree about the truth value of $\varphi(\alpha)$.

Now consider $\psi(x) : x$ is a cardinal: $\forall \alpha(\alpha < x \rightarrow \text{there is no injection from } x \text{ to } \alpha)$, which is Π_1^{ZFC} , so downwards absolute. In (T, \in) , take α least s.t. $(T, \in) \models \neg \varphi(\alpha)$. Then $(T, \in) \models \alpha$ is a cardinal. Clearly $V_\kappa \models \alpha$ is not a cardinal. So ψ is not upwards absolute.

Note that if λ is an uncountable cardinal in V_κ , then $\lambda \notin T$, so the downwards absoluteness of ψ is not very interesting because there aren't many cardinals.

Instead of building $\mathcal{H}^{V_\kappa}(\phi)$, why not build $H^* := \mathcal{H}^{V_\kappa}(\omega_1 + 1)$? Clearly $\omega_1 \in H$ now, and $\omega_1 \subseteq H$ as well, so ω_1 stays the same when we do Mostowski's Collapse; so $\omega_1 \subseteq T^*$ and $\omega_1 \in T^*$ where T^* is the new model obtained after Mostowski's collapse.

Now we have $V_\kappa \models \omega_1$ is a cardinal, so by downwards absoluteness, $T^* \models \omega_1$ is a cardinal. However, it might be the case that there are some other cardinals below ω_1 , so we can't say $\omega_1 = \aleph_1$.

Remember our goal is to deduce some results about CH. We've decided to go for transitive models of ZFC (as we don't want to be concerned about whether a function is still a function in a submodel). We've looked at 'inner models', and in particular, models of type V_α . We also know that if α is inaccessible, then $V_\alpha \models ZFC$. We've also found countable transitive submodel in V_α of ZFC, called T , by Mostowski collapse and L-S theorem.

One problem: this is *not* going to change the truth value of CH: Suppose CH is true in (M, \in) , so there's a bijection between \mathbb{R} and ω_1 . Similarly, if CH is false then there is no such bijection. An immediate problem is that we don't know what \mathbb{R} is; we might as well replace it by $\mathbb{P}(\mathbb{N})$, which is definitely in $V_{\omega+20}$ (safe enough), while $\omega_1 \in V_{\omega_1+1}$. But that means V_κ knows exactly if there is a bijection (say we can define this in $V_{\omega_1+20} \subseteq V_\kappa$) between \mathbb{R} and ω_1 because it has access to both of the levels, and it could just 'take a look' at it to determine the truth value. So $(M, \in) \models CH \iff (V_\kappa, \in) \models CH$.

But remember by L-S theorem we got a countable $H \prec V_\kappa$, so $(H, \in) \models CH \iff (V_\kappa, \in) \models CH$; and Mostowski collapse gives $(T, \in) \cong (H, \in)$, so $(T, \in) \models CH \iff (H, \in) \models CH$.

Summary: the method of finding countable transitive *elementary* submodels of V_κ is not going to change the value of CH. So let's look at different models.

1.6 The second construction: Models of hereditarily small sets

Let κ be a regular cardinal (e.g. $\kappa = \omega, \kappa = \omega_1$). Then x is called hereditarily of size $< \kappa$ if $|tcl(x)| < \kappa$, where $tcl(x)$ is the transitive closure of $x (= \bigcup_{n \in \mathbb{N}} t_n(x))$, where $t_0(x) := x$, $t_{n+1}(x) = \bigcup t_n(x)$.

The definition of $tcl(x)$ captures the intuition of ' x has size $< \kappa$, all elements of x have size $< \kappa$, all elements of elements of x have size $< \kappa$, etc.'

Remark. It's important that κ is regular for the intuition to work: suppose it's not, say let $\kappa = \aleph_\omega$. Now think of a tree that has branches of all finite lengths, where at level n we attach something of cardinality \aleph_n at the node. More formally, define

$$\begin{aligned} x_n^0 &= \aleph_n \\ x_n^{k+1} &= \{x_n^k\} \\ &\dots \\ \mathbb{X} &:= \{x_0^0, x_1^1, x_2^2, \dots\} \end{aligned}$$

Note that \mathbb{X} is countable, but in the 'tcl' notation above, the cardinality of $t_{n+1}(\mathbb{X})$ is \aleph_n , so $tcl(x)$, as the union of them, has cardinality \aleph_ω ! The problem is obviously because we could obtain \aleph_ω from fewer than \aleph_ω of smaller than \aleph_ω things, i.e. \aleph_ω is not regular.

From now let's forget about those singular κ , but only consider κ regular.

We make a few observations:

1. H_κ is transitive;
2. If $X \subseteq H_\kappa$ and $|X| < \kappa$, then $X \in H_\kappa$ (follows directly from regularity of κ and the definitions).

Example. Let $H_{\aleph_0} := HF$ (hereditarily finite).

We claim that $HF = V_\omega$.

Proof. $V_\omega \subseteq HF$. We need to show $V_n \subseteq HF \forall n$. Clearly $V_0 = \phi \subseteq HF$; if $V_n \subseteq HF$, and $Z \subseteq V_n$, then by observation 2, $Z \in HF$, so $\mathcal{P}(V_n) = V_{n+1} \subseteq HF$. Now we show $HF \subseteq V_\omega$. Suppose not, so there is $x \in HF \setminus V_\omega$; take such x with minimal rank α , so $x \in V_{\alpha+1} \setminus V_\alpha$. By minimality, if $y \in HF$ with $\rho(y) < \alpha$, then there is $k \in \mathbb{N}$ s.t. $\rho(y) = k$. We know $x \in \mathcal{P}(V_\alpha)$, so $x \subseteq V_\alpha \subseteq HF$. So $x \in HF$, so x is finite. Say $x = \{x_1, \dots, x_n\}$. By minimality, each of them is in V_{k_i} for some k_i , so $x \subseteq V_{\max(k_1, \dots, k_n)+1} \subseteq V_\omega$. \square

Example. $H_{\aleph_1} = HC$ (hereditarily countable).

We know:

- $Ord \cap HC = \omega_1$;
- $V_{\omega+2} \setminus HC \neq \phi$;
- $V_{\omega+1} \subseteq HC$.

Which axioms are true in HC? Let's check Pair: say $x, y \in HC$. then $\{x, y\} \subseteq HC$. But $|\{x, y\}| < \aleph_1$, so $\{x, y\} \in HC$.

Separation, foundation, extensionality, union all hold in HC as well quite easily. Replacement: let F be a function $HC \rightarrow HC$, and $x \in HC$. Consider $R := \{F(y); y \in x\}$, which exists in the universe, and we need to check whether it's in R . We know $|R| \leq |x| < \aleph_1$, and $R \subseteq HC$. By observation 2 we immediately get $R \in HC$. So replacement is trivial!

So we know Power Set must be false, or at least not provable in HC, else we've proved the existence of a model of ZFC!

Note that Power Set is trivial in ZFC and it's replacement that is troublesome there, while here it's the other way round.

We know that $\mathbb{N} \in HC$, and we also know that $\mathcal{P}(\mathbb{N}) \notin HC$. But that is not enough to disprove Power Set in HC; we need to show that there is no object in HC such that HC *thinks* that it is the power set of \mathbb{N} ; more formally, we need to show that for all $A \in HC$, $HC \models A$ is not a power set of \mathbb{N} , i.e. $\exists X(X \subseteq \mathbb{N} \wedge X \notin A)$.

Fix $A \in HC$, and presume that this might be the HC-powerset of \mathbb{N} .

Thus, if $X \subseteq \mathbb{N}$ and $X \in HC$, then $X \in A$. But if $X \subseteq \mathbb{N}$, then $X \subseteq HC$, and $|X| < \aleph_1$, so $X \in HC$.

So if A is any set s.t. $\forall X, X \subseteq \mathbb{N} \wedge X \in HC \rightarrow X \in A$, then A is uncountable, so $A \notin HC$. Contradiction. Thus $HC \models \neg PowerSet$.

—Lecture 8—

We've not been doing very useful things so far since what we've done was basically proving that many things are not useful. Even worse, the thing that we're going

to do now is the most pointless of all as it's almost identical to some problems in the example sheet.

We knew H_κ is a model of all of ZFC without power set if κ is regular.

Proposition. If κ is a strong limit (cannot be reached by taking power set), then $H_\kappa \models \text{PowerSet}$.

Proof. We'll show: if $x \in H_\kappa$, then $\mathcal{P}(x) \in H_\kappa$. This is much stronger than the Power Set axiom.

Certainly $\text{tcl}(\mathcal{P}(x)) = \mathcal{P}(x) \cup \text{tcl}(x)$. Now $|\text{tcl}(x)| < \kappa$ since $x \in H_\kappa$, and $|\mathcal{P}(x)| < \kappa$ as well since κ is a strong limit. So together we know $|\text{tcl}(\mathcal{P}(x))| < \kappa$, so $\mathcal{P}(x) \in H_\kappa$. \square

General idea: build inner models using *definability* properties. But the problem is: definability is not definable. This sounds like a typical joke from logicians, but it's actually a theorem.

Theorem. (Tarski Undefinability of Truth)

Let $(M, \in) \models \text{ZFC}$ (we'll just use this, but we'll see in the proof what set theory we actually need). We assume that the language of set theory $L_\in \subseteq M$ (so that sentences and formulas are actually elements of M). Consider the set S of sentences of L_\in and the set U of unary predicates (i.e. L_\in -formulas in one free variable).

A *truth predicate* would be a formula $T(x)$ in L_\in , so a unary predicate, i.e. $T \in U$, s.t. $(M, \in) \models \varphi \iff (M, \in) \models T(\varphi)$.

Claim: there can be no such truth predicate.

Before proving this, let's contrast this with the definition of truth we had before. Our previous result says if M is a set, then the concept $(M, \in) \models \varphi$ is Δ_0 (because we bound the quantifier by M , which is a set, so in some sense we had to use a parameter M in our previous result, which is not available in our current settings).

Proof. (idea: diagonalisation)

If $\varphi(x) \in U$, then we can ask whether $\varphi(\varphi)$ (which is a sentence, so $\in S$) is true. Let's assume that there is a truth predicate T , and define a unary formula $\delta(x) := \neg T(x(x))$ (the *diagonal*), where $x(x)$ is x applied to x if $x \in U$, and is ϕ otherwise (remember formulas are elements in M). Now apply δ to δ , so $\delta(\delta) \in S$. Now

$$M \models \neg T(\delta(\delta)) \iff M \models \delta(\delta) \iff M \models T(\delta(\delta))$$

by definition of δ and the fact that T was a truth predicate respectively. Contradiction. \square

Again, let $M \models \text{ZFC}$. We say that $x \in M$ is *definable* if there is a formula φ s.t. $\forall y \in M \ x = y \iff M \models \varphi(y)$.

We say that a formula D is called a *definition of definability* if $\forall x \in M, x$ is definable $\iff M \models D(x)$. By now we all know what we should expect next:

Theorem. (undefinability of definability)

There is no formula D that is a definition of definability.

Proof. Assume D is a definition of definability. Consider (informally now)

$$\alpha := \max\{\beta : \beta \text{ is not definable, but } \forall \gamma < \beta \exists \gamma' \gamma < \gamma' < \beta \text{ and } \gamma' \text{ is definable}\}$$

which is the supremum of the definable ordinals. This has to exist because there are only countably many formulas, and each formula only specifies at most one ordinal; this has to be uncountable since, say, \aleph_1 is definable. This is defined by the formula:

$y = \alpha \leftrightarrow y$ is an ordinal and $\neg D(y)$ and

$$\forall \gamma (\gamma < y \rightarrow \exists \gamma' (\gamma < \gamma' < y \wedge D(\gamma')))$$

This proves that α is definable in the sense of the definition, so $M \models D(\alpha)$; but one of the conjuncts in the definition implies $M \models \neg D(\alpha)$. Contradiction. \square

We now learned that *definability* is not going to work without keeping track of parameters. So we need to define definability with direct reference to what parameters are allowed.

Fix A , and $n \in \mathbb{N}$. We're going to define by recursion what it means to be a definable subset of A^n :

Define $Diag_{\in}(A, n, i, j) := \{s \in A^n : s_i \in s_j\}$ (read as *diagrams*),

$Diag_{=}(A, n, i, j) := \{s \in A^n : s_i = s_j\}$,

$Proj(A, R, n) := \{s \in A^n : \exists t \in R(t|_n = s)\}$ (where the formula in the last bracket means t restricted to n) (the intended meaning is for $R \subseteq A^{n+1}$). Now define

$Def(0, A, n) := \{Diag_{\in}(A, n, i, j) : i, j < n\} \cup \{Diag_{=}(A, n, i, j) : i, j < n\},$

$$\begin{aligned} Def(k+1, A, n) := & Def(k, A, n) \cup \{R \cap S : R, S \in Def(k, A, n)\} \\ & \cup \{A^n \setminus S : S \in Def(k, A, n)\} \\ & \cup \{Proj(A, R, n) : R \in Def(k, A, n+1)\} \end{aligned}$$

(corresponding to conjunctions, negations and existence quantifier). Lastly we define

$$Def(A, n) := \bigcup_{k \in \mathbb{N}} Def(k, A, n)$$

Observe that the definition of $Def(k+1, A, n)$ and $Def(0, A, n)$ are Δ_0 because all of the quantifiers are bounded by A . So the definition of $Def(A, n)$ is a recursive definition based on absolute notions, and thus absolute for transitive models (containing A , of course).

The next thing we are going to do (a bit of preview) is to use $Def(A, n)$ to define the *definable power set*. After that, define a *definable von Neumann hierarchy*. That's what we are going to do on Wednesday.

A reminder that we're going to have an example class this afternoon (330-5pm). I think it's in MR5?

—Lecture 9—

1.7 The constructible universe

Last time we discussed the construction of an inner model $L \subseteq M$ that is based on definability. Problems with definability are connected to Tarski's Undefinability of Truth; the *definable* fragment of truth is that where we fix the scope of existential quantifiers in advance.

Recall that we defined recursively $Def(A, n)$, which are definable subsets of A^n , where *definable* is interpreted in A .

Lemma. Let $X \subseteq A^n$. If there is a formula φ s.t.

$$(x_1, \dots, x_n) \in X \iff (A, \in) \models \varphi(x_1, \dots, x_n)$$

then $X \in Def(A, n)$.

Proof. Simple induction over complexity of φ . □

Lemma. In M , we have that $Def(A, n)$ is countable.
(We even have a concrete surjection $\mathbb{N} \rightarrow Def(A, n)$.)

Proof. There are only countably many formulas. □

Observation: in the definition of $Def(k, A, n)$, we only used notions absolute for transitive models of ZF. So, since $Def(A, n)$ was defined by recursion over $Def(k, A, n)$, also $Def(A, n)$ is absolute between transitive models.

Our goal is to find a definable power set. However, $Def(A, 1)$ is not a good candidate as it is always countable, so if A is uncountable, then there is $a \in A$ s.t. $\{a\} \notin Def(A, 1)$ by the above lemma.

We define

$$\mathcal{D}(A) := \{X \subseteq A : \exists n \exists s \in A^n \exists R \in Def(A, n+1) \text{ s.t. } X = \{a \in A : (a, s_0, \dots, s_{n-1}) \in R\}\}$$

and we'll call this the *definable power set*.

Observation: If X is informally definable with parameters from A , i.e. $X = \{a \in A : (A, \in) \models \varphi(a, p_1, \dots, p_n)\}$ for some $p_1, \dots, p_n \in A$, then $X \in \mathcal{D}(A)$.

Observation: As $Def(A, n)$ was absolute and the quantifiers in the definition of $\mathcal{D}(A)$ are all bounded, $\mathcal{D}(A)$ is absolute for transitive models.

Proposition. If A is transitive, then $\mathcal{D}(A)$ is transitive.

Proof. Suppose $x \in X \in \mathcal{D}(A)$. Then $x \in X \subseteq A$. So $x \subseteq A$ and $x \in A$ as A is transitive. We can then define x as a subset by the formula $v \in x = \varphi(v)$ (which selects exactly the elements of x , which form x): $x = \{z \in A : (A, \in) \models \varphi(z)\}$. \square

Let $L_0 := \phi$,
 $L_{\alpha+1} := \mathcal{D}(L_\alpha)$,
 $L_\lambda := \bigcup_{\alpha < \lambda} L_\alpha$.

We refer to the class $\bigcup_{\alpha \in \text{Ord}} L_\alpha$ as ' L ', or *the constructible universe*.

Properties:

- (1) If $\alpha \leq \omega$, then $V_\alpha = L_\alpha$.
- (2) For every α , L_α is transitive from proposition by induction.
- (3) $\alpha \leq \beta \rightarrow L_\alpha \subseteq L_\beta$;
- (4) $\text{Ord} \cap L_\alpha = \alpha$ (this and (3) are in sheet 2);
- (5) If $\alpha \geq \omega$, then $|L_\alpha| = |\alpha|$.

Proof. $\alpha = \omega$: $|L_\omega| = |V_\omega| = \aleph_0 = |\omega|$;

Suppose for all $\beta < \alpha$ we have $|L_\beta| \leq |\beta|$. Show that $|L_\alpha| < |\alpha|^+$:

- (a) α is a successor, say $\alpha = \beta + 1$. So $L_\alpha = L_{\beta+1} = \mathcal{D}(L_\beta)$; if we look carefully at the definition, we could get a surjection from $\aleph_0 \times \bigcup_{n \in \mathbb{N}} L_\beta^n$ onto $\mathcal{D}(L_\beta)$; but we assumed α is at least ω , $\aleph_0 \times \bigcup_{n \in \mathbb{N}} L_\beta^n$ has cardinality $|L_\beta|$ as well.
- (b) α is a limit. Let $\pi_\beta : \alpha \rightarrow L_\beta$ be a surjection. Then we find surjection from $\alpha \times \alpha \twoheadrightarrow L_\alpha$ by $(\gamma, \gamma') \rightarrow \pi_\gamma(\gamma')$. \square

Note that in the above we've only proved that the cardinality is at most what we wanted; but it is also at least that by property (4).

Now look at $V_{\omega+1}$ and $L_{\omega+1}$. They can't be the same, as the first is uncountable but the second is countable. So $L_{\omega+1} \subseteq V_{\omega+1}$, and $V_{\omega+1} \setminus L_{\omega+1} \neq \emptyset$. However, it is entirely possible that the missing part is collected later when we reach L_{ω_1} or later.

Definition. If x is constructible, $x \in L$, then $\rho_L(x) := \min\{\alpha : x \in L_{\alpha+1}\}$ (an analogue of the rank in V).

Definition. $V = L$ is the axiom of constructibility

$$\forall x \exists \alpha (x \in L_\alpha)$$

Note that this is a concrete sentence in L_∞ , as L_α can be recursively defined.

Lemma. If M is a transitive set model of $ZFC + (V = L)$, then there is a limit ordinal λ s.t. $M = L_\lambda$.

Note that we haven't shown that such M would exist (as $\neg V = L$ might well be a theorem in ZFC), but we know what it will look like *if* it exists.

We don't really have much time, so we'll present a quick proof here; more details next lecture.

Proof. Consider $\lambda := \text{Ord} \cap M$ (remember that M is a set, so λ is indeed a set, so an ordinal). Clearly, λ is a limit ordinal (both are transitive).

We claim that $M = L_\lambda$: suppose $x \in M$, by $V = L$ find $\alpha \in M$ s.t. $(M, \in) \models x \in L_\alpha$. L_α was defined by recursion from absolute notions, so $x \in L_\alpha$ is absolute. So $x \in L_\alpha \subseteq L_\lambda$.

Suppose $x \in L_\lambda$, so there is $\alpha < \lambda$ s.t. $x \in L_\alpha$ (this is absolute, so $(M, \in) \models x \in L_\alpha$), by choice of λ , $\alpha \in M$. \square

—Lecture 10—

What did we do so far? We defined the constructible universe, and we realized that our definitions were absolute for transitive models: if we write $\Phi(x, \alpha)$ the formula ' $x \in L_\alpha$ ', $\Psi(x, \alpha)$ the formula ' $x = L_\alpha$ ', then both Φ and Ψ are absolute for transitive models of set theory.

We also had the axiom ' $V = L$ ', which is a short hand for $\forall x \exists \alpha x \in L_\alpha$.

Proposition. If A is a transitive set model of $ZFC + (V = L)$, then $A = L_\alpha$ where $\alpha = \text{Ord} \cap A$.

Proof. ' $A \subseteq L_\alpha$ ': If $x \in A$, then by $(A, \in) \models V = L$, find β s.t. $(A, \in) \models x \in L_\beta$; by absoluteness, $x \in L_\beta \subseteq L_\alpha$.

' $L_\alpha \subseteq A$ ': If $x \in L_\alpha$, since $\alpha = \text{Ord} \cap A$ and $A \models ZFC$, α has to be a limit ordinal. So we can write $L_\alpha = \bigcup_{\beta < \alpha} L_\beta$. So find $\beta < \alpha$ and $x \in L_\beta$. Since $(A, \in) \models ZFC$, we know that A thinks ' L_β exists', or slightly more formally, there is X s.t. $(A, \in) \models \Psi(X, \beta)$. By absoluteness of Ψ , $\Psi(X, \beta)$ is actually true in the bigger universe M . So $X = L_\beta$. Therefore $L_\beta \in A$. But $x \in L_\beta$ and A is transitive. So $x \in A$. \square

However, as we said last time, we don't really know if there *is* actually a model of $ZFC + (V = L)$, so what we just did seems a bit pointless. Our next major goal shall then be:

Theorem. (Gödel, 1938)

If κ is inaccessible, then $L_\kappa \models ZFC + (V = L)$.

Corollary. $\text{IC} \implies$ there is a *countable* α s.t. $L_\alpha \models ZFC + (V = L)$.

Proof. Take $H := \mathcal{H}^{L_\kappa}(\phi) \prec L_\kappa$, the Skolem hull. H is therefore countable. Then take $T \cong H$ the Mostowski collapse, then $(T, \in) \equiv (L_\kappa, \in)$ (all sentences have the same truth value). Therefore $(T, \in) \models ZFC + (V = L)$ is a countable transitive model; then apply our previous proposition we know $T = L_\alpha$ for some ordinal α . But by a previous property $|L_\alpha| = |\alpha|$, so α has to be countable. \square

Contrast this with: If $V_\alpha \models ZFC$, then α cannot be countable: if α is countable, then there is a code for a surjection $f : N \twoheadrightarrow \alpha$ in $V_{\omega+1} \subseteq V_\alpha$, so $V_\alpha \models \text{'}\alpha \text{ is countable'}$; contradiction to $V_\alpha \models ZFC$. (some diagram)

Proof. (of theorem)

Extensionality follows from transitivity;

Pair: $x, y \in L_\kappa$, find α s.t. $x, y \in L_\alpha$ s.t. $\{x, y\} \subseteq L_\alpha$. Clearly the formula

$$\varphi(z, x, y) := z = x \vee z = y$$

defines the pair $\{x, y\}$. So by one of our previous lemmas, the pair lies in $\mathcal{D}(L_\alpha) = L_{\alpha+1} \subseteq L_\kappa$.

However, note that our lemma only says that if we have a formula φ , and we interpret it in L_α , then what we get will be in $D(L_\alpha)$. But luckily our φ here is simple enough that it doesn't matter where we interpret it within: its truth value is the same in any model that contains x and y .

The same proof takes care of, say, Union. Let's get to something that's more interesting. How about Power Set? This can't work directly, since there are just not enough elements in $D(L_\alpha)$.

Consider $x \in L_\kappa$. As before, $\alpha < \kappa$, so $x \in L_\alpha$, so $x \subseteq L_\alpha$ by transitivity, so $|x| \leq |L_\alpha| < \kappa$ because κ is strongly inaccessible.

Consider $\mathcal{P}(x)$ in M ; we know $|\mathcal{P}(x)| = 2^{|x|} < \kappa$ because κ is inaccessible, and $|L_\kappa \cap \mathcal{P}(x)| \leq |\mathcal{P}(x)| < \kappa$ (diagram).

Remark. Since $L_\kappa, \mathcal{P}(x)$ are sets in M , $L_\kappa \cap \mathcal{P}(x)$ is a set in V_κ , and it is *definable* in V_κ by the formula

$$z \in L_\kappa \cap \mathcal{P}(x) \iff \Phi(z, \kappa) \wedge z \subseteq x$$

But that's not good enough to prove that $L_\kappa \cap \mathcal{P}(x) \in L_\kappa$, as we actually need to find an $\alpha < \kappa$ in place of κ in the formula.

For each $z \in L_\kappa \cap \mathcal{P}(x)$, find the L -rank,

$$\alpha_z := \rho_L(z) < \kappa$$

and consider the set

$$\{\alpha_z : z \in L_\kappa \cap \mathcal{P}(x)\} \subseteq \kappa$$

of size $< \kappa$. Then by regularity of κ we know it must be bounded by some $\beta < \kappa$ s.t.

$$\{\alpha_z : z \in L_\kappa \cap \mathcal{P}(x)\} \subseteq \beta$$

So $L_\kappa \cap \mathcal{P}(x) \subset L_\beta$.

Now define

$$P := \{z : (L_\beta, \in) \models z \subseteq x\}$$

By our lemma, $P \in \mathcal{D}(L_\beta) = L_{\beta+1} \subseteq L_\kappa$; but $L_\kappa \models \forall z(z \in P \iff z \subseteq x)$, so L_κ is a model of the power set axiom. (see diagram)

—Lecture 11—

Example sheet 2 will be in an envelope next to C0.10 this afternoon.

We saw pairing and power set last time. Some of the others will be in the example sheet.

Let's consider separation, a more difficult one. Let $x \in L_\kappa$, φ be a formula, $a_1, \dots, a_k \in L_\kappa$.

Separation says that, informally, $\{z \in x : \varphi(z, a_1, \dots, a_n)\}$ exists.

More formally, $\{z \in x : L_\kappa \models \varphi(z, a_1, \dots, a_n)\}$ exists. At first glance we can use the same argument for power set: just find all these z are, and go to the next level and consider \mathcal{D} of it. Let's see how that *stops* working. For each $1 \leq i \leq n$, find $\alpha_i < \kappa$ s.t. $a_i \in L_{\alpha_i}$, and find $\alpha < \kappa$ s.t. $x \in L_\alpha$. Define $\beta := \max\{\alpha, \alpha_1, \dots, \alpha_n\}$.

So for any $z \in x$, we have $z, a_1, \dots, a_n \in L_\beta$ (by transitivity of L_β). So $\{z \in x : L_\beta \models \varphi(z, a_1, \dots, a_n)\} \in \mathcal{D}(L_\beta) = L_{\beta+1}$. Apparently this is not necessarily what we want: we have no idea if φ is absolute between L_β and L_κ ; in fact we have *some* idea, because this has to work for any φ , so there are definitely some formulas that are not absolute.

So we see the problem is that, in general, $\{z \in x : L_\beta \models \varphi(z, a_1, \dots, a_n)\} \neq \{z \in x : L_\kappa \models \varphi(z, a_1, \dots, a_n)\}$.

Note that our choice of β might be a terrible one, as we have no idea what it is; it might even be a successor ordinal. So we have to choose it more carefully.

Consider $\mathcal{H}^{L_\kappa}(L_\beta) \prec L_\kappa$, with cardinality the same as $|L_\beta| = |\beta| < \kappa$; and now consider its Mostowski collapse, $T \cong \mathcal{H}^{L_\kappa}(L_\beta)$, with Mostowski isomorphism $\pi : T \rightarrow \mathcal{H}^{L_\kappa}(L_\beta)$. T is transitive, $|T| = |\beta| < \kappa$, and $T \prec L_\kappa$.

[Some model theory recap: we have if $M \cong M'$, $M \models \varphi(x_1, \dots, x_n) \iff M' \models \varphi(\pi(x_1), \dots, \pi(x_n))$. $M \equiv M'$ means $M \models \varphi \iff M' \models \varphi$ for all sentences φ . Therefore isomorphism implies elementary equivalence. $M \prec M'$ means $M \models \varphi(x_1, \dots, x_n) \iff M' \models \varphi(x_1, \dots, x_n)$.]

We then have

$$T \xrightarrow{\pi} \mathcal{H}^{L_\kappa}(L_\beta) \xrightarrow{id} L_\kappa$$

$$\varphi(z, a_1, \dots, a_n) \quad \varphi(\pi(z), \pi(a_1), \dots, \pi(a_n)) \quad \varphi(\pi(z), \pi(a_1), \dots, \pi(a_n))$$

We know that the Mostowski collapse is the identity on transitive sets. So, if $X \subseteq L_\kappa$ is transitive, then $\pi|_X = id|_X$. Since L_β is transitive, and all z, a_1, \dots, a_n are in L_β , we get that $T \models \varphi(z, a_1, \dots, a_n) \iff L_\kappa \models \varphi(z, a_1, \dots, a_n)$.

Let's do the 'modified Skolem Hull construction' from sheet 1 q12: let $\alpha_0 := \text{Ord} \cap T$, α_{n+1} is the least γ s.t. L_γ contains L_{α_n} and a witness for each existential statement true with parameters in L_{α_n} . Let $\bar{\alpha} := \bigcup_{n \in \mathbb{N}} \alpha_n$. So $T \prec L_{\bar{\alpha}}$ with $\bar{\alpha} < \kappa$.

Now define the set via φ over $L_{\bar{\alpha}}$:

$$\begin{aligned} L_{\bar{\alpha}+1} &= \mathcal{D}(L_{\bar{\alpha}}) \ni \{z \in x : L_{\bar{\alpha}} \models \varphi(z, a_1, \dots, a_n)\} \\ &= \{z \in x : T \models \varphi(z, a_1, \dots, a_n)\} \ (T \prec L_{\bar{\alpha}}) \\ &= \{z \in x : \mathcal{H}^{L_\kappa}(L_\beta) \models \varphi(\pi(z), \pi(a_1), \dots, \pi(a_n))\} \\ &= \{z \in x : \mathcal{H}^{L_\kappa}(L_\beta) \models \varphi(z, a_1, \dots, a_n)\} \\ &= \{z \in x : L_\kappa \models \varphi(z, a_1, \dots, a_n)\} \end{aligned}$$

We only showed pairing, power set, and separation. The rest are similar. \square

Theorem. (The Condensation Lemma)

If κ is inaccessible, $x, y \in L_\kappa$, and $y \subseteq x$, then there is $\alpha < \kappa$ with $|\alpha| \leq |tcl(x)|$ s.t. $y \in L_\alpha$.

Proof. Consider $y \subseteq x$ and $tcl(x \cup \{y\}) =: t$. Clearly $|t| = |tcl(x)|$. Now consider $T \xrightarrow{\pi} \mathcal{H}^{L_\kappa}(t) \prec L_\kappa$, clearly all these things have the same cardinality. Now we know π is the identity on t , so in particular $\pi(y) = y$, $\forall z \in x \pi(z) = z$, and $\pi(x) = x$.

By our lemma, find β s.t. $T = L_\beta$ (we're now allowed to do this since $L_\kappa \models ZFC + (V = L)$, and therefore $L_\beta \equiv L_\kappa \models ZFC + (V = L)$).

Now y can be defined over L_β with parameters in L_β , namely viz. y (we know y is in there so we don't actually need to define it).

We just need to check β has the right cardinality: $|\beta| = |L_\beta| = |T| = |tcl(x)|$. \square

—Lecture 12—

Today we're going to prove that there is a model of ZFC+CH. This is especially convenient as today we're in the middle of the entire course (the 12th lecture), so we spent exactly half of the course on proving one side of what we wanted.

Recall that last time we had the condensation lemma: if κ is inaccessible, $x, y \in L_\kappa$, $y \subseteq x$, then there is α s.t. $|\alpha| \leq |tcl(x)|$ s.t. $y \in L_\alpha$.

Corollary. $x = \mathbb{N}$, $y \subseteq \mathbb{N}$, then there is $\alpha < \omega$ s.t. $y \in L_\alpha$.

Corollary. (1) $\mathcal{P}(\mathbb{N}) \cap L_\kappa \subseteq L_{\omega_1}$.

Observe that $\mathcal{P}(\mathbb{N}) \cap L_\kappa = \mathcal{P}^{L_\kappa}(\mathbb{N})$, where $\mathcal{P}^{L_\kappa}(\mathbb{N})$ refers to the unique $p \in L_\kappa$ s.t. $L_\kappa \models p = \mathcal{P}(\mathbb{N})$.

(2) $\mathcal{P}^{L_\kappa}(\mathbb{N}) \subseteq L_{\omega_1}$.

(3) $|\mathcal{P}^{L_\kappa}(\mathbb{N})| \leq |L_{\omega_1}| = |\omega_1| = \aleph_1$.

Hence it seems that we're extremely close to having a model of ZFC+CH. However, this ω_1 we're referring to here is, of course, in the model that we're building.

(A 'one-line proof' of the Condensation Lemma:

$$y \in L_\alpha = T \cong \mathcal{H}^{L_\kappa}(tcl(x) \cup \{y\}) \prec L_\kappa$$

)

Let's improve this to show $L_\kappa \models CH$. The key idea is that $L_\kappa \models ZFC$, the condensation lemma itself is a theorem of $ZFC(?)$, so we can just apply condensation lemma inside L_κ .

The problem is that our condensation lemma is a theorem of $ZFC+IC$ instead of ZFC .

Remark. So if you assume $ZFC+2IC$ in M , then this argument gets you that L_κ (where κ should be the second inaccessible cardinal now) is a model of CH , and you're actually done provided that you think $2IC$ is a reasonable thing to have.

But this feels a bit odd. Let's try to do this without $2IC$.

Work in L_κ , we know that $\mathcal{P}^{L_\kappa}(\mathbb{N}) \subseteq L_{\omega_1}$, where ω_1 is the ω_1 in M . Note that $\omega_1 < \kappa$, so in particular $L_{\omega_1} \subseteq L_\kappa$.

Now we could build $\mathcal{H}^{L_\kappa}(L_{\omega_1}) \prec L_\kappa$, and $\mathcal{P}^{L_\kappa}(\mathbb{N}) \subseteq L_\beta = T \cong \mathcal{H}^{L_\kappa}(L_{\omega_1})$. By the standard argument we know $\beta < \aleph_2 < \kappa$. But now $L_\beta \models ZFC + V = L$.

Run the condensation lemma proof for V_κ as M and L_β as L_κ , we then get

$$y \in L_\alpha = T \cong \mathcal{H}^{L_\beta}(\mathbb{N} \cup \{y\}) \prec L_\beta$$

Now the surrounding universe knows that α is countable, so L_κ knows α is countable. So if $\omega_1^{L_\kappa}$ is the ω_1 of L_κ , then $\mathcal{P}^{L_\kappa}(\mathbb{N}) \subseteq L_{\omega_1^{L_\kappa}}$.

So $L_\kappa \models 2^{\aleph_0} \leq \aleph_1$, i.e. $L_\kappa \models CH$. So we've proved:

Theorem.

$$Cons(ZFC + IC) \implies Cons(ZFC + CH)$$

Remark. (1) The same argument with $x = \lambda$ for some L_κ -cardinal λ gives us $L_\kappa \models 2^\lambda \leq \lambda^+$. So GCH holds: $\forall \lambda 2^\lambda = \lambda^+$.

(2) What about the inaccessible cardinal?

(a) If we have a transitive set model of ZFC , we can mimic this proof. Nevertheless we knew it was still stronger than $Cons(ZFC)$.

(b) There is a way of getting around that assumption as well by Lévy Reflection Theorem (see sheet 2): fix in advance some finite list Φ of sentences you wish to preserve, and find sufficiently large α s.t. Φ is absolute for V_α .

Go through all needed absoluteness results and lemmas and theorems, and collect, for each of them, φ , the finite set Φ_φ of axioms of ZFC we needed to prove them. Form $\Phi := \bigcup_{\varphi \text{ is relevant}} \Phi_\varphi$, which is a finite union of finitely many things. Apply Lévy to Φ and run the previous proof to get a model of $\Phi + CH$. Now consider all finite subsets Ψ s.t. $\Phi \subseteq \Psi \subseteq ZFC$, and get models of $\Psi + CH$. Compactness then gives a model of $ZFC + CH$.

(3) Consider

$$L = \bigcup_{\alpha \in Ord} L_\alpha \subseteq M$$

Our proof does not say what axioms hold in L , but using the Lévy reflection theorem, we can prove that if $V \models ZFC$, then $L \models ZFC + V = L$.

Now let's use this to solve one of the open problems in the 1910s, which follows as an almost trivial consequence of what we've done.

Consider the question about regular limit cardinals. Remember we had notions of regular/singular, successor/limit. We realize that we can prove that every successor cardinal is regular, and we have a lot of singular limit cardinals. So can we get regular limit cardinals?

If we strengthen 'limit' ($\forall \lambda < \kappa (\lambda^+ < \kappa)$) to 'strong limit' ($\forall \lambda < \kappa (2^\lambda < \kappa)$), then we showed that ZFC cannot prove the existence of regular strong limits (i.e. inaccessible cardinals).

Clearly, ZFC+GCH implies that every limit is a strong limit (because $\lambda^+ = 2^\lambda$). So ZFC+CH gives every regular limit is an inaccessible cardinal.

Proof. Assume that $M \models ZFC$ and that $ZFC \vdash$ there are regular limits. Towards a contradiction (with Gödel's incompleteness theorem), prove $M \models Cons(ZFC)$.

Consider $L \subset M$, by remark (3), $L \models ZFC + GCH$. By assumption, $ZFC \vdash \exists$ regular limit, so $L \models ZFC + GCH + (\exists \kappa \text{ } \kappa \text{ is a regular limit})$. But ZFC+GCH implies every regular limit is inaccessible, so $L \models ZFC + IC$. So $L \models \exists \kappa (L_\kappa \models ZFC)$, so $L \models Cons(ZFC)$; so $M \models Cons(ZFC)$ by absoluteness between transitive models. \square

2 Example Class 1

2.1 Question 1

The answer to this question really depends on how we formulate the axioms. For example, the pairing axiom

$$\forall x \forall y \exists p \forall z (z \in p \leftrightarrow z = x \vee z = y)$$

is sometimes formulated with the \leftrightarrow changed to \rightarrow , but of course they are equivalent with extensionality.

So there is no definite answer to this question; the message I want to convey here is that the axioms really only make sense when you have a set of those.

2.2 Question 2

Note that (a) is not Δ_0 , although it is Δ_0 under some first order logic.

2.3 Question 3

(d) if z is a function then $\mathcal{P}(\bigcup \bigcup z)$ is enough to bound the range and domain of the function.

2.4 Question 6

Example given on lecture: $X = \{(x, y) : x < \omega, y \in \{0, 1\}\}$;
 $R = \{(a, b) : a, b = X, a = (a_1, a_2), b = (b_1, b_2), (b_2 = a_2 \wedge a_1 < b_1) \vee (b_2 > a_2)\}$.
 Both of them are in $V_{\omega+1}$, and $(X, R) \in V_{\omega+4}$ or so.

2.5 Question 8

The general case is the fixed point theorem for normal functions. A normal function is a function $F : Ord \rightarrow Ord$ that is

- increasing: $\forall \alpha \alpha < F(\alpha)$;
- continuous: $\forall \lambda$ limits, $F(\lambda) = \bigcup_{\alpha < \lambda} F(\alpha)$.

Define $\gamma_0 := \gamma$, $\gamma_{n+1} := F(\gamma_n)$, $\gamma_\omega := \bigcup_{n \in \mathbb{N}} \gamma_n$. We basically just need to check that the fixed point happens at the third case (so that the cofinality is ω).

2.6 Question 9

(crude notes on board) κ is uncountable and regular. So we know $H_\kappa \models ZFC - PowerSet$, $V_\kappa \models ZFC - Replacement$. Let $\lambda < \kappa$ be a cardinal. Then $\lambda \in H_\kappa$ and every subset of λ is in H_κ . So $H_\kappa \models Powerset \implies P(\lambda) \in H_\kappa$, and $|P(\lambda)| \leq |tcl(P(\lambda))| < \kappa$. So κ is a strong limit. •

2.7 Question 10

Remember Hartog's theorem: $\forall x \exists \alpha$ a ordinal s.t. there is no injection from α into X ; remember that this is true in ZF.

Let $\Omega := \{R : R \text{ is a binary relation on } X, (X, R) \text{ is a well-order}\}$,
 $\bar{\Omega} := \{\alpha : \exists R \in \Omega, (\alpha, \in) \cong (X, R)\}$.

Note that this also tells us that, if we have two models of set theory which agree on real numbers, then they agree on the size of \aleph_1 .

2.8 Question 11

It's worth notice that Φ is not absolute in general (because we need strong limit for inaccessible cardinals).

For the last part just take V_{κ_2} . Also note that the statement holds for any finite number, or even countably many.

We can consider other set theories like $ZFC + \infty IC$: $\forall \alpha \exists \kappa > \alpha$ s.t. κ is inaccessible, or $ZFC +$ there is a cardinal κ which is inaccessible and $\forall \lambda < \kappa \exists \lambda' < \kappa' < \kappa$ s.t. κ' is inaccessible (so κ is an inaccessible limit of inaccessible cardinals).

2.9 Question 12

The idea is almost the same as the one we looked in class: build the Skolem Hull starting from quantifier-free formulas, adding witnesses of existential formulas at each step. By induction we get a countable elementary substructure. Now instead of that, for each step in building the skolem hull we take the V_{α_i} s.t. our structure is contained in V_{α_i} . By regularity of κ we know we won't reach V_κ in the end, say our supremum is α_ω . Note that it's possible that we've hit an inaccessible cardinal en route. So we need to ensure that we go up by at least one level at each stage to ensure that we're actually moving. Then use similar argument as q8.