
(a) relative spectral error (open: $N = 256$)
case A
case B
uniform
perturbed lattice
clustered
error = tol
$\|\Gamma_h - \Gamma_{\mathrm{ref}}\|_2 / \|\Gamma_{\mathrm{ref}}\|_2$
T9
$10^{-9}$
T7
$10^{-7}$
T5
$10^{-5}$
parameter set (tolerance accepted on A and B)
(b) absolute error, set T7
relative $10^{-7}$
box-length sets
$\|\Gamma_h - \Gamma_{\mathrm{ref}}\|_2$
$\|\Gamma_{\mathrm{ref}}\|_2$ (set by the closest pairs)
(c) probes through $\Gamma_h X$, set T7
$\|(\Gamma_h - \Gamma_{\mathrm{ref}})X\| / \|\Gamma_{\mathrm{ref}} X\|$
translation: $/(\|\Gamma_{\mathrm{ref}}\|_2 \|X\|)$
Gaussian
long. $k_{\min}$
transv. $k_{\min}$
translation
(d) box length, set T7 (3 seeds)
$\eta$=0.61
$\eta$=0.69
$\eta$=0.61
$\eta$=0.68
$\eta$=0.69
$\eta$=0.65
fixed $N = 64$ ($n$ varies)
fixed $n = n_A$ ($N$ varies)
box length $L$