

Introductory Mathematics for Computer Science (COMP0011)

Raphael Li

Year 1 Term 2, 2024–25

Contents

1	Complex numbers	2
1.1	Basic arithmetic with complex numbers, and complex conjugates	2
1.2	Visualising complex numbers	3
1.3	Exponential form	3
1.4	Converting between Cartesian and exponential forms	4
1.5	Visualising arithmetic on complex numbers	5
1.6	Roots of unity	6

1 Complex numbers

The foundation of the *complex numbers* is given by the imaginary unit i , defined either as $i = \sqrt{-1}$ or as $i^2 = -1$.

A complex number z can be written as $a + bi$, where $a, b \in \mathbb{R}$. The real numbers a and b are known as the *real part* and the *complex part* of z respectively.

The set of all complex numbers is denoted as \mathbb{C} . Note that the set of real numbers \mathbb{R} is a subset of \mathbb{C} . See figure 1.




Figure 1: The set of real numbers \mathbb{R} is a subset of the set of complex numbers \mathbb{C} . All real numbers are complex numbers.

1.1 Basic arithmetic with complex numbers, and complex conjugates

To add or subtract two complex numbers, we deal with the real and imaginary parts separately.

$$(2 + 3i) + (5 - 8i) = (2 + 5) + (3 + (-8))i = 7 - 5i \quad (\text{Addition})$$

$$(2 + 3i) - (5 - 8i) = (2 - 5) + (3 - (-8))i = -3 + 11i \quad (\text{Subtraction})$$

The multiplication of complex numbers is also straightforward as long as we bear in mind that $i^2 = -1$.

$$\begin{aligned} (3 + 4i)(-2 + 3i) &= -6 + 9i - 8i + 12i^2 \\ &= (-6 - 12) + (9 - 8)i \\ &= -18 + i \end{aligned}$$

To divide a complex number by another, e.g.

$$\frac{a + bi}{c + di}$$

we multiply both the numerator and denominator by $c - di$, which is obtained by flipping the sign of the imaginary part of the denominator. For example, if we want to compute

$$\frac{2 + 3i}{5 - 4i},$$

we flip the sign of the imaginary part of $5 - 4i$ to get $5 + 4i$. We then multiply both the numerator and denominator of the fraction by this $5 + 4i$ to get

$$\begin{aligned} \frac{2 + 3i}{5 - 4i} &= \frac{(2 + 3i)(5 + 4i)}{(5 - 4i)(5 + 4i)} \\ &= \frac{10 + 8i + 15i - 12}{25 + 20i - 20i + 16} \\ &= \frac{-2 + 23i}{41} \\ &= \frac{-2}{41} + \frac{23}{41}i. \end{aligned}$$

Notice how multiplying $5 - 4i$ with $5 + 4i$ produces the real number 41. By flipping the sign of the imaginary part of a complex number, we obtain what's called its *complex conjugate*. The complex conjugate of z is denoted as \bar{z} . By writing z as $a + bi$, we can easily prove that the product of any complex number with its conjugate must equal a real number:

$$z \times \bar{z} = (a + bi)(a - bi) = a^2 + b^2 \in \mathbb{R}.$$

1.2 Visualising complex numbers

Given some complex number $z = x + yi$, we can treat its real and imaginary parts as Cartesian coordinates, thus mapping it to a point on the 2D plane.




Figure 2: The complex number $z = x + yi$ as a point on the 2D plane

1.3 Exponential form

Recall that it is possible to express a point on a 2D plane using polar coordinates (R, θ) as well. Indeed, given any complex number $z = x + yi$, we can find its corresponding pair of values R and θ .




Figure 3: The position of a complex number on the 2D plane can be represented using polar coordinates.

Based on this idea, we introduce a new notation as follows.

If the position of a complex number z on the 2D plane can be represented by the polar coordinates (R, θ) , then we have

$$z = R \times e^{i\theta}$$

where $R, \theta \in \mathbb{R}$ and $R \geq 0$.

R is called the *absolute value* or *modulus* of z and is denoted as $|z|$. This represents the point's position from the origin.

θ is called the *argument* of z and is denoted as $\arg(z)$. This represents the angle from horizontal.

This way of representing complex numbers is known as the *exponential form*. (This is a natural result of Euler's formula $e^{i\theta} = \cos \theta + i \sin \theta$.)

Now consider two complex numbers expressed in exponential form.

$$\begin{aligned} z_1 &= R_1 \times e^{i\theta_1} \\ z_2 &= R_2 \times e^{i\theta_2} \end{aligned}$$

These two numbers are considered equal if both of the following conditions hold.

$$\begin{aligned} R_1 &= R_2 \\ \theta_1 &= \theta_2 + 2k\pi \end{aligned} \quad (\text{for some } k \in \mathbb{Z})$$

Note that the red part is necessary because a rotation of 2π radians has no effect on a point's position.

The exponential form makes the multiplication and division of complex numbers a lot easier.

Multiplication	Division
$\begin{aligned} (1 \times e^{\frac{\pi}{6}i}) \times (2 \times e^{-\frac{\pi}{4}i}) &= 2 \times e^{\frac{\pi}{6}i - \frac{\pi}{4}i} \\ &= 2 \times e^{-\frac{\pi}{12}i} \end{aligned}$	$\begin{aligned} \frac{1 \times e^{\frac{\pi}{6}i}}{2 \times e^{-\frac{\pi}{4}i}} &= \frac{1}{2} \times \frac{e^{\frac{\pi}{6}i}}{e^{-\frac{\pi}{4}i}} \\ &= 2 \times e^{\frac{5\pi}{12}i} \end{aligned}$

1.4 Converting between Cartesian and exponential forms

The methods used to convert between the Cartesian form $x + yi$ and the exponential form $R \times e^{i\theta}$ are outlined below.

- **Given the Cartesian form of a complex number, find its exponential form.**

Given the Cartesian form $z = x + yi$, we can find the modulus using Pythagoras' theorem.

$$|z| = \sqrt{x^2 + y^2}$$

The argument can be found using the arctangent.

$$\arg(z) = \arctan\left(\frac{y}{x}\right)$$

- **Given the exponential form of a complex number, find its Cartesian form.**

Given the exponential form $z = R \times e^{i\theta}$, we can find the Cartesian coordinates using simple trigonometry.

$$\begin{aligned} x &= R \cos \theta \\ y &= R \sin \theta \end{aligned}$$

To speed up conversion processes, it is often useful to memorize the Cartesian coordinates of some special points on the unit circle. See figure 4 and table 1.




Figure 4: It is important to know the coordinates of points on the circle corresponding to classic angles.

θ (radians)	$\pi/6$	$\pi/4$	$\pi/3$
θ (degrees)	30°	45°	60°
$\sin \theta$	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$
$\cos \theta$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$

Table 1: The values of $\sin \theta$ and $\cos \theta$ for some classic angles θ .

1.5 Visualising arithmetic on complex numbers

When visualised on the 2D plane, the addition of complex numbers is similar to that of vectors. We join the arrows in a tip-to-tail manner in order to determine the sum, as shown in figure 5.




Figure 5: Addition of complex numbers.

The above figure also illustrates another key idea. Notice how in the figure on the right, the vectors of z_1 , z_2 and $z_1 + z_2$ form a triangle. This means their absolute values must fulfil the triangle inequality.

$$|z_1| + |z_2| \geq |z_1 + z_2|$$

To visualise multiplication we consider the exponential form. As shown in figure 6, when two complex numbers are multiplied, their arguments are added together to produce a rotation, while their moduli are multiplied.




Figure 6: Multiplication of complex numbers.

1.6 Roots of unity