
FedBCa public dataset (Zenodo 10409145)
Patients with urothelial carcinoma, MRI ≤ 2 weeks before TURBT or cystectomy,
no prior treatment; four Chinese centers, 2019–2022
Released cases with T2WI, expert lesion mask(s) and pathology label
221 patients · 275 lesions · no exclusions in this study
(descriptor text states 228 patients; its Table 1 and the released files contain 221)
Center 1
Siemens Skyra
119 patients (30 MIBC)
160 lesions (30 MIBC)
multifocal patients: 18
Center 2
United Imaging uMR780
47 patients (25 MIBC)
48 lesions (25 MIBC)
multifocal patients: 1
Center 3
GE MR750w
24 patients (12 MIBC)
32 lesions (13 MIBC)
multifocal patients: 5
Center 4
Siemens Verio
31 patients (16 MIBC)
35 lesions (17 MIBC)
multifocal patients: 3
Patient-level analysis (primary)
221 patients · 83 MIBC (37.6%)
patient score = maximum lesion probability
sensitivity: unifocal patients only, n = 194 (77 MIBC)
Lesion-level analysis (secondary, Part B)
275 lesions · 85 MIBC (30.9%)
MIBC = pT2–T4; NMIBC = pTa/pT1
patient-clustered bootstrap for all CIs
Leave-one-center-out validation: each center held out once (4 folds);
all preprocessing, tuning and model selection within the three training centers.
Grade and T stage were never used as predictors.