Deep Learning Enhanced Virtual Sensing for Water Quality Estimation

March 2, 2024

Efficient monitoring of water quality parameters is crucial for the successful management of aquaculture systems, ensuring the health and growth of aquatic life. Traditional sensor-based systems are often cost-prohibitive, limiting their adoption, particularly in resource-constrained settings. In this research, we introduce a cost-effective method that leverages virtual sensing to estimate dissolved oxygen (DO) levels using solubility and interdependency equations. Furthermore, we propose deep learning-enhanced virtual sensing (DLEVS) by employing deep recurrent neural networks (DL-RNN) to accurately estimate DO.

Virtual Sensing Using Solubility Equations

Our virtual sensing (VS) methodology leverages established solubility equations to estimate water quality parameters, with a specific focus on dissolved oxygen (DO). By utilizing mathematical models derived from comprehensive datasets, our VS system offers a cost-effective and practical alternative to physical sensors. The flexibility of this approach allows us to estimate DO concentrations irrespective of the source of water, making it applicable to various water bodies.

VS Capabilities Irrespective of Water Source

The VS system, based on solubility equations, is designed to estimate water quality parameters across diverse water sources. Through mathematical models, the system predicts parameters that are challenging to measure directly through physical sensors. This flexibility ensures accurate estimations for a wide range of scenarios.

Deep Learning Enhancement for Accuracy

To enhance the accuracy of our VS system, we have integrated deep learning recurrent neural network (DL-RNN) models through our Dissolved Oxygen Enhanced Virtual Sensing (DLEVS) mechanism. This mechanism aims to compensate for potential inaccuracies in the VS-based DO estimation through a multi-step process.

DL-RNN Training and Error Prediction

The DL-RNN model undergoes training using a variety of water quality datasets to effectively predict errors in VS-based DO estimates. This comprehensive training process ensures the adaptability and high predictive accuracy of the model across a diverse range of water sources. The utilization of multiple datasets contributes to the robustness of the DL-RNN, allowing it to capture the intricacies and variations present in different water quality scenarios.

Real-Time Measurements and DL Enhancement

In real-time scenarios, salinity and temperature are measured, and the VS system estimates DO using solubility equations. The DL-RNN model predicts errors based on the dynamic characteristics of the specific water source. By subtracting the predicted error from the VS-estimated DO, we obtain an enhanced DO estimate that is more accurate and reflective of real-time conditions.

Conclusion

In conclusion, our VS system, leveraging solubility equations and DL-RNN models, demonstrates the capability to estimate water quality parameters irrespective of the source of water. The integration of DL-RNN models through the DLEVS mechanism enhances the accuracy of DO estimation by predicting and compensating for potential errors. Real-time measurements of salinity and temperature, coupled with virtual sensing and DL enhancement, provide a comprehensive and accurate approach to water quality estimation.