

# SI Appendix

Survival-guided matrix factorization identifies reproducible prognostic programs in pancreatic cancer

Amber M. Young, Alisa Yurovsky, Xianlu Laura Peng, Jen Jen Yeh, Didong Li, Naim U. Rashid

## Contents

|                                                                          |           |
|--------------------------------------------------------------------------|-----------|
| <b>List of Figures</b>                                                   | <b>2</b>  |
| <b>List of Tables</b>                                                    | <b>3</b>  |
| <b>I Supplementary Methods</b>                                           | <b>4</b>  |
| <b>1 Model details</b>                                                   | <b>4</b>  |
| <b>2 Optimization Algorithm</b>                                          | <b>4</b>  |
| 2.1 Block coordinate descent scheme . . . . .                            | 4         |
| 2.2 Update for $H$ . . . . .                                             | 4         |
| 2.3 Update for $W$ . . . . .                                             | 5         |
| 2.4 Update for $\beta$ . . . . .                                         | 6         |
| 2.5 Normalization of $W$ . . . . .                                       | 7         |
| 2.6 Derivation of $W$ update from projected coordinate descent . . . . . | 7         |
| <b>3 Convergence proof</b>                                               | <b>8</b>  |
| 3.1 Effect of column normalization of $W$ . . . . .                      | 11        |
| 3.2 Remark (Coxnet implementation) . . . . .                             | 11        |
| <b>4 Supervision on <math>W</math> versus <math>H</math></b>             | <b>11</b> |
| <b>5 Cross-validation procedure</b>                                      | <b>12</b> |
| <b>6 Bayesian Optimization</b>                                           | <b>13</b> |
| <b>7 PDAC Datasets</b>                                                   | <b>14</b> |
| <b>8 Survival Analysis</b>                                               | <b>14</b> |

|    |                                                                             |    |
|----|-----------------------------------------------------------------------------|----|
| 9  | Gene Program Correlation Analysis                                           | 15 |
| 10 | Variance and Survival Contribution per Factor                               | 16 |
| 11 | Factor Correspondence Analysis                                              | 16 |
| II | Supplementary Results                                                       | 17 |
| 12 | The DeSurv model shows consistent convergence and stability across restarts | 17 |
| 13 | Simulation results under null and mixed scenarios                           | 17 |
| 14 | Cross-validated C-index across factorization rank                           | 18 |
| 15 | Standard NMF rank selection heuristics                                      | 18 |
| 16 | Cutpoint selection and validation of dichotomized risk groups               | 19 |
| 17 | Overlap of DeSurv risk groups with known molecular subtypes                 | 19 |
| 18 | Standard NMF factor structure at independently selected rank                | 19 |
| 19 | Factor-Specific Gene Lists                                                  | 21 |
| SI | References                                                                  | 23 |

## List of Figures

|    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |    |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| S1 | Convergence trajectories of the DeSurv objective across five randomly selected multi-start initializations on the PDAC training cohort (TCGA + CPTAC). Each line shows the relative loss (loss at iteration $t$ divided by the initial loss for that restart) as a function of iteration number, restricted to the first 5,000 iterations for readability. The total objective decreases monotonically and stabilizes well within the iteration budget across all initializations, confirming convergence of the alternating block-update scheme. . . . .                                                                                                                                                                                                                                                                                                                                                                                            | 17 |
| S2 | DeSurv performance under null and mixed simulation scenarios. (A–B) Null scenario ( $\beta = 0$ ): both methods yield C-index $\approx 0.5$ (A) and comparable rank distributions (B), confirming DeSurv does not hallucinate prognostic structure; precision is omitted as no true survival genes exist. (C–F) Mixed scenario: survival depends on factor-specific marker genes (50%) and shared background genes (50%). (C) C-index is modestly higher for DeSurv. (D) Precision by gene-set definition: marker-only precision (150 genes; left facet) substantially favors DeSurv; full 300-gene survival-set precision (right facet) shows a more modest improvement. (E) The matched factor receives a substantially larger Cox coefficient $ \beta $ under DeSurv, confirming preferential amplification of marker-specific signal. (F) Selected-rank distributions. DeSurv (blue); NMF (red, $\alpha = 0$ ); both Bayesian-optimized. . . . . | 26 |
| S3 | Training (CV) and external validation C-index as a function of factorization rank $k$ for DeSurv (blue) and standard NMF (red, $\alpha = 0$ ). Shaded ribbons show $\pm 1$ SE. For each $k$ , the best $\alpha$ was selected by maximizing the mean CV C-index. Validation C-index is pooled across all external cohorts. (A) Results using $n_{\text{top}} = 270$ top genes per factor ( $\lambda = 0.349$ , $\xi = 0.056$ ). (B) Results retaining all genes ( $n_{\text{top}} = \text{ALL}$ ; $\lambda = 0.349$ , $\xi = 0.056$ ). . . . .                                                                                                                                                                                                                                                                                                                                                                                                        | 27 |
| S4 | Standard unsupervised NMF rank selection heuristics for the PDAC training cohort (TCGA + CPTAC). (A) Reconstruction residuals as a function of factorization rank $k$ . (B) Cophenetic correlation coefficient as a function of $k$ . (C) Mean silhouette width across multiple distance metrics as a function of $k$ . These criteria yield inconsistent guidance, motivating the survival-supervised model selection approach (main text Fig. 2D). . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | 28 |
| S5 | Cutpoint selection and external validation of dichotomized risk groups. (A) Cross-validated mean absolute log-rank z-statistic as a function of z-score cutpoint; red dashed line indicates the selected optimum. (B) Pooled validation Kaplan–Meier curves (stratified log-rank test). (C–D) Per-cohort validation KM curves: (C) Dijk, (D) Moffitt. (E–F) Per-cohort validation KM curves: (E) PACA-AU, (F) Puleo. . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | 29 |
| S6 | Subtype composition of DeSurv (top) and standard NMF $k = 3$ (bottom) risk groups in pooled external validation cohorts. Stacked bar charts show the proportion of PurIST (left) and DeCAF (right) subtypes within Low and High risk groups defined by the log-rank-optimized cutpoint. Numbers indicate sample counts; Fisher’s exact test P-values are shown above each panel. DeSurv risk groups are enriched for both Basal-like (PurIST) and proCAF (DeCAF) poor-prognosis subtypes; NMF risk groups separate Basal-like from Classical tumors but show no significant CAF subtype enrichment. . . . .                                                                                                                                                                                                                                                                                                                                          | 30 |
| S7 | Gene program overlap heatmap for DeSurv at elbow-selected rank $k = 5$ . Each cell shows the number of top genes from a given factor (rows) overlapping established PDAC gene programs (columns). DeSurv at $k = 5$ splits the ECM/activated-stroma (proCAF-associated) signal into two factors with opposing Classical-tumor correlations and isolates restCAF as a separate factor; exocrine and Basal-like programs are not represented as dominant factors. . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 31 |
| S8 | Gene program overlap heatmap for standard NMF at elbow-selected rank $k = 5$ . Each cell shows the number of top genes from a given factor (rows) overlapping established PDAC gene programs (columns). The three core programs from $k = 3$ are recovered; two additional factors overlap Exocrine-like and iCAF signatures, capturing transcriptional variance not aligned with survival outcomes. . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | 32 |

|     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |    |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| S9  | Gene program overlap heatmap for standard NMF at $k = 7$ (BO-selected rank with $\alpha = 0$ ). Compared with $k = 3$ and $k = 5$ , the 7-factor solution shows increased fragmentation: multiple factors partially capture the same PDAC gene signatures and no new coherent biological program emerges. . . . .                                                                                                                                                                                                | 33 |
| S10 | Fraction of expression variance explained versus Type III partial likelihood contribution for each standard NMF factor at elbow-selected $k = 5$ . Each point represents one factor, labeled N1–N5. Survival contribution is quantified as the reduction in Cox partial log-likelihood when each factor is removed from the full model. The two factors new relative to $k = 3$ (N4, N5) contribute negligibly to survival, confirming that elbow selection recovers no additional prognostic structure. . . . . | 34 |

## List of Tables

|    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |    |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| S1 | Summary of PDAC cohorts used in this study. $N$ denotes the number of samples after all quality-control filters (histology, survival validity, dataset-specific exclusions). Patient-level metadata: OS = overall survival (time and event indicator); PurIST = tumor subtype classification (Rashid et al. 2020); DeCAF = CAF subtype classification (Peng et al. 2020). Expression data and molecular classifications for the validation cohorts were originally compiled for the DeCAF study (Peng et al. 2020). . . . . | 15 |
| S2 | External validation C-index by method and cohort. DeSurv at BO-selected rank ( $k = 3$ ) is compared with standard NMF at matched rank ( $k = 3$ ), elbow-selected rank ( $k = 5$ ), and BO-selected rank ( $k = 7, \alpha = 0$ ). Higher values indicate better discrimination. . . . .                                                                                                                                                                                                                                    | 20 |
| S3 | Pooled external validation hazard ratios (per SD of linear predictor) for DeSurv at $k = 3$ and standard NMF at BO-selected $k = 7$ ( $\alpha = 0$ ). Adjusted models include PurIST and DeCAF molecular classifiers as covariates, with stratification by dataset. . . . .                                                                                                                                                                                                                                                 | 20 |
| S4 | Top 270 genes for DeSurv factor D1 (Factor D1 (Classical tumor + restCAF stroma)), ranked by factor specificity score $s_{ij}$ . The 270-gene cutoff corresponds to the Bayesian optimization–selected projection dimension ( $n_{\text{top}}$ ). . . . .                                                                                                                                                                                                                                                                   | 21 |
| S5 | Top 270 genes for DeSurv factor D2 (Factor D2 (proCAF stroma)), ranked by factor specificity score $s_{ij}$ . . . . .                                                                                                                                                                                                                                                                                                                                                                                                       | 22 |
| S6 | Top 270 genes for DeSurv factor D3 (Factor D3 (Basal-like tumor)), ranked by factor specificity score $s_{ij}$ . . . . .                                                                                                                                                                                                                                                                                                                                                                                                    | 23 |

## Part I

# Supplementary Methods

## 1 Model details

Let  $X \in \mathbb{R}_{\geq 0}^{p \times n}$  denote the nonnegative gene expression matrix with  $p$  genes and  $n$  subjects. DeSurv approximates  $X \approx WH$  with  $W \in \mathbb{R}_{\geq 0}^{p \times k}$  and  $H \in \mathbb{R}_{\geq 0}^{k \times n}$ , and links the shared program matrix  $W$  to survival via an elastic-net-penalized Cox model.

Let  $y \in \mathbb{R}_{> 0}^n$  be the observed survival times,  $\delta \in \{0, 1\}^n$  the event indicators, and  $Z = W^\top X \in \mathbb{R}^{k \times n}$  the sample-wise factor scores. We write  $Z_i$  for the  $i$ th column of  $Z$ , and  $n_{\text{event}} = \sum_{i=1}^n \delta_i$ . For convenience we recall the DeSurv loss from Equation 1 in the main text and rewrite it as

$$\begin{aligned} \mathcal{L}(W, H, \beta) = & \\ & (1 - \alpha) \left( \frac{1}{2np} \|X - WH\|_F^2 + \frac{\lambda_H}{2nk} \|H\|_F^2 \right) \\ & - \alpha \left( \frac{2}{n_{\text{event}}} \ell(W, \beta) - \lambda \{ \xi \|\beta\|_1 + \frac{1-\xi}{2} \|\beta\|_2^2 \} \right) \end{aligned}$$

where  $\|\cdot\|_F$  is the Frobenius norm and  $\beta \in \mathbb{R}^k$  are the Cox coefficients. The Cox log-partial likelihood  $\ell(W, \beta)$  is

$$\ell(W, \beta) = \sum_{i=1}^n \delta_i \left[ Z_i^\top \beta - \log \left( \sum_{j: y_j \geq y_i} \exp(Z_j^\top \beta) \right) \right]. \quad (1)$$

Hyperparameters satisfy  $\alpha \in [0, 1)$ ,  $\lambda_H \geq 0$ ,  $\lambda \geq 0$ , and  $\xi \in [0, 1]$ . The constants  $1/(2np)$ ,  $2/n_{\text{event}}$ , and  $1/(2nk)$  are chosen for numerical convenience and do not affect the minimizer.

## 2 Optimization Algorithm

### 2.1 Block coordinate descent scheme

Algorithm S1 summarizes the block coordinate descent (BCD) scheme used to minimize  $\mathcal{L}(W, H, \beta)$ . At each outer iteration, we update  $H$ , then  $W$ , then  $\beta$  while holding the other blocks fixed.

### 2.2 Update for $H$

Conditional on  $W$ , the loss  $\mathcal{L}(W, H, \beta)$  reduces to a strictly convex quadratic function of  $H$ . We adopt the standard NMF multiplicative update with  $\ell_2$  penalty:

$$H \leftarrow \max \left( H \odot \frac{W^\top X}{W^\top W H + \lambda_H H + \varepsilon_H}, \varepsilon_H \right), \quad (2)$$

where  $\odot$  denotes elementwise multiplication, all divisions are elementwise, and  $\varepsilon_H > 0$  is a small floor to prevent entries from becoming exactly zero. This update is equivalent to a majorization-minimization step and guarantees a nonincreasing reconstruction term conditional on  $W$  (Lee and Seung 2001; Pascual-Montano et al. 2006). The  $\varepsilon_H$ -floor ensures that iterates remain in the interior of the nonnegative orthant, which simplifies the convergence analysis.

---

**Algorithm S1** DeSurv block coordinate descent

---

**Input:**  $X \in \mathbb{R}_{\geq 0}^{p \times n}$ , survival times  $y \in \mathbb{R}_{\geq 0}^n$ , event indicators  $\delta \in \{0, 1\}^n$ , tolerance  $tol$ , maximum iterations  $maxit$ , supervision parameter  $\alpha$ , rank  $k$ , penalties  $\lambda_H$ ,  $\lambda$ , and  $\xi$

**Output:** Fitted DeSurv parameters  $(W, H, \beta)$

- 1: Initialize  $W^{(0)}, H^{(0)}$  with positive entries (e.g.,  $W^{(0)}, H^{(0)} \sim \text{Uniform}(0, \max X)$ )
  - 2: Initialize  $\beta^{(0)}$  (e.g.,  $\beta^{(0)} = \mathbf{1}$ )
  - 3:  $loss = \mathcal{L}(W^{(0)}, H^{(0)}, \beta^{(0)})$
  - 4:  $eps = \infty, t = 0$
  - 5: **while**  $eps \geq tol$  **and**  $t < maxit$  **do**
  - 6:   **(H-update)**
  - 7:     Update  $H^{(t+1)}$  from  $H^{(t)}$  using the multiplicative rule in Eq. 2,
  - 8:     holding  $W^{(t)}$  and  $\beta^{(t)}$  fixed.
  - 9:   **(W-update)**
  - 10:    Update  $W^{(t+1)}$  from  $W^{(t)}$  using the hybrid multiplicative rule in Eq. 6,
  - 11:    holding  $H^{(t+1)}$  and  $\beta^{(t)}$  fixed, and perform backtracking to ensure
  - 12:     $\mathcal{L}$  does not increase.
  - 13:   **( $\beta$ -update)**
  - 14:    Update  $\beta^{(t+1)}$  from  $\beta^{(t)}$  using a Newton-like step for the Cox loss
  - 15:    (Eq. 7), holding  $W^{(t+1)}$  and  $H^{(t+1)}$  fixed.
  - 16:     $lossNew = \mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t+1)})$
  - 17:     $eps = |lossNew - loss|/|loss|$
  - 18:     $loss = lossNew$
  - 19:     $t = t + 1$
  - 20: **end while**
  - 21: **return**  $\hat{W} = W^{(t+1)}, \hat{H} = H^{(t+1)}, \hat{\beta} = \beta^{(t+1)}$
- 

## 2.3 Update for $W$

For  $W$ , we construct a hybrid multiplicative update that balances the contributions of the NMF and Cox gradients. Let

$$\nabla_W \mathcal{L}_{\text{NMF}}(W, H) = \frac{1}{np} (WH - X)H^\top,$$

and let

$$\nabla_W \mathcal{L}_{\text{Cox}}(W, \beta) = \frac{2}{n_{\text{event}}} \nabla_W \ell(W, \beta).$$

where  $\nabla_W \ell(W, \beta)$  denotes the Cox gradient with respect to  $W$ . A closed-form expression for  $\nabla_W \ell$  is

$$\nabla_W \ell(W, \beta) = \sum_{i=1}^n \delta_i \left( x_i - \frac{\sum_{j: y_j \geq y_i} x_j \exp(x_j^\top W \beta)}{\sum_{j: y_j \geq y_i} \exp(x_j^\top W \beta)} \right) \beta^\top, \quad (3)$$

where  $x_i$  denotes the  $i$ th column of  $X$ .

We define the gradient-balancing factor

$$\zeta^{(t)} = \frac{\|(1 - \alpha) \nabla_W \mathcal{L}_{\text{NMF}}(W^{(t)}, H^{(t+1)})\|_F^2}{\|\alpha \nabla_W \mathcal{L}_{\text{Cox}}(W^{(t)}, \beta^{(t)})\|_F^2}, \quad (4)$$

and clip  $\zeta^{(t)}$  to avoid extreme values:  $\zeta^{(t)} \leftarrow \min(\zeta^{(t)}, \zeta_{\max})$  with  $\zeta_{\max} = 10^6$  in all experiments.

The multiplicative factor for  $W$  is then

$$R^{(t)} = \frac{\frac{(1-\alpha)}{np} X H^{(t+1)\top} + \frac{2\alpha}{n_{\text{event}}} \zeta^{(t)} \nabla_W \ell(W^{(t)}, \beta^{(t)})}{\frac{(1-\alpha)}{np} W^{(t)} H^{(t+1)} H^{(t+1)\top} + \varepsilon_W}, \quad (5)$$

and the proposed update is

$$W^{(t+1)} = \max\left(W^{(t)} \odot R^{(t)}, \varepsilon_W\right), \quad (6)$$

with a small floor  $\varepsilon_W > 0$ . When  $\alpha = 0$ , this reduces to the standard multiplicative update for NMF (Lee and Seung 2001); for  $\alpha > 0$ , the Cox gradient perturbs the update in a direction that decreases the supervised loss.

Because the combined multiplicative step does not, by itself, guarantee a nonincrease in the full loss  $\mathcal{L}$ , we embed it in a backtracking line search.

---

**Algorithm S2**  $W$  update with backtracking

---

**Input:** Value of  $W$  and the previous iteration  $t$ :  $W^{(t)}$ , the multiplicative update term at the current iteration  $R^{(t+1)}$ , the max number of backtracking updates  $max\_bt$ , the backtracking parameter  $\rho \in (0, 1)$

**Output:**  $W^{(t+1)}$

```

1:  $\theta = 1$ 
2:  $b = 1$ 
3:  $flag\_accept = FALSE$ 
4: while  $b \leq max\_bt$  do
5:    $W_{cand}^{(t+1)} = W^{(t)} \odot [(1 - \theta) + \theta R^{(t+1)}]$ 
6:   (Column normalization)
7:     Compute  $D = \text{diag}(\|W_{(:,1)cand}^{(t+1)}\|_2, \dots, \|W_{(:,k)cand}^{(t+1)}\|_2)$ 
8:     and set  $W_{cand}^{(t+1)} \leftarrow W_{cand}^{(t+1)} D^{-1}$ ,  $H_{cand}^{(t+1)} \leftarrow D H^{(t+1)}$ , and  $\beta_{cand}^{(t)} \leftarrow D \beta^{(t)}$ 
9:     if  $\mathcal{L}(W_{cand}^{(t+1)}, H_{cand}^{(t+1)}, \beta^{(t)}) \leq \mathcal{L}(W^{(t)}, H^{(t+1)}, \beta^{(t)})$  then
10:       $W^{(t+1)} = W_{cand}^{(t+1)}$ 
11:       $H^{(t+1)} = H_{cand}^{(t+1)}$ 
12:       $\beta^{(t)} = \beta_{cand}^{(t)}$ 
13:       $flag\_accept = TRUE$ 
14:      break
15:    end if
16:     $\theta = \theta * \rho$ 
17:     $b = b + 1$ 
18: end while
19: if  $flag\_accept = FALSE$  then
20:    $W^{(t+1)} = W^{(t)}$ 
21: end if

```

---

The column normalization preserves both  $WH$  and  $W\beta$ , and therefore leaves the loss  $\mathcal{L}$  invariant up to numerical error. Backtracking guarantees that the accepted  $W$  update does not increase  $\mathcal{L}$ .

## 2.4 Update for $\beta$

Conditional on  $(W, H)$ , the loss in  $\beta$  reduces to a convex elastic-net-penalized Cox problem:

$$\min_{\beta \in \mathbb{R}^k} \left\{ -\frac{2\alpha}{n_{\text{event}}} \ell(W, \beta) + \lambda \left( \xi \|\beta\|_1 + \frac{1-\xi}{2} \|\beta\|_2^2 \right) \right\}.$$

We solve this subproblem by cyclic coordinate descent following (Simon et al. 2011). Writing  $\ell(\beta) = \ell(W, \beta)$  and  $\tilde{\eta}_i = Z_i^\top \beta$ , the update for coordinate  $r$  has the closed form

$$\hat{\beta}_r = \frac{S\left(\frac{1}{n} \sum_{i=1}^n w(\tilde{\eta})_i z_{i,r} \left[ v(\tilde{\eta})_i - \sum_{j \neq r} z_{i,j} \beta_j \right], \lambda \xi\right)}{\frac{1}{n} \sum_{i=1}^n w(\tilde{\eta})_i z_{i,r}^2 + \lambda(1 - \xi)}, \quad (7)$$



where  $S(a, \tau) = \text{sign}(a) \max(|a| - \tau, 0)$  is the soft-thresholding operator, and  $w(\tilde{\eta})$ ,  $v(\tilde{\eta})$  are standard local quadratic approximations of the Cox log-partial likelihood (Simon et al. 2011). We iterate coordinate updates until the subproblem converges to numerical tolerance, yielding a minimizer in  $\beta$  for the current  $W$ .

## 2.5 Normalization of $W$

After each accepted  $W$  update, we normalize the columns of  $W$  as  $W \leftarrow WD^{-1}$ , and adjust  $H$  and  $\beta$  as

$$H \leftarrow DH, \quad \beta \leftarrow D\beta,$$

where  $D$  is the diagonal matrix of column  $\ell_2$ -norms of  $W$ . This preserves the reconstruction  $WH$  and the linear predictor  $W\beta$ , leaving both the NMF and Cox terms in the loss unchanged. Normalization prevents degeneracy in the scale-nonidentifiable factorization and keeps columns of  $W$  comparable in magnitude and interpretable as gene programs.

## 2.6 Derivation of $W$ update from projected coordinate descent

The  $W$  update can be derived directly from the projected coordinate descent update with a specific step size.

Recall that the overall loss function is

$$\mathcal{L}(W, H, \beta) = \frac{(1-\alpha)}{2np} \|X - WH\|_F^2 - \frac{2\alpha}{n_{event}} \ell(W, \beta) + \lambda \left( \xi \|\beta\|_1 + \frac{(1-\xi)}{2} \|\beta\|_2^2 \right).$$

where  $\ell(W, \beta)$  is the log-partial likelihood for the Cox model. Let  $\nabla_W \ell$  represent the derivative of  $\ell(W, \beta)$  with respect to  $W$ . Then the derivative of the overall loss with respect to  $W$  is

$$\frac{\partial L}{\partial W} = \frac{(1-\alpha)}{np} (WHH^T - XH^T) - \frac{2\alpha}{n_{event}} \nabla_W \ell.$$

Then the gradient descent update rule at iteration  $t$  is

$$\begin{aligned} W^{(t)} &= W^{(t-1)} - \gamma \left( \frac{\partial L}{\partial W} \right) \\ &= W^{(t-1)} - \gamma \left( \frac{(1-\alpha)}{np} (WHH^T - XH^T) - \frac{2\alpha}{n_{event}} \nabla_W \ell \right). \end{aligned}$$

Let the step size  $\gamma$  be defined as in (Lee and Seung 2001):

$$\gamma = \frac{np}{(1-\alpha)} \frac{W}{WHH^T}.$$

Then the update becomes

$$\begin{aligned} W^{(t)} &= W^{(t-1)} - \frac{np}{(1-\alpha)} \frac{W}{WHH^T} \left( \frac{(1-\alpha)}{np} (WHH^T - XH^T) - \frac{2\alpha}{n_{event}} \nabla_W \ell \right) \\ &= \frac{W}{WHH^T} XH^T + \frac{2\alpha np}{n_{event}(1-\alpha)} \nabla_W \ell \\ &= W \odot \frac{XH^T + \frac{2\alpha np}{n_{event}(1-\alpha)} \nabla_W \ell}{WHH^T} \\ &= W \odot \frac{\frac{(1-\alpha)}{np} XH^T + \frac{2\alpha}{n_{event}} \nabla_W \ell}{\frac{(1-\alpha)}{np} WHH^T}. \end{aligned}$$

Finally, projected coordinate descent projects the  $W$  update in to the positive space

$$W^{(t)} = \max \left( W \odot \frac{\frac{(1-\alpha)}{np} XH^T + \frac{2\alpha}{n_{event}} \nabla_W \ell}{\frac{(1-\alpha)}{np} WHH^T}, 0 \right).$$

Since  $W \geq 0$  this is equivalent to

$$W^{(t)} = W \odot \max \left( \frac{\frac{(1-\alpha)}{np} XH^T + \frac{2\alpha}{n_{event}} \nabla_W \ell}{\frac{(1-\alpha)}{np} WHH^T}, 0 \right),$$

which matches the multiplicative form used in the software implementation.

### 3 Convergence proof

We show that, under mild regularity conditions, the block coordinate descent (BCD) Algorithm S1 converges to a stationary point of the DeSurv loss function

$$\mathcal{L}(W, H, \beta).$$

For clarity, we first analyze the algorithm *without* the column normalization of  $W$  inside the loop, and then argue in Section 3.1 that the normalization step preserves stationarity of limit points.

Throughout, let  $\theta = (W, H, \beta)$  and denote  $\mathcal{L}(\theta) = \mathcal{L}(W, H, \beta)$ . The feasible set is

$$\Theta = \{(W, H, \beta) : W \in \mathbb{R}_{\geq 0}^{p \times k}, H \in \mathbb{R}_{\geq 0}^{k \times n}, \beta \in \mathbb{R}^k\}.$$

**Assumption 1 (Regularity and parameter space).** We assume:

- (i) The data  $X \in \mathbb{R}_{\geq 0}^{p \times n}$ ,  $y \in \mathbb{R}_{> 0}^n$ , and  $\delta \in \{0, 1\}^n$  are fixed and bounded.
- (ii) The hyperparameters satisfy  $\lambda_H > 0$ ,  $\lambda > 0$ ,  $\alpha \in [0, 1)$ , and  $\xi \in [0, 1]$ .
- (iii) The initial iterate  $\theta^{(0)} = (W^{(0)}, H^{(0)}, \beta^{(0)})$  lies in  $\Theta$  and satisfies  $W^{(0)}, H^{(0)} > 0$  elementwise.

**Assumption 2 (Block updates).** At each outer iteration  $t$ , the three block updates in Algorithm S1 satisfy:

- (i) *H-update.* For fixed  $(W^{(t)}, \beta^{(t)})$ , the update  $H^{(t)} \mapsto H^{(t+1)}$  is given by the multiplicative rule

$$H \leftarrow H \odot \frac{W^\top X}{W^\top W H + \lambda_H H},$$

applied at  $(W^{(t)}, H^{(t)})$ . This rule preserves nonnegativity and yields a nonincreasing value of the reconstruction term conditional on  $W^{(t)}$  and  $\beta^{(t)}$ ; see e.g. (Lee and Seung 2001; Pascual-Montano et al. 2006; Lin 2007).

- (ii) *W-update.* For fixed  $(H^{(t+1)}, \beta^{(t)})$ , the update  $W^{(t)} \mapsto W^{(t+1)}$  is the hybrid multiplicative step followed by backtracking as in Algorithm S2, producing  $W^{(t+1)}$  such that

$$\mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t)}) \leq \mathcal{L}(W^{(t)}, H^{(t+1)}, \beta^{(t)}).$$

If no candidate satisfies the Armijo-type condition within the allowed number of backtracking steps, the algorithm sets  $W^{(t+1)} := W^{(t)}$ .

- (iii)  $\beta$ -update. For fixed  $(W^{(t+1)}, H^{(t+1)})$ , the update  $\beta^{(t)} \mapsto \beta^{(t+1)}$  is obtained by running the coordinate descent method of Simon et al. (2011) on the convex elastic-net Cox subproblem until convergence. Thus

$$\beta^{(t+1)} \in \arg \min_{\beta \in \mathbb{R}^k} \mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta),$$

and

$$\mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t+1)}) \leq \mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t)}).$$

**Lemma 1 (Continuity and bounded level sets).** Under Assumption 1,  $\mathcal{L} : \Theta \rightarrow \mathbb{R}$  is continuous, and the initial sublevel set

$$\mathcal{S}_0 := \{\theta \in \Theta : \mathcal{L}(\theta) \leq \mathcal{L}(\theta^{(0)})\}$$

is nonempty, closed, and bounded.

*Proof.* The NMF term  $\|X - WH\|_F^2$  is a polynomial in the entries of  $W$  and  $H$ , hence is continuously differentiable. The penalty  $\|H\|_F^2$  is continuous as well. The Cox partial log-likelihood is a smooth function of the linear predictor  $\eta_i = x_i^\top W\beta$ , which is linear in  $(W, \beta)$ , so this term is continuously differentiable in  $(W, \beta)$ . The  $\ell_2$  penalty on  $\beta$  is smooth, and the  $\ell_1$  penalty is convex and lower semicontinuous. Therefore  $\mathcal{L}$  is continuous on  $\Theta$ .

The constraints  $W, H \geq 0$  define closed convex cones, and  $\beta \in \mathbb{R}^k$  is unconstrained. Because  $\lambda_H > 0$  and  $\lambda > 0$ , the terms  $\frac{\lambda_H}{nk} \|H\|_F^2$  and  $\lambda \frac{1-\xi}{2} \|\beta\|_2^2$  dominate the objective as  $\|H\|_F \rightarrow \infty$  or  $\|\beta\|_2 \rightarrow \infty$ , respectively. Thus the sublevel set  $\mathcal{S}_0$  is bounded in  $(H, \beta)$ . Since  $W \geq 0$  and  $X$  is fixed, the reconstruction term and Cox term being bounded prevent  $W$  from diverging while  $\mathcal{L}$  remains below  $\mathcal{L}(\theta^{(0)})$ , so  $W$  is bounded on  $\mathcal{S}_0$ . Because  $\Theta$  is closed and  $\mathcal{L}$  is continuous,  $\mathcal{S}_0$  is closed. Nonemptiness follows from  $\theta^{(0)} \in \mathcal{S}_0$ .  $\square$

**Lemma 2 (Monotone descent and existence of limit points).** Under Assumptions 1 and 2, Algorithm S1 (without  $W$  normalization) generates a sequence  $\{\theta^{(t)}\}_{t \geq 0} \subset \mathcal{S}_0$  such that

$$\mathcal{L}(\theta^{(t+1)}) \leq \mathcal{L}(\theta^{(t)}) \quad \forall t,$$

and  $\{\mathcal{L}(\theta^{(t)})\}$  converges to a finite limit  $\mathcal{L}^*$ . Moreover,  $\{\theta^{(t)}\}$  is bounded and therefore admits at least one limit point.

*Proof.* By Assumption 2(i),

$$\mathcal{L}(W^{(t)}, H^{(t+1)}, \beta^{(t)}) \leq \mathcal{L}(W^{(t)}, H^{(t)}, \beta^{(t)}).$$

By Assumption 2(ii),

$$\mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t)}) \leq \mathcal{L}(W^{(t)}, H^{(t+1)}, \beta^{(t)}).$$

By Assumption 2(iii),

$$\mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t+1)}) \leq \mathcal{L}(W^{(t+1)}, H^{(t+1)}, \beta^{(t)}).$$

Combining these inequalities gives

$$\mathcal{L}(\theta^{(t+1)}) \leq \mathcal{L}(\theta^{(t)}) \quad \forall t,$$

so  $\{\mathcal{L}(\theta^{(t)})\}$  is monotonically nonincreasing. Since  $\mathcal{L}$  is bounded below on  $\Theta$ , the sequence converges to some finite  $\mathcal{L}^*$ .

Each iterate lies in  $\mathcal{S}_0$  by construction, and  $\mathcal{S}_0$  is bounded by Lemma 1. Therefore  $\{\theta^{(t)}\}$  is bounded and has at least one limit point by the Bolzano–Weierstrass theorem.  $\square$

**Lemma 3 (Blockwise optimality of limit points).** Let  $\theta^* = (W^*, H^*, \beta^*)$  be any limit point of  $\{\theta^{(t)}\}$  under Assumptions 1–2. Then:

- (i)  $H^*$  satisfies the KKT conditions for

$$\min_{H \geq 0} \mathcal{L}(W^*, H, \beta^*).$$

(ii)  $W^*$  satisfies the KKT conditions for

$$\min_{W \geq 0} \mathcal{L}(W, H^*, \beta^*).$$

(iii)  $\beta^*$  is the unique minimizer of

$$\min_{\beta \in \mathbb{R}^k} \mathcal{L}(W^*, H^*, \beta),$$

and satisfies the KKT conditions for the elastic-net Cox subproblem.

*Proof.* Let  $\{t_j\}$  be a subsequence such that  $\theta^{(t_j)} \rightarrow \theta^* = (W^*, H^*, \beta^*)$ . By continuity of  $\mathcal{L}$ ,  $\mathcal{L}(\theta^{(t_j)}) \rightarrow \mathcal{L}^*$ .

(i) Conditional on  $(W, \beta)$ , the  $H$ -subproblem is

$$\min_{H \geq 0} \frac{(1 - \alpha)}{np} (\|X - WH\|_F^2 + \frac{\lambda_H}{nk} \|H\|_F^2) + \text{const in } H,$$

a strictly convex quadratic program. The multiplicative update for  $H$  is known to be a descent method whose fixed points coincide with the KKT points of this constrained subproblem (Lee and Seung 2001; Lin 2007). Since the full objective  $\mathcal{L}(\theta^{(t)})$  converges, the per-iteration decrease in  $\mathcal{L}$  due to the  $H$ -update must vanish along  $\{t_j\}$ . If  $H^*$  were not KKT for the  $H$ -subproblem given  $(W^*, \beta^*)$ , there would exist a feasible descent direction for  $H$  at  $(W^*, \beta^*)$ , implying that for sufficiently large  $j$  the  $H$ -update would still produce a strict decrease in  $\mathcal{L}$ , contradicting convergence. Hence  $H^*$  satisfies the KKT conditions.

(ii) Conditional on  $(H, \beta)$ , the  $W$ -subproblem is differentiable and convex in  $W$  (sum of a quadratic term and a negative Cox partial log-likelihood in a linear predictor). The hybrid multiplicative step with backtracking can be viewed as a projected gradient-like update onto the nonnegative orthant: as shown in the derivation above, the multiplicative direction is equivalent to a negative gradient step with the adaptive step size of Seung et al., projected onto the nonnegative orthant. Under mild conditions on the search direction and step sizes, any limit point of such a projected gradient scheme is a KKT point for the constrained subproblem; see, for example, (Tseng 2001; Grippo and Sciandrone 2000). If  $W^*$  did not satisfy the KKT conditions, there would be a feasible descent direction at  $(W^*, H^*, \beta^*)$  and a sufficiently small step that reduces  $\mathcal{L}$ , which the line search would eventually accept. This would contradict the fact that  $\mathcal{L}(\theta^{(t_j)}) \rightarrow \mathcal{L}^*$ . Therefore  $W^*$  is KKT for the  $W$ -subproblem.

(iii) For fixed  $(W, H)$ , the  $\beta$ -subproblem is the convex elastic-net penalized Cox objective. The coordinate descent algorithm converges to the unique minimizer. By Assumption 2(iii),  $\beta^{(t+1)}$  is taken to be this minimizer at each iteration. Passing to the limit along  $\{t_j\}$  shows that  $\beta^*$  is the unique minimizer of the  $\beta$ -block given  $(W^*, H^*)$  and hence satisfies its KKT conditions.  $\square$

**Theorem 1 (Convergence to a stationary point).** Under Assumptions 1 and 2, the sequence  $\{\theta^{(t)}\}$  produced by Algorithm S1 (without  $W$  normalization) satisfies:

- (a) The objective values  $\{\mathcal{L}(\theta^{(t)})\}$  are monotonically nonincreasing and converge to a finite limit  $\mathcal{L}^*$ .
- (b) Every limit point  $\theta^* = (W^*, H^*, \beta^*)$  of  $\{\theta^{(t)}\}$  is a stationary point of  $\mathcal{L}$  on  $\Theta$ , i.e. it satisfies the first-order KKT conditions for

$$\min_{\theta \in \Theta} \mathcal{L}(\theta).$$

*Proof.* Part (a) is Lemma 2. For part (b), Lemma 3 shows that any limit point  $\theta^*$  is blockwise optimal: each block is optimal (in the KKT sense) when the others are held fixed.

The DeSurv loss can be written as

$$\mathcal{L}(\theta) = f(W, H, \beta) + g(\beta) + I_{\{W \geq 0\}}(W) + I_{\{H \geq 0\}}(H),$$

where  $f$  is continuously differentiable,  $g(\beta) = \lambda \xi \|\beta\|_1$  is a separable convex (possibly nondifferentiable) penalty, and  $I_C$  denotes the indicator of a closed convex set  $C$ . This is the smooth+nonsmooth composite form considered in block coordinate descent analyses such as (Tseng 2001). In this setting, any point that is optimal with respect to each block individually (a block coordinatewise minimizer) is a stationary point for the full problem: equivalently,

$$\begin{aligned} 0 &\in \nabla_{W,H} f(W^*, H^*, \beta^*) + \partial I_{\{W \geq 0\}}(W^*) + \partial I_{\{H \geq 0\}}(H^*), \\ 0 &\in \nabla_{\beta} f(W^*, H^*, \beta^*) + \partial g(\beta^*), \end{aligned}$$

which are precisely the KKT conditions for  $\min_{\theta \in \Theta} \mathcal{L}(\theta)$ . Hence every limit point of Algorithm S1 is stationary.  $\square$

### 3.1 Effect of column normalization of $W$

The above analysis omits the column-normalization step for  $W$  used in Algorithm S2. We briefly argue that this normalization does not affect stationarity of limit points.

Let  $D$  be a diagonal matrix with strictly positive diagonal entries. The transformation

$$(W, H, \beta) \mapsto (W', H', \beta') = (WD^{-1}, DH, D\beta)$$

preserves both the product  $WH$  and the linear predictor  $W\beta$ , and hence leaves  $\mathcal{L}$  invariant. The column-normalization step is exactly such a transformation, with  $D$  chosen from the column norms of  $W$ .

Let  $\{\theta^{(t)}\}$  be the sequence generated by the algorithm without normalization, and let  $\{\tilde{\theta}^{(t)}\}$  be the sequence with normalization. For each  $t$  there exists a diagonal  $D^{(t)}$  with strictly positive entries such that

$$\tilde{\theta}^{(t)} = (W^{(t)} D^{(t), -1}, D^{(t)} H^{(t)}, D^{(t)} \beta^{(t)}),$$

and  $\mathcal{L}(\tilde{\theta}^{(t)}) = \mathcal{L}(\theta^{(t)})$ . Thus limit points of  $\{\tilde{\theta}^{(t)}\}$  are obtained from limit points of  $\{\theta^{(t)}\}$  by such invertible diagonal scalings.

Since the KKT conditions are expressed in terms of the gradients with respect to  $WH$  and  $W\beta$ , and these quantities are invariant under the above scaling, stationarity is preserved under the transformation. Therefore Theorem 1 implies that every limit point of the *normalized* algorithm is also a stationary point of  $\mathcal{L}$  (up to this scaling equivalence), which establishes convergence of the implementation used in practice.

### 3.2 Remark (Coxnet implementation)

Our implementation updates the  $\beta$  block using a Coxnet-style coordinate descent that relies on an approximate Hessian. Consequently, the formal optimality proof is established for an idealized version of the  $\beta$  update that assumes exact second-order information. Nonetheless, in empirical applications the approximate update is numerically stable and yields a monotone decrease in the objective, consistent with the behavior predicted by the idealized analysis.

## 4 Supervision on $W$ versus $H$

Classical nonnegative matrix factorization (NMF) is non-identifiable: for any invertible matrix  $R$  with nonnegative entries, the factorization  $(W, H)$  can be transformed to  $(WR, R^{-1}H)$  without changing the reconstruction error, yielding multiple valid solutions (Donoho and Stodden 2004; Laurberg et al. 2008; Gillis 2014). Although normalizing columns of  $W$  or rows of  $H$  resolves the trivial scaling ambiguity, nonnegative mixing transformations persist, and empirical studies consistently show that the sample-loading matrix  $H$  is

more sensitive to initialization and local minima than the program matrix  $W$  (Brunet et al. 2004; Kim and Park 2008; Gillis and Glineur 2012).

In DeSurv, survival supervision enters through the projection  $Z = W^\top X$  in the partial log-likelihood, so that the gradient of the DeSurv loss with respect to the programs is

$$\nabla \mathcal{L} = (1 - \alpha) \nabla_W \mathcal{L}_{\text{NMF}}(W, H) - \alpha \nabla_W \mathcal{L}_{\text{Cox}}(W, \beta),$$

and therefore the update rule for  $W$  as in Equation 6 relies on both the reconstruction term and the supervision term. As such, the gene-program definitions are directly shaped by their association with survival. This mechanism amplifies genes aligned with the hazard gradient and suppresses those that dilute prognostic structure, ensuring that the learned programs encode survival-relevant biological variation.

By contrast, if supervision were applied to the sample loadings  $H$ , the model could reduce the Cox loss by redistributing patient-specific coefficients while leaving the gene-level programs  $W$  nearly unchanged, a behavior documented in multiple supervised variants of NMF and supervised topic models, where supervision applied only to the coefficient matrix modifies  $H$  but not the basis  $W$  (Cai et al. 2011; Blei and McAuliffe 2007).

Supervising  $W$  also provides a practical advantage for **portability**: because  $W$  defines gene-level programs, new samples can be embedded by the closed-form projection  $Z_{\text{new}} = X_{\text{new}}^\top W$ . In contrast, supervising  $H$  would not yield such a mapping; instead, obtaining sample loadings for external datasets would require solving a nonnegative least-squares (NNLS) problem for each new sample, introducing optimization noise and reducing reproducibility.

Together, these considerations motivate supervising through  $W$ , which shapes the gene programs toward prognostic directions, stabilizes solutions across initializations, and yields biologically interpretable signatures that generalize across cohorts.

## 5 Cross-validation procedure

Algorithm S3 describes the cross validation procedure with  $F$  folds and  $R$  initializations. We define the c-index using comparable pairs  $\mathcal{P} = \{(i, j) : y_i < y_j, \delta_i = 1\}$  and linear predictors  $\hat{\eta}_i$ :

$$\hat{c} = \frac{1}{|\mathcal{P}|} \sum_{(i,j) \in \mathcal{P}} [\mathbb{1}(\hat{\eta}_i > \hat{\eta}_j) + \frac{1}{2} \mathbb{1}(\hat{\eta}_i = \hat{\eta}_j)]. \quad (8)$$

---

**Algorithm S3** Cross-validation for DeSurv pipeline

---

**Input:**

Number of folds  $F$   
Number of initializations  $R$   
Observed data  $(X, y, \delta)$   
Hyperparameters  $k, \alpha, \lambda, \xi$ , and  $\lambda_H$

**Output:** The cross-validated c-index

```
1: Divide subjects into  $F$  folds.
2: for  $f = 1$  to  $F$  do
3:   Split data into training and validation  $(X_{(-f)}, y_{(-f)}, \delta_{(-f)})$ , and  $(X_{(f)}, y_{(f)}, \delta_{(f)})$  sets
4:   for  $r = 1$  to  $R$  do
5:     set.seed( $r$ )
6:     Apply Algorithm S1 with inputs  $(X_{(-f)}, y_{(-f)}, \delta_{(-f)})$ , and  $(k, \alpha, \lambda, \xi, \lambda_H)$ 
7:     Obtain  $\hat{W}$  and  $\hat{\beta}$  as output from Algorithm S1
8:     Compute the estimated linear predictor:  $\hat{\eta} = X_{(f)}^T \hat{W} \hat{\beta}$ 
9:     Compute c-index, denoted  $\hat{c}_{(f)r}$ , according to Equation 8
10:   end for
11: end loop over  $r$ 
12:   Compute average c-index across initializations:  $\hat{c}_{(f)} = \frac{1}{R} \sum_{r=1}^R \hat{c}_{(f)r}$ 
13: end for
14: end loop over  $f$ 
15: Compute final cross-validated c-index:  $\hat{c} = \frac{1}{F} \sum_{f=1}^F \hat{c}_{(f)}$ 
16: return Final estimate  $\hat{c}$ 
```

---

## 6 Bayesian Optimization

We treat hyperparameter tuning as a black-box optimization problem over a bounded domain. Let  $\theta$  collect the tuned parameters (at minimum  $\theta = (k, \alpha, \lambda, \xi)$ , and optionally discrete choices such as the signature size  $n_{\text{top}}$ ). For a given  $\theta$  we run the cross-validation procedure in Algorithm S3, averaging the c-index across folds and random initializations to form the objective

$$\hat{c}(\theta) = \frac{1}{F} \sum_{f=1}^F \hat{c}_{(f)}(\theta).$$

Because  $\hat{c}(\theta)$  is expensive to evaluate and non-differentiable with respect to  $\theta$ , we use Bayesian optimization (BO) to adaptively select candidate configurations.

We place a Gaussian-process (GP) prior on the response surface  $f(\theta) \sim \mathcal{GP}(0, k(\theta, \theta'))$  with a Mat'ern kernel and automatic relevance determination length-scales. Continuous parameters are rescaled to  $[0, 1]$  and penalties are optimized on a log scale; integer-valued coordinates (e.g.,  $k$  or  $n_{\text{top}}$ ) are proposed in the continuous space and rounded inside the evaluator. After an initial batch of evaluations, the GP is refit after each new observation and used to guide acquisition.

At iteration  $t$ , let  $\mu_t(\theta)$  and  $\sigma_t(\theta)$  denote the GP posterior mean and standard deviation, and let  $f_\star$  be the best observed score. We select new candidates by maximizing expected improvement (EI),

$$\text{EI}(\theta) = (\mu_t(\theta) - f_\star - \kappa) \Phi(Z(\theta)) + \sigma_t(\theta) \phi(Z(\theta)), \quad Z(\theta) = \frac{\mu_t(\theta) - f_\star - \kappa}{\sigma_t(\theta)},$$

where  $\Phi$  and  $\phi$  are the standard normal cdf/pdf and  $\kappa \geq 0$  controls the exploration margin. EI is evaluated over a candidate pool sampled from the bounded search region, and the maximizer is evaluated next. The loop continues until the iteration budget is exhausted or the improvement plateaus.

For model selection, we take the highest observed  $\hat{c}(\theta)$  as the optimal configuration. When fold-level standard errors are available, we apply a one-standard-error rule for  $k$ , selecting the smallest rank whose mean

performance lies within one standard error of the best. The final DeSurv model is refit at the selected parameters using the consensus-based initialization described above, and the BO-chosen  $n_{\text{top}}$  (when tuned) is used to truncate each program for downstream analysis.

The parameter  $n_{\text{top}}$  controls which genes enter the survival linear predictor. After factorization, each sample’s score for factor  $j$  is computed as  $\widetilde{W}_j^\top x$ , where  $\widetilde{W}_j$  retains only the  $n_{\text{top}}$  genes with the largest factor-specificity scores  $s_{ij}$  (SI Appendix, Section~9); genes outside this set still shape the factorization but do not enter the linear predictor. BO selects  $n_{\text{top}}$  jointly with  $k$ ,  $\alpha$ ,  $\lambda$ , and  $\xi$  by optimizing cross-validated C-index. When  $\alpha = 0$  (no survival supervision), gene selection and survival outcomes are decoupled, so varying  $n_{\text{top}}$  has negligible effect on precision in the null simulation.

## 7 PDAC Datasets

We analyzed PDAC cohorts from TCGA-PAAD and CPTAC-PDAC as training data (Raphael et al. 2017; Ellis et al. 2013). External validation used independent cohorts from Dijk, Moffitt, PACA-AU (array and RNA-seq), and Puleo (Dijk et al. 2020; Moffitt et al. 2015; Zhao et al. 2018; Puleo et al. 2018). Training models were fit on the combined TCGA and CPTAC expression matrices after harmonizing gene identifiers; validation datasets were processed with the same gene set and transformations before prediction and clustering.

Dataset-specific inclusion and preprocessing steps were as follows. For TCGA-PAAD, we excluded samples lacking tumor grade and mapped features to gene symbols. For CPTAC, we retained samples annotated as PDAC by histology. For Dijk, all 90 samples passed quality-control filters and were retained. For Moffitt, only primary tumor specimens were kept. For PACA-AU array and RNA-seq, we retained primary PDAC tumors and collapsed duplicated gene symbols by median; the RNA-seq cohort received a “\_seq” suffix on sample IDs to avoid collisions with the array cohort. Puleo required no additional dataset-specific filtering beyond survival QC.

Across cohorts, samples without valid survival outcomes (nonpositive time, missing time, or missing event indicator) were removed. Expression matrices were analyzed on a log scale: RNA-seq cohorts as  $\log_2(\text{TPM} + 1)$ ; microarray cohorts in platform-normalized log-scale intensities. For each training cohort (TCGA-PAAD and CPTAC) separately, genes were ranked by mean expression and by variance independently; the 3,000 genes with the lowest sum of these two ranks were retained, selecting genes that are both highly expressed and highly variable. The intersection of these two gene sets yielded 1,970 genes used for model training. Validation datasets were restricted to this same 1,970-gene set. Within-sample rank transformation was applied to all datasets to mitigate platform effects, with validation data transformed identically to training.

Table S1 summarizes the cohorts, platforms, sample sizes after quality control, and patient-level metadata used in this study. Overall survival time and event indicators were available for all cohorts. PurIST (Rashid et al. 2020) and DeCAF (Peng et al. 2026) molecular classifications, applied to bulk expression data as described in those references, were used as covariates in the covariate-adjusted external validation analysis. Gene expression data and molecular classifications for the validation cohorts were originally compiled for the DeCAF study (Peng et al. 2026); training data were obtained directly from GDC.

## 8 Survival Analysis

Survival analyses used overall survival time with event indicators as provided by each cohort. For fitted DeSurv models, sample-level program scores were computed as  $Z = \widetilde{W}^\top X$  and standardized using the training mean and standard deviation before constructing the linear predictor  $\hat{\eta} = Z\hat{\beta}$ . Prognostic performance was summarized using the concordance index as defined above. For visualization and group-level testing, we generated Kaplan–Meier curves and log-rank tests for risk groups defined by median splits of  $\hat{\eta}$  within each dataset to avoid cross-cohort scaling differences. When multiple datasets were pooled, Cox models were



Table S1: Summary of PDAC cohorts used in this study.  $N$  denotes the number of samples after all quality-control filters (histology, survival validity, dataset-specific exclusions). Patient-level metadata: OS = overall survival (time and event indicator); PurIST = tumor subtype classification (Rashid et al. 2020); DeCAF = CAF subtype classification (Peng et al. 2020). Expression data and molecular classifications for the validation cohorts were originally compiled for the DeCAF study (Peng et al. 2020).

| Cohort            | Platform   | $N$ | Role       | Accession       | Metadata          | Reference             |
|-------------------|------------|-----|------------|-----------------|-------------------|-----------------------|
| TCGA-PAAD         | RNA-seq    | 144 | Training   | GDC (TCGA-PAAD) | OS, PurIST, DeCAF | Raphael et al. (2017) |
| CPTAC             | RNA-seq    | 129 | Training   | GDC (CPTAC-3)   | OS, PurIST, DeCAF | Ellis et al. (2013)   |
| Dijk              | RNA-seq    | 90  | Validation | E-MTAB-6830     | OS, PurIST, DeCAF | Dijk et al. (2020)    |
| Moffitt           | Microarray | 123 | Validation | GSE71729        | OS, PurIST, DeCAF | Moffitt et al. (2015) |
| PACA-AU (array)   | Microarray | 63  | Validation | GSE36924        | OS, PurIST, DeCAF | Zhao et al. (2018)    |
| PACA-AU (RNA-seq) | RNA-seq    | 52  | Validation | EGAS00001000154 | OS, PurIST, DeCAF | Zhao et al. (2018)    |
| Puleo             | Microarray | 288 | Validation | E-MTAB-6134     | OS, PurIST, DeCAF | Puleo et al. (2018)   |
| Training total    |            | 273 |            |                 |                   |                       |
| Validation total  |            | 616 |            |                 |                   |                       |

stratified by dataset to allow cohort-specific baseline hazards. All p-values reported for survival differences between groups correspond to two-sided log-rank tests.

## 9 Gene Program Correlation Analysis

To assess the biological identity of learned factors, we evaluated the correspondence between factor-specific gene rankings and established PDAC gene programs. For each factor column  $j$  of the gene program matrix  $W$ , we identified the top 50 genes by factor specificity score. Each column of  $W$  was first scaled by its column maximum to produce  $W_{ij}^* = W_{ij} / \max_i W_{ij}$ , placing all factors on a common  $[0, 1]$  scale. The specificity score for gene  $i$  in factor  $j$  was then defined as

$$s_{ij} = W_{ij}^* - \max_{j' \neq j} W_{ij'}^*.$$

Genes were ranked in descending order of  $s_{ij}$ , and the top 50 per factor were retained. The union of these gene sets across all factors was then intersected with the genes present in any reference program to form the set of genes used in the correlation analysis.

For each factor  $j$  and each reference gene program  $G_k$ , we computed the Spearman rank correlation between the continuous factor loadings  $W_{.j}$  (restricted to the intersection gene set) and a binary indicator vector  $\mathbf{1}[\text{gene} \in G_k]$ . P-values were derived from the asymptotic approximation to the Spearman test statistic and adjusted for multiple comparisons using the Benjamini–Hochberg procedure applied within each factor column independently. Gene programs were included in the visualization only if at least one factor yielded a Spearman correlation exceeding 0.2. Asterisks (\*) indicate adjusted p-value  $< 0.1$ .

Reference gene programs included Classical and Basal-like tumor cell signatures from Moffitt et al. (Moffitt et al. 2015); iCAF and myCAF cancer-associated fibroblast signatures from Elyada et al. (Elyada et al. 2019); promoting (proCAF) and restraining (restCAF) cancer-associated fibroblast subtypes from the DeCAF study (Peng et al. 2026); proCAF and restCAF subtypes from SCISSORS (Leary et al. 2023); tumor, stromal, and immune compartment signatures from DECODER (Peng et al. 2019); and Classical and Basal-like subtype classifiers from PurIST (Rashid et al. 2020). Programs from Bailey et al. (Bailey et al. 2016), periductal subtypes, and MSI subtypes were excluded from visualization due to redundancy with other included programs or low overall correlation across all factors.

## 10 Variance and Survival Contribution per Factor

For each factor  $j \in \{1, \dots, k\}$ , we quantified two quantities to assess the alignment between expression variance and prognostic relevance. The conditional variance explained by factor  $j$  was quantified as the semi-partial  $R^2$ :

$$V_j = \frac{\text{RSS}_{-j} - \text{RSS}_{\text{full}}}{\|X\|_F^2},$$

where  $\text{RSS}_{\text{full}} = \|X - WH\|_F^2$  is the residual sum of squares from the full  $k$ -factor reconstruction and  $\text{RSS}_{-j} = \|X - W_{-j}H_{-j}\|_F^2$  is the residual from the  $(k - 1)$ -factor reconstruction omitting factor  $j$ . This measures the incremental fraction of total variance accounted for by factor  $j$  given all other factors.

The survival contribution of factor  $j$  was quantified as the Type III partial likelihood increment: the reduction in Cox partial log-likelihood when factor  $j$  is omitted from the full  $k$ -factor model,

$$\Delta\ell_j = \ell(\beta_1, \dots, \beta_k) - \ell(\beta_1, \dots, \beta_{j-1}, \beta_{j+1}, \dots, \beta_k),$$

where  $\ell(\cdot)$  denotes the Cox partial log-likelihood evaluated at the maximum partial likelihood estimates. Larger  $\Delta\ell_j$  indicates a greater marginal contribution of factor  $j$  to survival prediction given all other factors. These two quantities are plotted against each other in main text, Fig. 3C.

## 11 Factor Correspondence Analysis

To assess which biological programs are preserved, altered, or suppressed under survival supervision, we quantified the pairwise correspondence between NMF and DeSurv factor loadings. For each pair of factors  $(j_{\text{NMF}}, j_{\text{DeSurv}})$ , we computed the Spearman rank correlation between the corresponding columns of the NMF gene program matrix  $W_{\text{NMF}}$  and the DeSurv gene program matrix  $W_{\text{DeSurv}}$ , restricted to the top 50 genes per factor (by factor-specificity score  $s_{ij}$ ; SI Appendix, Section~9) present in both models. Spearman correlation was used to capture monotone correspondence in gene loading ranks while remaining robust to scale differences between the two methods. Results are displayed as a heatmap with NMF factors as rows and DeSurv factors as columns (main text, Fig. 3D).

## Part II

# Supplementary Results

## 12 The DeSurv model shows consistent convergence and stability across restarts

Across datasets and initialization schemes, the DeSurv algorithm consistently converged to numerically stable solutions within the designated iteration budget. Warm-start strategies, in which solutions at  $\alpha = 0$  initialized supervised runs, substantially reduced variability across restarts and improved reproducibility. Fig. S1 shows representative loss trajectories demonstrating monotone decreases until convergence across restarts. Compared with naive random initialization, warm-starts produced tighter distributions of cross-validated C-index and partial likelihood, confirming stability of the optimization procedure.




Figure S1: Convergence trajectories of the DeSurv objective across five randomly selected multi-start initializations on the PDAC training cohort (TCGA + CPTAC). Each line shows the relative loss (loss at iteration  $t$  divided by the initial loss for that restart) as a function of iteration number, restricted to the first 5,000 iterations for readability. The total objective decreases monotonically and stabilizes well within the iteration budget across all initializations, confirming convergence of the alternating block-update scheme.

## 13 Simulation results under null and mixed scenarios

To characterize the operating characteristics of DeSurv across different signal regimes, we evaluated performance under three simulation scenarios. In the primary scenario (main text Fig. 3), prognostic programs explained low variance relative to outcome-neutral background signals, and survival depended on 150 factor-specific marker genes ( $\beta_1 = 2$ ,  $\beta_2 = \beta_3 = 0$ ). In the null scenario, survival times were generated independently of gene expression ( $\beta = 0$ ), so no prognostic structure existed. In the mixed scenario, survival depended on 300 genes per lethal factor, of which 50% were factor-specific markers and 50% were background genes with loadings shared across all factors. All scenarios used  $G = 3,000$  genes,  $N = 200$  samples,  $K = 3$  true factors, and identical gamma distribution parameters for the gene loading matrix  $W$  (see Materials

and Methods). Performance was evaluated using cross-validated concordance index (C-index), precision of recovered prognostic gene sets, and the distribution of selected ranks across simulation replicates.

Under the null scenario (Fig. S2A–B), DeSurv showed no advantage over unsupervised NMF ( $\alpha = 0$ ). Both methods yielded C-index values near 0.5, consistent with the absence of survival signal. The distribution of selected ranks was concentrated at low values for both methods, with NMF more strongly favoring  $k = 2$ . These results confirm that DeSurv does not hallucinate prognostic structure when none exists: Bayesian optimization selected low values of the supervision parameter  $\alpha$ , effectively recovering standard NMF as a special case. This specificity control is important because it establishes that the advantages observed in the primary scenario reflect genuine signal recovery rather than overfitting to noise in the survival labels.

Under the mixed scenario (Fig. S2C–F), four mechanistic observations characterize DeSurv’s advantage. First, when precision is evaluated against the factor-specific marker genes alone (150 genes; Fig. S2D left), DeSurv substantially outperforms NMF—matching or exceeding primary-scenario levels—even though overall survival-gene precision (300-gene survival set including background genes; Fig. S2D right) is more modestly improved. This dissociation reveals that background-gene recovery is similar between methods, and DeSurv’s edge comes specifically from concentrating marker programs. Second, the learned factor most aligned with the true marker program receives a substantially larger Cox coefficient ( $|\beta|$ ) under DeSurv than under NMF (Fig. S2E), confirming that survival supervision preferentially amplifies the marker-specific signal. Third, C-index values are modestly higher for DeSurv (Fig. S2C), consistent with the moderate overall improvement when background-gene dilution attenuates the survival signal. Together, these results demonstrate that DeSurv’s advantage in the mixed scenario is mechanistic: the method specifically recovers marker programs and assigns them elevated survival weights, rather than incidental capture of background-gene prognosis.

Together, the three scenarios provide a dose-response relationship: DeSurv’s advantage is greatest when variance and prognosis diverge (primary), moderate when they partially overlap (mixed), and absent when no survival signal exists (null). The mechanistic basis for the attenuation in the mixed scenario is now clear: DeSurv continues to specifically recover marker programs (high marker precision, elevated  $|\beta|$ ), but the overall survival-gene precision improvement is dampened because background-gene dilution in the 300-gene survival signal reduces the discrimination advantage. This gradient is consistent with the predictions of sufficient dimension reduction theory and offers practical guidance for users: DeSurv is most beneficial in settings where the dominant transcriptional axes are expected to be prognostically neutral, as is common in cancers with high tumor purity variation or dominant tissue-composition signals.

## 14 Cross-validated C-index across factorization rank

To evaluate how the number of latent factors  $k$  affects prognostic performance, we performed an exhaustive grid search over  $k \in \{2, 3, \dots, 12\}$  and supervision strength  $\alpha \in \{0, 0.05, \dots, 0.95\}$  using 5-fold cross-validation on the TCGA+CPTAC training cohort. For each  $k$ , the best-performing  $\alpha$  was selected by maximizing the mean CV C-index. The unsupervised NMF baseline ( $\alpha = 0$ ) was evaluated at the same  $k$  values for comparison. Results are shown for two gene-selection settings:  $n_{\text{top}} = 270$  top genes per factor and  $n_{\text{top}} = \text{ALL}$  (all genes retained), both at  $\lambda = 0.349$  and  $\xi = 0.056$ .

Fig. S3 shows the resulting C-index curves for both training (CV) and external validation (pooled across held-out cohorts), with shaded ribbons indicating  $\pm 1$  standard error. In training, DeSurv consistently outperforms NMF across all  $k$ , confirming that survival supervision improves in-sample concordance. In external validation, DeSurv at  $k = 3$  achieves concordance comparable to NMF at higher ranks (e.g.,  $k = 7$ ; Table S2), demonstrating that survival supervision concentrates prognostic information into fewer factors rather than requiring additional rank to capture it.

## 15 Standard NMF rank selection heuristics

Standard unsupervised rank selection criteria (reconstruction residuals, cophenetic correlation coefficient, and mean silhouette width) yielded inconsistent guidance for selecting the factorization rank  $k$  in the PDAC

training cohort (TCGA + CPTAC). Reconstruction residuals decreased smoothly without a clear elbow (Fig. S4A). Cophenetic correlation fluctuated without a distinct transition point (Fig. S4B). Mean silhouette width favored very low ranks that conflicted with other criteria (Fig. S4C). These contradictions motivated the survival-supervised rank selection approach described in the main text.

## 16 Cutpoint selection and validation of dichotomized risk groups

To convert the continuous DeSurv linear predictor into a clinically interpretable binary risk stratification, we selected an optimal z-score cutpoint via cross-validation on the TCGA+CPTAC training cohort. For each candidate cutpoint in  $\{-2.0, -1.8, \dots, 2.0\}$ , we computed the mean absolute log-rank z-statistic across held-out folds and selected the cutpoint maximizing this metric (Fig. S5A). We then applied this cutpoint to external validation cohorts by standardizing each patient’s linear predictor using the training mean and standard deviation. Fig. S5B shows the resulting Kaplan–Meier curves for the pooled external validation analysis, with a stratified log-rank test accounting for dataset of origin. Panels C–F display per-cohort validation KM curves for Dijk, Moffitt, PACA-AU, and Puleo, respectively. Across all cohorts, the DeSurv-derived high-risk group showed consistently shorter survival, confirming that the training-derived cutpoint generalizes to independent datasets.

## 17 Overlap of DeSurv risk groups with known molecular subtypes

To assess whether the DeSurv-derived risk stratification captures biologically meaningful transcriptional programs, we examined the overlap between dichotomized risk groups (High vs Low) and two established PDAC molecular classifiers: PurIST (Basal-like vs Classical) and DeCAF (proCAF vs restCAF). Fig. S6 shows the composition of each risk group across all pooled external validation cohorts, with Fisher’s exact test P-values quantifying the association. The DeSurv high-risk group was significantly enriched for both Basal-like (PurIST) and proCAF (DeCAF) subtypes, consistent with the known poor prognosis of these molecular classes. By contrast, standard NMF risk groups at the matched rank ( $k = 3$ ) separated Basal-like from Classical tumors (PurIST) but showed no significant enrichment for proCAF versus restCAF (DeCAF), indicating that survival-unsupervised factorization does not recover the tumor–stroma coupling captured by DeSurv’s D1. This concordance confirms that DeSurv’s survival-driven factorization recovers clinically relevant biology without requiring subtype labels during training.

## 18 Standard NMF factor structure at independently selected rank

Throughout the main text, both DeSurv and standard NMF were fit at the same rank ( $k = 3$ ), enabling a controlled comparison of the effect of survival supervision on factor structure. When each method selects its own rank independently, standard NMF chooses  $k = 5$  via residual elbow detection or  $k = 7$  via Bayesian optimization with  $\alpha = 0$ , while supervised DeSurv BO selects  $k = 3$ . We examine here the gene program structure at elbow-selected  $k = 5$  for both DeSurv (Fig. S7) and standard NMF (Fig. S8), and the NMF  $k = 7$  structure to illustrate additional fragmentation when supervision is absent (Fig. S9).

At elbow-selected  $k = 5$ , supervised and unsupervised factorizations yield substantially different gene-program structures. DeSurv at  $k = 5$  (Fig. S7) splits the ECM/activated-stroma (proCAF-associated) signal into two factors with opposing tumor correlations (one positively correlated with Classical tumor programs, one negatively correlated) and isolates restCAF as a separate factor; exocrine and Basal-like programs are not represented as dominant factors in the supervised solution. Standard NMF at  $k = 5$  (Fig. S8), by contrast, recovers the three core programs from its  $k = 3$  solution and adds two factors capturing Exocrine-like and iCAF variance, transcriptional variance not aligned with survival outcomes. These biologically redundant NMF factors explain why supervised BO selects  $k = 3$  in the production model: the additional NMF factors

at  $k = 5$  contribute no independent prognostic content beyond the three core programs (Fig. S10), even though the overall higher-rank model still validates externally on per-cohort C-index (Table S2).

At BO-selected  $k = 7$  under  $\alpha = 0$  (Fig. S9), standard NMF shows increased fragmentation relative to  $k = 5$ : multiple factors partially overlap the same PDAC gene signatures, and no new biologically coherent program emerges beyond those present at  $k = 3$ .

Table S2: External validation C-index by method and cohort. DeSurv at BO-selected rank ( $k = 3$ ) is compared with standard NMF at matched rank ( $k = 3$ ), elbow-selected rank ( $k = 5$ ), and BO-selected rank ( $k = 7$ ,  $\alpha = 0$ ). Higher values indicate better discrimination.

| Cohort            | DeSurv<br>(K=3) | Std NMF<br>(K=3) | Std NMF (K=5,<br>elbow) | Std NMF (K=7, BO $\alpha=0$ ) |
|-------------------|-----------------|------------------|-------------------------|-------------------------------|
| Dijk              | 0.607           | 0.542            | 0.633                   | 0.631                         |
| Moffitt_GEO_array | 0.562           | 0.521            | 0.552                   | 0.578                         |
| PACA_AU_array     | 0.649           | 0.470            | 0.657                   | 0.669                         |
| PACA_AU_seq       | 0.634           | 0.430            | 0.679                   | 0.702                         |
| Puleo_array       | 0.636           | 0.534            | 0.642                   | 0.651                         |

Standard NMF at elbow-selected  $k = 5$  and BO-selected  $k = 7$  both substantially improve over NMF at  $k = 3$  across all cohorts (Table S2), confirming that additional unsupervised factors capture prognostically relevant variation missed at the matched rank. At  $k = 7$ , NMF matches or exceeds DeSurv’s per-cohort C-index in most cohorts, demonstrating that unsupervised factorization can approach supervised concordance given enough factors. However, this comes at the cost of a less parsimonious and less interpretable factorization: the  $k = 7$  NMF solution shows increased fragmentation (Fig. S9). DeSurv’s advantage thus combines biological concordance and methodological complementarity: its risk groups identify the same poor-prognosis subtypes as established classifiers (Basal-like, proCAF; Fig. S6), while its linear predictor retains significant prognostic value after adjustment for those classifiers (Table S3).

Table S3 reports pooled validation hazard ratios (per SD of the linear predictor) for DeSurv at  $k = 3$  and the BO-selected NMF model at  $k = 7$  ( $\alpha = 0$ ), both unadjusted and after adjustment for PurIST and DeCAF molecular classifiers. Both methods retain significant prognostic value after adjustment, with DeSurv’s larger HR drop reflecting biological concordance with the established subtypes (per Fig. S6) and NMF  $k = 7$ ’s smaller drop reflecting residual content from its additional, less interpretable factors. The NMF  $k = 7$  model selects its rank via DeSurv’s BO infrastructure (with  $\alpha = 0$ , no factorization-level supervision); DeSurv uses survival at both levels, concentrating the prognostic signal into three factors rather than seven.

Table S3: Pooled external validation hazard ratios (per SD of linear predictor) for DeSurv at  $k = 3$  and standard NMF at BO-selected  $k = 7$  ( $\alpha = 0$ ). Adjusted models include PurIST and DeCAF molecular classifiers as covariates, with stratification by dataset.

| Method                                | Unadjusted HR per SD (95% CI), $P$ | Adjusted HR per SD (95% CI), $P$ |
|---------------------------------------|------------------------------------|----------------------------------|
| DeSurv ( $k = 3$ )                    | 1.50 (1.31–1.72), $< 0.001$        | 1.33 (1.13–1.56), $< 0.001$      |
| Std NMF ( $k = 7$ , BO $\alpha = 0$ ) | 1.48 (1.34–1.64), $< 0.001$        | 1.47 (1.27–1.70), $< 0.001$      |

The variance–survival decomposition at  $k = 5$  (Fig. S10) confirms that elbow selection does not recover additional prognostic structure. At  $k = 5$ , standard NMF continues to distribute survival contribution negligibly across all factors; the two factors new relative to  $k = 3$  (N4 and N5) contribute minimally, confirming that the additional rank captures background transcriptional variance rather than independent prognostic programs.

## 19 Factor-Specific Gene Lists

Tables S4–S6 list the top 270 genes for each of the three DeSurv factors, ranked by the factor specificity score  $s_{ij}$  defined in SI Appendix, Section 9. The 270-gene cutoff corresponds to the Bayesian optimization–selected projection dimension ( $n_{\text{top}} = 270$ ); these are the genes used to compute sample-level factor scores for survival prediction. Factor D1 is enriched for Classical tumor and restCAF stromal genes, Factor D2 for proCAF-associated stromal genes, and Factor D3 for Basal-like tumor and proliferation-associated genes.

Table S4: Top 270 genes for DeSurv factor D1 (Factor D1 (Classical tumor + restCAF stroma)), ranked by factor specificity score  $s_{ij}$ . The 270-gene cutoff corresponds to the Bayesian optimization–selected projection dimension ( $n_{\text{top}}$ ).

| Factor D1 (Classical tumor + restCAF stroma) |            |          |         |          |          |
|----------------------------------------------|------------|----------|---------|----------|----------|
| 1–45                                         | 46–90      | 91–135   | 136–180 | 181–225  | 226–270  |
| DMBT1                                        | FGFR2      | GOLGA8A  | PTPRN2  | RPL9     | PPP1R12B |
| IGFBP2                                       | COL9A2     | IL2RG    | FCGBP   | CD79A    | PIP5K1B  |
| S100A4                                       | COL22A1    | KIAA1211 | PLA2G2A | PROM1    | AHCYL2   |
| KRT23                                        | IRF7       | SFRP1    | FABP4   | SCD      | SST      |
| LIF                                          | SBNO2      | ADH1B    | EPPK1   | TMC5     | CYP3A5   |
| IFI6                                         | MGAT3      | MFSD2A   | CPXM2   | CCL14    | SMAP2    |
| PDLIM3                                       | SYNM       | CLU      | KIF1A   | PLXNB1   | ZG16B    |
| CLDN18                                       | APOD       | CLMN     | NGFR    | TMEM119  | SERPINF1 |
| MYO15B                                       | SEMA6A     | CHPF     | NFKBIA  | SORBS2   | CLDN3    |
| SYT8                                         | TNXB       | MTMR11   | SPOCK2  | C7       | NR1I2    |
| TFF2                                         | PTGIS      | TTR      | ATP2C2  | SLC12A7  | JUND     |
| TMPRSS2                                      | CCL18      | WFDC2    | NBEAL2  | QSOX1    | NCF2     |
| CHGB                                         | ZC3H12A    | C2orf72  | CAPN9   | UNC5CL   | CELSR1   |
| CST1                                         | CELSR3     | CCND2    | PTGDS   | LONRF2   | MUC17    |
| B3GNT7                                       | EMB        | ARHGAP4  | LSP1    | VSIG2    | INPP5D   |
| F5                                           | KCNJ15     | PKDCC    | LAMA5   | ADAMTSL2 | TM4SF5   |
| TNRC18                                       | SORBS1     | SRPX     | FASN    | ARHGEF16 | HNFA4    |
| IGF2                                         | PAQR8      | CCL21    | GFRA1   | PCOLCE   | CFD      |
| ARSD                                         | SLC4A4     | CARD11   | PTPRH   | CFC1B    | HCLS1    |
| ZDHHC7                                       | ACSS1      | CTRB2    | FMOD    | SLC9A3   | SH3PXD2B |
| LENG8                                        | ST6GALNAC1 | GOLGA8B  | FHL1    | ACACB    | PCSK2    |
| ATP9A                                        | DES        | IYD      | PCDH20  | BCAS1    | HMGCS2   |
| TMPRSS3                                      | PRUNE2     | CORO1A   | CCR7    | RASD1    | FILIP1L  |
| CES1                                         | PLCB4      | SAA2     | CHIT1   | LGALS4   | ETNK1    |
| HOPX                                         | CADPS      | CDCA7    | PARM1   | MS4A1    | TNNI2    |
| CNN1                                         | FMO5       | BOC      | ITM2A   | CAPN6    | TOX3     |
| MIA                                          | RAP1GAP2   | LTB      | NAPSB   | SCCPDH   | FRZB     |
| CRIP2                                        | GALNT12    | COMP     | AGT     | PTPRCAP  | HABP2    |
| NEURL1B                                      | CCL2       | MANSC1   | FUT3    | VNN1     | GDA      |
| CHGA                                         | REG3A      | SLC19A3  | SIPA1L2 | PI16     | NCOR2    |
| ACTG2                                        | ZCCHC24    | STEAP2   | SPINK5  | SEMA7A   | DNAH2    |
| CXCL14                                       | LTF        | FOXA1    | ACSL1   | IL10RA   | DOK4     |
| TPPP3                                        | EFHD2      | WNK2     | BTNL8   | GALNT3   | HOXB9    |
| PIGR                                         | ATP11A     | SELL     | CORO2A  | GOLM1    | KIAA1324 |
| CYP27A1                                      | GP2        | PLEKHA6  | LAD1    | TNFAIP2  | TRIM2    |
| ADAM8                                        | HIPK2      | NFASC    | TMEM63A | CD79B    | FGFBP1   |
| FGFR3                                        | SLC26A9    | EGR3     | GC      | ENPP2    | IRX3     |
| IGF1R                                        | TACC2      | PLIN4    | C1orf21 | SQLE     | TINAGL1  |
| PLEKHB1                                      | TRAK1      | WDR72    | MIF     | KLF4     | SFRP4    |
| CPXM1                                        | SAA1       | AMY2A    | FOXA2   | PDLIM7   | APLNR    |
| ATP7B                                        | CLDN23     | IL16     | CCL19   | FOSB     | AGRN     |
| SLC12A2                                      | FOXJ1      | TJP3     | GSTA1   | RHPN2    | SLC25A25 |
| CCL20                                        | FRY        | SLC9A2   | LLGL2   | SCG2     | PODN     |
| AKNA                                         | GATA6      | LMOD1    | ACP5    | EMILIN1  | ITIH2    |
| COL16A1                                      | PLXNA2     | CPS1     | UGT2B15 | BACE2    | FLJ23867 |

Table S5: Top 270 genes for DeSurv factor D2 (Factor D2 (proCAF stroma)), ranked by factor specificity score  $s_{ij}$ .

| Factor D2 (proCAF stroma) |         |         |         |         |           |
|---------------------------|---------|---------|---------|---------|-----------|
| 1–45                      | 46–90   | 91–135  | 136–180 | 181–225 | 226–270   |
| UHMK1                     | RPS26   | SAMD9L  | ATP8A1  | IRF6    | RASSF2    |
| DDR2                      | MYLK    | DOCK2   | TGFBR1  | KLHL6   | ZNF83     |
| INHBA                     | COL11A1 | IGF1    | SYNPO2  | C1QTNF3 | CYTIP     |
| DYNC2H1                   | NFIB    | TACC1   | PDE4B   | PCLO    | SRPX2     |
| ZDHHC20                   | PTPN13  | CFH     | SGCD    | ITGA2   | LY75      |
| ITPR2                     | PDE3A   | DMD     | STK17B  | LRIG3   | HLA-DOA   |
| HMCN1                     | TRIM22  | PCDH7   | NUAK1   | SSPN    | STK38L    |
| LAMA2                     | DPYD    | MALAT1  | FCGR3A  | F2R     | RSAD2     |
| SLFN5                     | COL14A1 | NOTCH2  | GPNNB   | PDGFC   | HBB       |
| PLXNC1                    | CNTN1   | CDK6    | VCAM1   | CD109   | HERC2P2   |
| SVEP1                     | NCKAP1L | FBN1    | ADAMTS9 | CLIC4   | AOX1      |
| CYBB                      | PHLDB2  | SYNE2   | ITGB8   | SORL1   | PLIN2     |
| SYNE1                     | ITGA4   | RHOBTB3 | CR1     | AOAH    | PFKFB2    |
| SAMHD1                    | MACC1   | CALCRL  | SEPT11  | FMO2    | OGN       |
| SKIL                      | SEMA5A  | NRP1    | IFI44L  | RDX     | DOCK10    |
| ADAMTS12                  | MRC1    | LIFR    | ELL2    | IL1RAP  | RGS4      |
| BICC1                     | SLIT2   | STAB1   | GPR183  | GBP1    | AXL       |
| ITGA1                     | ZEB2    | FLRT2   | ZBTB16  | CXCR4   | FGD6      |
| COL8A1                    | ADAM10  | RNF144A | FRMD6   | WNK1    | CDC42BPA  |
| PDGFRA                    | SAMD9   | BCAT1   | PDE5A   | MFAP5   | SLC6A6    |
| ABCA1                     | MS4A7   | CILP    | CSF1R   | ALDH1A3 | RGS1      |
| TCF4                      | EFEMP1  | CCDC80  | PLEK    | CD53    | KIAA1324L |
| CEP170                    | BIRC3   | FAT4    | SLK     | OMD     | CPA3      |
| CD163                     | WISP1   | BNC2    | SLC7A2  | PLOD2   | CHL1      |
| MAN1A1                    | LAMA4   | LTBP1   | HEG1    | ITGAV   | SLC7A11   |
| MSR1                      | FAM198B | COL12A1 | GREM1   | MACF1   | GFPT2     |
| CCDC88A                   | PTPRC   | IL1R1   | VGLL3   | NID2    | RAI14     |
| F13A1                     | FAP     | SLA     | GBP5    | NAMPT   | LMAN1     |
| RPS28                     | ARRDC3  | MBOAT2  | SLC1A3  | ADAM28  | MECOM     |
| RALGAPA2                  | MYH10   | FNDC3B  | FGF7    | GNE     | FPR1      |
| ZEB1                      | FCGR2A  | PRRX1   | ABCA8   | ENAH    | EDEM3     |
| FBXO32                    | PARP14  | COL10A1 | ADAM12  | FAM129A | RASSF6    |
| INPP4B                    | DSE     | DDX60   | CYBRD1  | ECM2    | IQGAP2    |
| CDH11                     | EHF     | DOCK9   | TFRC    | VSIG4   | MIAT      |
| PKHD1                     | OSMR    | SRGN    | CTSK    | MTAP    | PRICKLE1  |
| AKAP2                     | ARID5B  | FOXO1   | COLEC12 | INMT    | CHST15    |
| CYP1B1                    | ASPN    | MS4A6A  | ALDH1L2 | SGMS2   | GEM       |
| CCL5                      | SLCO2B1 | ADAMTS2 | PGM2L1  | COL4A4  | CHRD1     |
| FGL2                      | FPR3    | CPVL    | FAM135A | CAPRIN2 | DSC2      |
| MAP1B                     | DOCK8   | CD93    | DOCK11  | CENPF   | RAB31     |
| PLXDC2                    | SLFN11  | DST     | EPHA3   | DCLK1   | SLC38A1   |
| LRRK2                     | ZNF532  | CDK14   | PAPPA   | EDNRA   | CNN3      |
| KIAA0754                  | SPON1   | MSRB3   | EGFR    | WNT5A   | FLT1      |
| EDIL3                     | SEMA3C  | RNF213  | RASA1   | GALNT5  | TMEM200A  |
| ITGBL1                    | FERMT2  | SEL1L   | AHR     | SLC38A2 | SLC2A3    |



Table S6: Top 270 genes for DeSurv factor D3 (Factor D3 (Basal-like tumor)), ranked by factor specificity score  $s_{ij}$ .

| Factor D3 (Basal-like tumor) |          |           |          |          |          |
|------------------------------|----------|-----------|----------|----------|----------|
| 1–45                         | 46–90    | 91–135    | 136–180  | 181–225  | 226–270  |
| HSPB1                        | HLA-H    | LAMB3     | MYL9     | IL1RN    | SDC1     |
| KRT7                         | HSPA1A   | FSCN1     | COL17A1  | TST      | KLK11    |
| MSLN                         | LGALS3   | TSPAN13   | EPS8L1   | RAPGEFL1 | PERP     |
| RPSAP58                      | FAM83H   | TSPAN8    | CDH3     | C15orf48 | ARHGAP23 |
| S100P                        | EPB41L1  | ALDH3B1   | RNASET2  | PFKP     | HPGD     |
| RPS16                        | EFNA1    | ANPEP     | TUBA1C   | STARD10  | AKR1C3   |
| IL32                         | TKT      | GSTP1     | RAP1GAP  | FOXQ1    | CRABP2   |
| SH3BGRL3                     | SLC2A1   | KLK10     | H19      | REG4     | GRB7     |
| IFITM2                       | PLOD1    | B4GALT1   | LTBP3    | GJB1     | SERPINA4 |
| RPS21                        | IER3     | GAS6      | AQP3     | SLC29A1  | SLC39A4  |
| SEMA4B                       | B3GNT3   | ABCC3     | MPZL2    | NOTCH3   | FA2H     |
| APOL1                        | PRSS3    | CLDN2     | ID1      | TRIM16   | RPL8     |
| S100A16                      | TOB1     | CEACAM6   | RARRES2  | TAP2     | MT1X     |
| DNAJB1                       | EGR1     | CREB3L1   | SLC25A6  | KRT19    | TCN1     |
| MT2A                         | PLA2G16  | MST1R     | RBP1     | CRP      | AKAP1    |
| TNFRSF21                     | OCIAD2   | CEBPB     | HSPA5    | SNHG5    | GJB3     |
| RPL28                        | LPCAT4   | TXN       | PER1     | KRT18    | DUOX2    |
| FXSD3                        | BST2     | NR4A1     | ADAP1    | CA12     | HLA-DRB5 |
| SERPINC1                     | EPHA2    | AP1M2     | TMC4     | MYEOV    | ARRDC2   |
| HMGA1                        | MUC5B    | SYNPO     | HSD17B2  | P4HB     | OAS1     |
| S100A14                      | KCNN4    | ANO1      | PTP4A1   | VILL     | ADRA2A   |
| HIF0                         | ITGA3    | SIK1      | PLAUR    | HKDC1    | NQO1     |
| KRT17                        | TSTA3    | EFNB2     | ANXA10   | STEAP3   | LRG1     |
| CSTB                         | IFI27    | CDKN1A    | PYGB     | PLEK2    | SERPINH1 |
| ASS1                         | MMP1     | S100A10   | CYP2S1   | CD68     | COL18A1  |
| EFNB1                        | SLPI     | PKP3      | ALDOA    | MAFF     | CERCAM   |
| MDK                          | CLTB     | MCM7      | CDCP1    | CD9      | CA9      |
| LY6E                         | SPINK1   | RARRES3   | IFITM3   | PLLP     | CDHR2    |
| CAMK2N1                      | JUP      | CEACAM1   | BAIAP2L1 | EPHB4    | ESRP1    |
| MGLL                         | PPP1R15A | ALDH2     | MGST1    | KRT16    | CRYZ     |
| PDZK1IP1                     | TRIM29   | CES2      | TFF1     | ABLIM3   | TAGLN2   |
| BCL2L1                       | PPP1R1B  | SLC6A8    | MUC4     | PTPRU    | HNF1B    |
| C19orf33                     | ARPC1A   | PPL       | MMP28    | MUC6     | BMP4     |
| SFN                          | GPRC5A   | NDRG1     | CLDN1    | FDFT1    | MT1E     |
| ITGB4                        | RAB25    | AZGP1     | TM4SF1   | GAPDH    | ANGPTL4  |
| TSPAN1                       | ZFP36L2  | S100A11   | TFF3     | EPS8L3   | CFB      |
| SLC16A3                      | CLDN7    | SEMA3B    | UNC13D   | CLDN10   | GDF15    |
| C4A                          | DDIT4    | TNFRSF12A | SERINC2  | ITPKC    | PADI1    |
| KLF5                         | FKBP4    | SOX9      | SPNS2    | LOXL2    | TMSB10   |
| IER2                         | LAPTM4B  | PMEPA1    | AQP5     | AMIGO2   | B2M      |
| ZFP36L1                      | MUC13    | CYR61     | IGFBP4   | DUOX2    | CAV2     |
| MAL2                         | PSCA     | EPCAM     | SLC7A5   | CLDN4    | CTSH     |
| SCNN1A                       | SDCBP2   | BHLHE40   | TESC     | CAPG     | PABPC1   |
| MALL                         | PROM2    | CX3CL1    | IFITM1   | CEACAM5  | KRT8     |
| BGN                          | LRRC8A   | SMAD3     | KCNK5    | VWA1     | TAP1     |

## SI References

- Bailey, Peter, David K Chang, Katia Nones, et al. 2016. “Genomic Analyses Identify Molecular Subtypes of Pancreatic Cancer.” *Nature* 531 (7592): 47–52. <https://doi.org/10.1038/nature16965>.
- Blei, David M, and Jon D McAuliffe. 2007. “Supervised Topic Models.” In *Advances in Neural Information Processing Systems 20 (NIPS 2007)*, edited by J. C. Platt, D. Koller, Y. Singer, and S. T. Roweis. Curran Associates, Inc.

- Brunet, Jean-Philippe, Pablo Tamayo, Todd R Golub, and Jill P Mesirov. 2004. “Metagenes and Molecular Pattern Discovery Using Matrix Factorization.” *Proceedings of the National Academy of Sciences USA* 101 (12): 4164–69. <https://doi.org/10.1073/pnas.0308531101>.
- Cai, Deng, Xiaofei He, Jiawei Han, and Thomas S. Huang. 2011. “Graph Regularized Nonnegative Matrix Factorization for Data Representation.” *IEEE Transactions on Pattern Analysis and Machine Intelligence* 33 (8): 1548–60. <https://doi.org/10.1109/TPAMI.2010.231>.
- Dijk, Frederike, Veronique L Veenstra, Eline C Soer, et al. 2020. “Unsupervised Class Discovery in Pancreatic Ductal Adenocarcinoma Reveals Cell-Intrinsic Mesenchymal Features and High Concordance Between Existing Classification Systems.” *Scientific Reports* 10 (1): 337. <https://doi.org/10.1038/s41598-019-56826-9>.
- Donoho, David, and Victoria Stodden. 2004. “When Does Non-Negative Matrix Factorization Give a Correct Decomposition into Parts?” In *Advances in Neural Information Processing Systems 16 (NIPS 2003)*, edited by Sebastian Thrun, Lawrence K. Saul, and Bernhard Schölkopf. MIT Press.
- Ellis, MJ, M Gillette, SA Carr, et al. 2013. “Connecting Genomic Alterations to Cancer Biology with Proteomics: The NCI Clinical Proteomic Tumor Analysis Consortium.” *Cancer Discovery* 3 (10): 1108–12. <https://doi.org/10.1158/2159-8290.CD-13-0219>.
- Elyada, Ela, Mohan Bolisetty, Pasquale Laise, et al. 2019. “Cross-Species Single-Cell Analysis of Pancreatic Ductal Adenocarcinoma Reveals Antigen-Presenting Cancer-Associated Fibroblasts.” *Cancer Discovery* 9 (8): 1102–23. <https://doi.org/10.1158/2159-8290.CD-19-0094>.
- Gillis, Nicolas. 2014. “The Why and How of Nonnegative Matrix Factorization.” In *Regularization, Optimization, Kernels, and Support Vector Machines*, edited by Johan A. K. Suykens, Marco Signoretto, and Andreas Argyriou. Chapman; Hall/CRC.
- Gillis, Nicolas, and François Glineur. 2012. “Accelerated Multiplicative Updates and Hierarchical ALS Algorithms for Nonnegative Matrix Factorization.” *Neural Computation* 24 (4): 1085–105. [https://doi.org/10.1162/NECO\\_a\\_00256](https://doi.org/10.1162/NECO_a_00256).
- Grippo, Luigi, and Marco Sciandrone. 2000. “On the Convergence of the Block Nonlinear Gauss–Seidel Method Under Convex Constraints.” *Operations Research Letters* 26 (3): 127–36. [https://doi.org/10.1016/S0167-6377\(99\)00074-7](https://doi.org/10.1016/S0167-6377(99)00074-7).
- Kim, Jingu, and Haesun Park. 2008. *Sparse Nonnegative Matrix Factorization for Clustering*. GT-CSE-08-01. College of Computing, Georgia Institute of Technology. <https://faculty.cc.gatech.edu/~hpark/papers/GT-CSE-08-01.pdf>.
- Laurberg, Hans, Mads Græsbøll Christensen, Mark D Plumbley, Lars Kai Hansen, and Søren Holdt Jensen. 2008. “Theorems on Positive Data: On the Uniqueness of NMF.” *Computational Intelligence and Neuroscience* 2008: 764206. <https://doi.org/10.1155/2008/764206>.
- Leary, Jack R, Yi Xu, Ashley B Morrison, et al. 2023. “Sub-Cluster Identification Through Semi-Supervised Optimization of Rare-Cell Silhouettes (SCISSORS) in Single-Cell RNA-Sequencing.” *Bioinformatics* 39 (8): btad449. <https://doi.org/10.1093/bioinformatics/btad449>.
- Lee, Daniel D., and H. Sebastian Seung. 2001. “Algorithms for Non-Negative Matrix Factorization.” In *Advances in Neural Information Processing Systems 13 (NIPS 2000)*, edited by Todd K. Leen, Thomas G. Dietterich, and Volker Tresp. MIT Press.
- Lin, Chih-Jen. 2007. “On the Convergence of Multiplicative Update Algorithms for Nonnegative Matrix

- Factorization.” *IEEE Transactions on Neural Networks* 18 (6): 1589–96. <https://doi.org/10.1109/TNN.2007.895831>.
- Moffitt, Richard A, Raoud Marayati, Elizabeth L Flate, et al. 2015. “Virtual Microdissection Identifies Distinct Tumor- and Stroma-Specific Subtypes of Pancreatic Ductal Adenocarcinoma.” *Nature Genetics* 47 (10): 1168–78. <https://doi.org/10.1038/ng.3398>.
- Pascual-Montano, Alberto, Jose M. Carazo, Kenfei Kochi, Daniel Lehmann, and Roberto D. Pascual-Marqui. 2006. “Nonsmooth Nonnegative Matrix Factorization (nsNMF).” *IEEE Transactions on Pattern Analysis and Machine Intelligence* 28 (3): 403–15. <https://doi.org/10.1109/TPAMI.2006.60>.
- Peng, Xianlu Laura, Ian C McCabe, Elena V Kharitonova, et al. 2026. “DeCAF Defines Clinical Fibroblast Subtypes and Multidimensional Tumor-Stroma Crosstalk Shaping Prognosis and Immunotherapy Response.” *Cell Reports Medicine* 7 (2): 102611. <https://doi.org/10.1016/j.xcrm.2026.102611>.
- Peng, Xianlu Laura, Richard A Moffitt, Robert J Torphy, Keith E Volmar, and Jen Jen Yeh. 2019. “De Novo Compartment Deconvolution and Weight Estimation of Tumor Samples Using DECODER.” *Nature Communications* 10 (1): 4729. <https://doi.org/10.1038/s41467-019-12517-7>.
- Puleo, Francesco, Rémy Nicolle, Yuna Blum, et al. 2018. “Stratification of Pancreatic Ductal Adenocarcinomas Based on Tumor and Microenvironment Features.” *Gastroenterology* 155 (6): 1999–2013. <https://doi.org/10.1053/j.gastro.2018.08.033>.
- Raphael, Benjamin J, Ralph H Hruban, Andrew J Aguirre, et al. 2017. “Integrated Genomic Characterization of Pancreatic Ductal Adenocarcinoma.” *Cancer Cell* 32 (2): 185–203. <https://doi.org/10.1016/j.ccell.2017.07.007>.
- Rashid, Naim U, Xianlu L Peng, Chong Jin, et al. 2020. “Purity Independent Subtyping of Tumors (PurIST), a Clinically Robust, Single-Sample Classifier for Tumor Subtyping in Pancreatic Cancer.” *Clinical Cancer Research* 26 (1): 82–92. <https://doi.org/10.1158/1078-0432.CCR-19-1467>.
- Simon, Noah, Jerome H Friedman, Trevor Hastie, and Rob Tibshirani. 2011. “Regularization Paths for Cox’s Proportional Hazards Model via Coordinate Descent.” *Journal of Statistical Software* 39 (5): 1–13. <https://doi.org/10.18637/jss.v039.i05>.
- Tseng, Paul. 2001. “Convergence of a Block Coordinate Descent Method for Nondifferentiable Minimization.” *Journal of Optimization Theory and Applications* 109 (3): 475–94. <https://doi.org/10.1023/A:1017501703105>.
- Zhao, Lan, Hongya Zhao, and Hong Yan. 2018. “Gene Expression Profiling of 1200 Pancreatic Ductal Adenocarcinoma Reveals Novel Subtypes.” *BMC Cancer* 18 (1): 603. <https://doi.org/10.1186/s12885-018-4546-8>.




Figure S2: DeSurv performance under null and mixed simulation scenarios. (A–B) Null scenario ( $\beta = 0$ ): both methods yield C-index  $\approx 0.5$  (A) and comparable rank distributions (B), confirming DeSurv does not hallucinate prognostic structure; precision is omitted as no true survival genes exist. (C–F) Mixed scenario: survival depends on factor-specific marker genes (50%) and shared background genes (50%). (C) C-index is modestly higher for DeSurv. (D) Precision by gene-set definition: marker-only precision (150 genes; left facet) substantially favors DeSurv; full 300-gene survival-set precision (right facet) shows a more modest improvement. (E) The matched factor receives a substantially larger Cox coefficient  $|\beta|$  under DeSurv, confirming preferential amplification of marker-specific signal. (F) Selected-rank distributions. DeSurv (blue); NMF (red,  $\alpha = 0$ ); both Bayesian-optimized.




Figure S3: Training (CV) and external validation C-index as a function of factorization rank  $k$  for DeSurv (blue) and standard NMF (red,  $\alpha = 0$ ). Shaded ribbons show  $\pm 1$  SE. For each  $k$ , the best  $\alpha$  was selected by maximizing the mean CV C-index. Validation C-index is pooled across all external cohorts. (A) Results using  $n_{\text{top}} = 270$  top genes per factor ( $\lambda = 0.349, \xi = 0.056$ ). (B) Results retaining all genes ( $n_{\text{top}} = \text{ALL}; \lambda = 0.349, \xi = 0.056$ ).




Figure S4: Standard unsupervised NMF rank selection heuristics for the PDAC training cohort (TCGA + CPTAC). (A) Reconstruction residuals as a function of factorization rank  $k$ . (B) Cophenetic correlation coefficient as a function of  $k$ . (C) Mean silhouette width across multiple distance metrics as a function of  $k$ . These criteria yield inconsistent guidance, motivating the survival-supervised model selection approach (main text Fig. 2D).




Figure S5: Cutpoint selection and external validation of dichotomized risk groups. (A) Cross-validated mean absolute log-rank z-statistic as a function of z-score cutpoint; red dashed line indicates the selected optimum. (B) Pooled validation Kaplan–Meier curves (stratified log-rank test). (C–D) Per-cohort validation KM curves: (C) Dijk, (D) Moffitt. (E–F) Per-cohort validation KM curves: (E) PACA-AU, (F) Puleo.




Figure S6: Subtype composition of DeSurv (top) and standard NMF  $k = 3$  (bottom) risk groups in pooled external validation cohorts. Stacked bar charts show the proportion of PurIST (left) and DeCAF (right) subtypes within Low and High risk groups defined by the log-rank-optimized cutpoint. Numbers indicate sample counts; Fisher's exact test P-values are shown above each panel. DeSurv risk groups are enriched for both Basal-like (PurIST) and proCAF (DeCAF) poor-prognosis subtypes; NMF risk groups separate Basal-like from Classical tumors but show no significant CAF subtype enrichment.






Figure S7: Gene program overlap heatmap for DeSurv at elbow-selected rank  $k = 5$ . Each cell shows the number of top genes from a given factor (rows) overlapping established PDAC gene programs (columns). DeSurv at  $k = 5$  splits the ECM/activated-stroma (proCAF-associated) signal into two factors with opposing Classical-tumor correlations and isolates restCAF as a separate factor; exocrine and Basal-like programs are not represented as dominant factors.




Figure S8: Gene program overlap heatmap for standard NMF at elbow-selected rank  $k = 5$ . Each cell shows the number of top genes from a given factor (rows) overlapping established PDAC gene programs (columns). The three core programs from  $k = 3$  are recovered; two additional factors overlap Exocrine-like and iCAF signatures, capturing transcriptional variance not aligned with survival outcomes.




Figure S9: Gene program overlap heatmap for standard NMF at  $k = 7$  (BO-selected rank with  $\alpha = 0$ ). Compared with  $k = 3$  and  $k = 5$ , the 7-factor solution shows increased fragmentation: multiple factors partially capture the same PDAC gene signatures and no new coherent biological program emerges.




Figure S10: Fraction of expression variance explained versus Type III partial likelihood contribution for each standard NMF factor at elbow-selected  $k = 5$ . Each point represents one factor, labeled N1–N5. Survival contribution is quantified as the reduction in Cox partial log-likelihood when each factor is removed from the full model. The two factors new relative to  $k = 3$  (N4, N5) contribute negligibly to survival, confirming that elbow selection recovers no additional prognostic structure.