Mollecular and Cellular Biology

Lecture – 2

Department of CSE, DIU



CONTENTS

- 1. Cell
 - Eukaryotes VS Prokaryotes
- 2. Cell Division
 - Mitosis VS Meiosis
- 3. Nucleic Acids
 - 3.1. DeoxyriboNucleic Acid (DNA)
 - * DNA Structure
 - * DNA Replication
 - 3.2. RiboNucleic Acid (RNA)
 - * RNA Structure
 - * Major RNA Types

1. Cell


Let's learn about Eukaryotes, Prokaryotes and Cell Division

What is Life made of?


















Cells

- Fundamental working units of every living system.
- Cell specialization in multi-cellular organism.
- **Tissues** are groups of cells for a particular function.
 - Fourteen major tissue types
 - Bone, muscle, nerve etc.
- Organs are formed
- More than 200 different cell types
 - With lots of variety in every sense
 - But the genetic code is same












Blood Bone

Nerve

Muscle

Fat

Life begins with Cell




- A cell is the smallest structural unit of an organism that is capable of sustained independent functioning
- All cells have some common features

2 types of Cells

- 1. Eukaryotic Cells
- 2. Prokaryotic Cells


Eukaryotic Cells



- Single or Multi Cell
- Are called Eukaryotes
- Have Nucleus
- Have membrane bounded organelles
- Have chromosomes inside Nucleus
- Seen in most of the life forms

Prokaryotic Cells

- ▶ Single Cell organism
- ▶ Are called Prokaryotes
- ⊳No Nucleus
- ▷No other membrane bounded organelles
- ▷One piece of rolled up DNA floating in cellular fluid
- ▶ Mostly some forms of very ancient Bacteria




2. Cell Division

2 types of Cell Division

- Mitosis Cell Division
- Miosis Cell Division


Mitosis Cell Division



- Happens in Somatic Cells (general body cells)
- Cell and chromosome, both gets divided only once
- No cross over
- Takes part in healing and repair

Meiosis Cell Division

- Happens in Germ Cells
- Cell is divided twice, but chromosome only once
- Cross over happens
- Takes part in formation of gametes and maintenance of chromosome number in the race



-centrosome


3. Nucleic Acid

All Life depends on 3 critical molecules

- DNAs
 - Hold information on how cell works
- RNAs
 - Act to transfer short pieces of information to different parts of cell
 - Provide templates to synthesize into protein
- Proteins
 - Form enzymes that send signals to other cells and regulate gene activity
 - Form body's major components (e.g. hair, skin, etc.)
 - Are life's laborers!

Building Blocks of Nucleic acids

- DNA/RNA are polymeric chain on nucleotides
- Three parts of Nucleotides
 - a nitrogenous base,
 - a five-carbon-atom sugar and
 - a phosphate group




- Phosphate Molecule
- Deoxyribose Sugar
- Base

Adenine, Cytosine, Guanine and Thymine

Nucleic acids Bases

 Adenine (A), Guanine (G), Cytosine (C) and Thymine (T), Uracil (U)




3.1 DeoxyriboNucleic Acid (DNA)

Carrier of genetic instructions


DNA Structure

- Double Helix Structure (Watson and Crick, Nature 1953)
- ▶ Two complimentary antiparallel strands, one runs from 5' to 3' end and another runs from 3' to 5' end
- ▶3 major parts Nitrogenous Base, 5-Carbon Deoxyribose Sugar and Phosphate Group
- ▶ Four nitrogenous bases Adenine (A), Cytosine (C), Guanine (G), Thymine (T)
- ►A-T is Double Hydrogen Bond and G-C is Triple Hydrogen Bond
- DNA is more stable than RNA due to its Deoxyribose Sugar Structure



© 2007 Encyclopædia Britannica, Inc.

DNA Replication



▶Initiation

- Helicase enzyme unwinds DNA strands
- Replication fork is created
- RNA Primer is created by Primase enzyme
- Primer is starting point of elongation

▶ Elongation

- New DNA Strand grows 1 base at a time as complimentary of leading strand (5' to 3')
 - DNA Polymerase enzyme controls it
- Complimentary strand of lagging strand is created in small fragments called Okazaki Fragments (3' to 5')
- ▶ Termination
- Exonuclease enzyme removes all the primer sequences from new strands
 - Again, DNA Polymerase fills the gaps
 - DNA Ligase enzyme seals all the gaps


* DNA Replication is Semi-Conservative, because, in new sets of DNA, one strand is newly created but the other strand comes from the ancestor.

3.2 RiboNucleic Acid (RNA)

Protein Coding and Carrier

RNA Structure

- ▶ Single Helix Structure
- Single Strand which generally runs from 5' to 3'
- ▶3 major parts Nitrogenous Base, 5-Carbon Ribose Sugar and Phosphate Group
- ▶ Four nitrogenous bases Adenine (A), Cytosine (C), Guanine (G), Uracil (U)
- ►A-U is Double Hydrogen Bond and G-C is Triple Hydrogen Bond
- ▶RNA is less stable than DNA due to its Ribose Sugar's structure



RNA Types

Messenger RNA (mRNA)

Carries a genes coding message for protein from Nucleus to Ribosome

Transfer RNA (tRNA)

Transfers specific amino acid sequence to ribosome to form Protein

Ribosomal RNA (rRNA)

Protein and rRNA combinedly forms ribosome

Non-Coding RNA

Not translated into protein. Ex – tRNA, rRNA

Catalytic RNA

Catalyze chemical reaction.

Double Stranded RNA

Contains complementary strands like DNA. Induces gene expression.

